

Antidiarrheal activity of the combination of ethanol extract of Tumeric.

by Sumi Wijaya

FILE	7-ANTIDIARRHEAL_ACTIVITY_OF_THE.PDF (966K)	WORD COUNT	3066
TIME SUBMITTED	25-SEP-2020 03:16PM (UTC+0700)	CHARACTER COUNT	17077
SUBMISSION ID	1396635428		

Antidiarrheal Activity of The Combination of Ethanol Extracts of Turmeric Rhizomes, Gall oak Leaves, Guava Leaves and Meniran Herbs

Authors Sumi Wijaya^{*}, Lisa Soegianto

Afiliation Faculty of Pharmacy, Widya Mandala Catholic University, Surabaya, East Java, Indonesia

Keyword

- ☞ Antibacterial
- ☞ Antidiarrheal
- ☞ Intestinal transit
- ☞ Protection method

Received 7 September 2015

Revised 24 April 2017

Accepted 6 November 2017

***Corresponding author**

Sumi Wijaya

Raya Kalisari Selatan 1 street,

Surabaya, East Java

sumiwijaya@yahoo.com

ABSTRACT

Diarrheal disease is an endemic disease in Indonesia and also a potential disease that can cause death. In practice, the use of traditional medicines derived from plants still being considered by society due to its viability, economic factors and safety reasons. Several plants worked as antidiarrheal agents with the mechanism actions of (1) antibacterial (inhibited the growth of *Escherichia coli*, *Shigella dysenteriae*, *Shigella flexneri*, *Staphylococcus aureus* and *Salmonella Typhi*); (2) reduced the contraction of intestinal as results in the reduction of passage and the reduction of frequency of defecation or; (3) both mechanism of actions. Thus the purpose of the present study is to evaluate antidiarrheal activity of the combination of plant extracts which have been proved to have antidiarrheal activity. Turmeric (*Curcuma domestica*) rhizomes, Gall oak (*Quercus lusitanica*) leaves, Guava (*Psidium guajava*) leaves and Meniran (*Phyllanthus niruri*) herbs were used for this combination. The purpose of this combination was to minimize the doses (based on literature reviews Turmeric at 5% and Meniran 10% inhibited the growth of *Escherichia coli* and *Staphylococcus aureus*, meanwhile guava leaves at 150 mg/kg BW and Gall oak leaves at 10% reduced diarrhea in animal tested). Maceration was chosen for the extraction method using ethanol 96% as the solvent. Protection and transit intestinal method with loperamide HCl as comparison drug and castor oil for the induction of diarrhea were used for antidiarrheal activity, meanwhile diffusion and dilution methods using, inhibition zones, minimum inhibition concentration and minimum bactericidal concentration parameters were used for antibacterial activity. Tetracycline HCl with the concentration of 30 µg was used for the control. The results revealed that the combination of plant extract with the doses of 200 mg/KgBW (1:1:1:1) reduced diarrhea in mice with relaxation of smooth muscle and showed weak antibacterial effects on *Escherichia coli* and *Staphylococcus aureus* but strongly inhibited the growth of *Salmonella Typhi*.



INTRODUCTION

Diarrhea is an alteration in normal bowel movement and is characterized by an increase in the water content, volume, or frequency of stools (Guerrant et al. 2001). Diarrheal diseases are a major problem in Third World countries and are responsible for the death of millions of people each year (Shoba and Thomas 2001). Diarrheal disease is an endemic disease in Indonesia and also a potential disease that can cause death. In practice, the use of traditional medicines derived from plants still being considered by society due to its viability, economic factors and safety reasons. Several plants worked as antidiarrheal agent with the mechanism actions of (1) antibacterial (inhibited the growth of *Escherichia coli*, *Shigella dysenteriae*, *Shigella flexneri*, *Staphylococcus aureus* and *Salmonella Typhi*); (2) reduced the contraction of intestinal as results in the reduction of pain and the reduction of frequency of defecation or; (3) both mechanism of actions. The antibacterial activity include the potency of diminished toxin caused by bacteria especially bacteria that caused food poisoning. Thus the purpose of the present study is to evaluate antidiarrheal activity of the combination of plant extracts which have been proved to have antidiarrheal activity. Turmeric (*Curcuma domestica*) rhizomes, Gall oak (*Quercus lusitanica*) leaves, Guava (*Psidium guajava*) leaves and Meniran (*Phyllanthus niruri*) herbs were used for this combination. The purpose of this combination was to minimize the doses being used. Turmeric at 5% and Meniran 10% inhibited the growth of *Escherichia coli* and *Staphylococcus aureus*, meanwhile guava leaves at 150 mg/kg BW and Gall oak leaves at 10% reduced diarrhea in animal tested (Dzulkarnain et al. 1978; Adyana et al. 2004; Ximenes 2010; Desfita 2011). Protection and transit intestinal method with loperamide HCl as comparison drug and castor oil for the induction of diarrhea were used for antidiarrheal activity, meanwhile diffusion dilution methods using inhibition zones, minimum inhibition concentration and minimum bactericidal concentration parameters were used for antibacterial activity. Tetracycline HCl with the concentration of 30 µg was used for the control.

METHODS

Drugs and chemicals

Ethanol (CV. Lestari, Surabaya, Indonesia), Loperamide (Kimia Farma, Surabaya, Indonesia) was

purchase from local pharmacy, castor oils (Brataco, Surabaya, Indonesia), Nutrient Broth (E.Merck, Germany), Nutrient Agar (E. Merck, Germany)

Plant materials

Turmeric (*C. domestica*) rhizomes, Gall oak (*Q. lusitanica*) leaves, Guava (*P.guajava*) leaves and Meniran (*P niruri*) herbs were obtained in dried powder form from International Herbs Laboratories research.

Animal and Bacteria tested

Mice used for the animal tested were purchased from Animal Laboratories at Faculty of Pharmacy Widya Mandala Catholic University. *S. aureus*, *E. coli* and *Salmonella Typhi* were obtained from Microbiology Laboratories at Faculty of Pharmacy Widya Mandala Catholic University.

Preparation of extracts

Each dried powder of Turmeric (*C. domestica*) rhizomes, Gall oak (*Q. lusitanica*) leaves, Guava (*P. guajava*) leaves and Meniran (*P. niruri*) herbs was soaked in alcohol 96% and leave it for overnight then filtered and evaporated using water bath to dryness. The thick extracts were stored in desiccators until further use.

Antidiarrheal Activity – Protection Method

Animals were divided into three groups of five animals in each group. Group I received normal saline (2ml/KgBW) and served as control. Groups II received standard drug (loperamide 2 mg/kg) and served as standard. Group III received the combination of ethanol extract of Turmeric (*C. domestica*) rhizomes, Gall oak (*Q. lusitanica*) leaves, Guava (*P. guajava*) leaves and Meniran (*P. niruri*) herbs with the doses of 200 mg/KgBW (1:1:1:1). Diarrhea was induced all the overnight fasted animals by administering 1 ml of castor oil orally. The test extracts and the standard drug were administered one hour prior to the treatment of castor oil. Each mice was housed separately and observed for diarrheal episode, for a period of 4 hours. During that period, number and weight of diarrheal feces were taken and noted at every half an hour. The total number of diarrheal feces and percent protection was calculated. The antidiarrheal activity was determined in terms of percentage protection. The data of stool weight was expressed as Mean ± SEM (Awounters et al. 1978).



Antidiarrheal Activity – Transit Intestinal Method

The mice were divided into three groups of five animals each and fasted for 18 h but water was freely provided. The first group (control group) received orally normal saline (2 ml/kg BW), Group II receive standard drug (loperamide 2 mg/KgBW) while the third group was given orally the combination of plant extract in doses of 200 mg/kg BW (1:1:1:1). Thirty minutes later, each animal was given 1 ml of charcoal meal (10% activated charcoal in 0.5% gum acacia) via the oral route. All animals were sacrificed 30 min thereafter and the distance covered by the charcoal meal in intestine, from the pylorus to the caecum was measured and expressed as percentage of distance moved (Mascolo 1994).

Antibacterial Activity – Inhibition zones Determination by Well Diffusion Method

Ten 5×10^8 CFU/ml of suspension contained 1.5×10^8 CFU/ml was poured uniformly to solidified 20 mL Nutrient Agar (NA) and the inoculum was allowed to dry for 5 min. Holes of 6 mm in diameter were made in the seeded agar. Aliquot of 20 μ l from the combination of plant extract (10,000 ppm, 1:1:1:1) and tetracycline (30 μ g) were added into each well on the seeded medium and allowed to stand on the bench for 1 h for proper diffusion and thereafter incubated at 37°C for 24 h. The resulting inhibition zones were measured in millimeters (mm). All the tests were performed in duplicate. Tetracycline (30 μ g) served as positive controls.

Antibacterial Activity – MIC Determination by Microdilution Method

The 96-well plates were prepared by dispensing 50 μ l of Nutrient Broth, into each well. A 50 μ l from the stock solution of tested extracts (concentration of 10,000 ppm) was added into the first row of the plate. Then, twofold, serial dilutions were performed by using a micropipette. The obtained concentration range was from 5000 – 9.77 ppm, and then added 10 μ l of inocula to each well except a positive control (inocula were adjusted to contain approximately 1.5×10^5 CFU/mL. Plant extract with media was used as a positive control and inoculum with media was used as a negative control. The test plates were incubated at 37°C for 18 h. After 18 h 50 μ l of a 0.01% solution of 2, 3, 5-triphenyl tetrazolium chloride (TTC) was added to the wells and the plate was incubated for another hour. Since the colorless tetrazolium salt is reduced to red

colored product by biological active bacteria, the inhibition of growth can be detected when the solution in the well remains clear after incubation with TTC. MIC was defined as the lowest sample concentration showing no color change (clear) and exhibited complete the inhibition of growth (Abu-Shanab et al. 2004)

Antibacterial Activity – MBC Determination by Agar Well Diffusion Method

The lowest sample concentration showing no growth of bacterial tested from MIC determination used as the concentration sample for this method. Aliquot of 10 μ l from the sample were spread on the surface of plate containing Nutrient agar. The plate was incubated at 37°C for 24 h. The presence of colonies was considered a evidence of bacteriostatic action, while the absence of colonies indicated bactericidal activity. All the tests were performed in duplicate.

RESULTS AND DISCUSSION

Our results showed that the combination of extract inhibited castor oil-induced diarrhea in mice, even though not powerful as loperamid (Table 1). Several mechanisms had been previously proposed to explain the diarrheal effect of castor oil. These include inhibition of intestinal $\text{Na}^+ \text{K}^+$ ATPase activity, thus reducing normal fluid absorption (Gaginella and Bass 1978), activation of adenylate cyclase or mucosal cAMP-mediated active secretion (Capasso et al. 1994) stimulation of prostaglandin formation (Galvez et al. 1993) and platelet activating factor (Pinto et al. 1992). Most recent nitric oxide has been claimed to contribute to the diarrheal effect of castor oil (Mascolo et al. 1996). However, it is well documented that castor oil produces diarrhoea due to its most active component ricinoleic acid through a hypersecretory response (Ammon et al. 1974). Therefore it can be assumed that the antidiarrheal action of the extract was mediated by an antisecretory mechanism. Furthermore, the combination of ethanol extract significantly reduced intestinal transit as evidenced by the decrease in the distance traveled by charcoal meal (Table 2). These results also show that the extract suppressed the propulsion of charcoal meal thereby increasing the absorption of water and electrolytes. Antidiarrheal properties of the combination of plant extract was found to be due to tannins (Gall oak (*Q. lusitanica*) leaves and Guava (*P. guajava*) leaves),



Table 1. Effect of combination of extracts Turmeric (*C. domestica*) rhizomes, Gall oak (*Q. lusitanica*) leaves, Guava (*P. guajava*) leaves and Meniran (*P. niruri*) herbs on castor oil induced diarrhea in mice

Group Treatment (p.o)	Total number of diarrheal feces (g)	% Inhibition of diarrhea
I. Saline (2 ml/Kg)	0.5904 ± 1.08%	-
II. Loperamide (5 mg/Kg)	0.3350 ± 0.33%	43.26
III. CE (200 mg/Kg)	0.5140 ± 0.51%	12.94

Value are expressed as mean ±SEM (n=5). CE: combination of extracts Turmeric rhizomes, Gall oak leaves, Guava leaves and Meniran herbs (1:1:1:1)

Table 2. Effect of combination of extracts Turmeric (*Curcuma domestica*) rhizomes, Gall oak (*Quercus lusitanica*) leaves, Guava (*Psidium guajava*) leaves and Meniran (*Phyllanthus niruri*) herbs on small intestinal transits in mice

Group Treatment (p.o)	Inhibition (%)
I. Saline (2 ml/Kg)	-
II. Loperamide (5 mg/Kg)	52.24
III. CE (200 mg/Kg)	96.08

CE: combination of extracts Turmeric rhizomes, Gall oak leaves, Guava leaves and Meniran herbs (1:1:1:1)

Table 3. Diameter inhibition zones, minimum inhibition concentration and minimum bactericidal concentration of combination of extracts Turmeric (*C. domestica*) rhizomes, Gall oak (*Q. lusitanica*) leaves, Guava (*P. guajava*) leaves and Meniran (*P. niruri*) herbs

Bakteri		Tetracycline (ppm)	CE (ppm)
<i>Eschericia coli</i>	DIZ	-	-
	MIC	2500	5000
	MBC	2500	5000
<i>Salmonella thypii</i>	DIZ	-	-
	MIC	625	1250
	MBC	625	1250
<i>Staphylococcus aureus</i>	DIZ	-	-
	MIC	2500	5000
	MBC	2500	5000

CE: CE: combination of extracts Turmeric rhizomes, Gall oak leaves, Guava leaves and Meniran herbs (1:1:1:1) DIZ: Diameter inhibition zones; MIC: minimum inhibition concentration; MBC: minimum bactericidal concentration; -: no activity at the concentration of the extracts tested.

flavonoids (Meniran (*P. niruri*) herbs, and curcuminoids (Turmeric (*C. domestica*) rhizomes) (Longanga *et al.* 2000). The antidiarrheal activity of flavonoids has been ascribed to their ability to inhibit intestinal motility and hydro-electrolytic secretions (Rao *et al.* 1997) which are altered in this intestinal condition. In vitro and in vivo experiments have shown that flavonoids are able to inhibit the intestinal secretory response induced by prostaglandins E₂ (Sanchez *et al.* 1997). In addition, flavonoids present antioxidant properties which are presumed to be responsible for the inhibitory effects exerted upon several enzymes including those involved in the arachidonic acid metabolism (Mora *et al.* 1990).

Tannins present in antidiarrheal plants denature proteins in the intestinal mucosa by forming protein tannates which reduces secretion. The antibacterial activity of combination of ethanol extract of Turmeric (*C. domestica*) rhizomes, Gall oak (*Q. lusitanica*) leaves, Guava (*P. guajava*) leaves and Meniran (*P. niruri*) herbs against *S. aureus*, *E. coli* and *S. Thypi* was assessed by evaluating the presence of DIZ, MIC and MBC values. Results (Table 3) showed that the combination of extracts have weak antibacterial effects on *E. coli* and *S. aureus* but strongly inhibited the growth of *S. Thypi*. The absence of inhibition zone does not necessarily mean that compounds are inactive. For example, non-



polar compounds may not diffuse into the culture medium (Moreno et al. 2006).

CONCLUSION

The results revealed that the combination of plant extract with the doses of 200 mg/KgBW (1:1:1:1) reduced diarrhea in mice with relaxation of smooth muscle and showed weak antibacterial effects on *Escherichia coli* and *Staphylococcus aureus* but strongly inhibited the growth of *Salmonella* Thypi.

ACKNOWLEDGMENT

The author sincerely thanks to PT. International Herbs Research Laboratories International and Traditional Medicines Research Center Widya Mandala Catholic University for funding this research.

REFERENCES

- Abu-Shanab B, Adwan G, Abu-Safiya D, Jarrar N, Adwan K. 2004. Antibacterial Activities of Some Plant Extracts Utilized in Popular Medicine in Palestine. *Turkish Journal of Biology*. 28:99-102.
- Adyana IK, Yulinah E, Sigit JI, Fisheri KN, Insanu M. 2004. Efek ekstrak daun jambu biji daging buah putih dan jambu biji daging buah merah sebagai antidiare. *Acta Pharmaceutica Indonesia*. XXIX (1):19-27.
- Ammon HV, Thomas PJ, Phillips SF. 1974. Effect of the oleic acid and ricinoleic acid net jejunal water and electrolyte movement. *Journal of Clinical Investigation*. 53:374-379.
- Awouters F, Niemegeers CJE, Lenaerts FM, Janseen PAJ. 1978. Delay of castor oil diarrhoea in rats; a new way to evaluate inhibitors of prostaglandin biosynthesis. *Journal of Pharmacy Pharmacology*. 30:41-45.
- Capasso F, Mascolo N, Izzo A.A, Gagarella TS. 1994. Dissociation of castor oil-induced diarrhea and intestinal mucosal injury in rat: effect of NG-nitro-L-arginine methyl ester. *British Journal of Pharmacology*; 113: 1127-1130.
- Desfita V. 2011. Aktivitas Antimikroba Ekstrak Herba Meniran terhadap *Staphylococcus aureus* dan *Escherichia coli*. [Skripsi]. Universitas Sumatera Utara.
- Dzulkarnain B., Wahjoedi, B., Bakar, S., dan Prasetyaningsih N. 1978. Pengaruh ekstrak beberapa tanaman antidisentri dan antidiare terhadap usus kelinci terisolasi. *Hemera Zoa* 70 (1): 27 – 36.
- Gaginella TS and Bass P. 1978. Laxatives: an update on mechanism of action. *Life Science*. 23: 1001-1010.
- Galvez J, Zavzuelo A, Crespo, ME, Lorente MD, Ocete MA, Jimenez J. 1993. Anti-diarrhoeic activity of *Euphorbia hirta* extract and isolation of an active flavonoid constituent. *Planta Medica*, 59: 333-336.
- Guerrant RL, Van Gilder T, Steiner TS, Theilman MN, Slutsker L, Tauxe RV. 2001. Practice guidelines for the management of infectious diarrhea. *Clinical Infectious Disease*, 32: 331-35
- Longanga Otshudi A, Vercruyse A, Foriers A. 2000. Contribution to the ethnobotanical, phytochemical and pharmacological studies of traditionally used medicinal plant in the treatment of dysentery and diarrhea in Lomela area, Democratic Republic of Congo (DRC). *Journal of Ethnopharmacology*, 71(3): 411-423.
- Mascolo, N., Izzo, A.A., Avotore, G., Barboto, F., Capasso, F., 1994. Nitric oxide and castor oil included diarrhea, *Journal of Pharmacology and Experimental therapeutics*, 268: 291-292.
- Mascolo N, Izzo AA, Gagarella TS, Capasso F., 1996. Relationship between nitric oxide and platelet-activating factor in castor oil-induced mucosal injury in the rat duodenum. *Naunyn-Schmiedeberg's Archives of Pharmacology*, 353:680-684.
- Mora A, Paya M, Rios JL, Alcaraz MJ, 1990. Structure activity relationships of polymethoxy flavones and other flavonoids as inhibitors of non-enzymic lipid peroxidation. *Biochemical Pharmacology*, 36: 317-322.
- Moreno S., Scheyer T., Romano C.S., Vojnov A.A. 2006. Antioxidant and antimicrobial activities of rosemary extracts linked to their polyphenol composition. *Free Radical Research*, 40: 223-231.
- Pinto A, Autore G, Mascolo N, Sorrentino R, Biondi A, Izzo AA, Capasso F. 1992. Time course of PAF formation by gastrointestinal tissue in rats after castor oil challenge. *Journal of Pharmacy and Pharmacology*, 44: 224-226.
- Rao VSN, Santos FA, Sobreika TT, Souza MF, Melo LL, Silveira ER., 1997. Investigation on the gastroprotective and antidiarrheal properties of ternatin, a tetramethoxyflavone from *Egletes viscoso*. *Planta Medica*, 63: 1496-1497.



- Sanchez de Medina F, Galvez J, Gonzalez M, Zarzuelo A, Barrett KE. 1997. Effects of quercetin on epithelial chloride secretion. *Life Science*, 61: 2049-2055.
- Shoba FG, Thomas M. 2001. Study of antidiarrheal activity of four medicinal plants in castor oil-induced diarrhea. *Journal of Ethnopharmacology*, 76: 73-76.
- Ximenes, J.M.C. 2010, 'Efek infusa rhizoma *Curcuma domestica* Vall (rimpang Kunyit) dan *Zingiber officinale* Rosc. (rimpang jahe) terhadap motilitas usus pada mencit putih galur Swiss Webster menggunakan metode transit intestinal', [Skripsi], Sarjana Kedokteran, Universitas Kristen Maranatha, Bandung.



Antidiarrheal activity of the combination of ethanol extract of Tumeric.

ORIGINALITY REPORT

% **17**
SIMILARITY INDEX

% **8**
INTERNET SOURCES

% **15**
PUBLICATIONS

% **5**
STUDENT PAPERS

PRIMARY SOURCES

1 tpcj.org Internet Source % **2**

2 Ivana Radojevic, Sava Vasic, Ljiljana Comic, Srecko Trifunovic, Marina Mijajlovic, Milos Nikolic, Gordana Radic. "Antibacterial and antibiofilm screening of new platinum(IV) complexes with some s-alkyl derivatives of thiosalicylic acid", Kragujevac Journal of Science, 2017
Publication % **1**

3 Uddin, S.J.. "Antidiarrhoeal activity of the methanol extract of the barks of *Xylocarpus moluccensis* in castor oil- and magnesium sulphate-induced diarrhoea models in mice", Journal of Ethnopharmacology, 20051003
Publication % **1**

4 Perumal, Shanmugapriya, Roziahanim Mahmud, Suthagar Pillai, Wei Cai Lee, and Surash Ramanathan. "Antimicrobial Activity and Cytotoxicity Evaluation of *Euphorbia hirta* (L.) % **1**

Extracts from Malaysia", APCBEE Procedia,
2012.

Publication

5

Submitted to Gauhati University

Student Paper

% 1

6

www.insidemoringa.com

Internet Source

% 1

7

Rosa Ventura-Martínez, Oscar Rivero-Osorno,
Claudia Gómez, María Eva González-Trujano.
"Spasmolytic activity of Rosmarinus officinalis L.
involves calcium channels in the guinea pig
ileum", Journal of Ethnopharmacology, 2011

Publication

% 1

8

Submitted to Universiti Teknologi Malaysia

Student Paper

% 1

9

Daiane Voss-Rech, Cátia Silene Klein, Vânia
Helena Techio, Gerson Neudi Scheuermann,
Gilberto Rech, Laurimar Fiorentin. "Antibacterial
activity of vegetal extracts against serovars of
Salmonella", Ciência Rural, 2011

Publication

% 1

10

Nonglak Laoprom, Araya Sangprom,
Patcharaporn Chaisri. "Antioxidant and
antibacterial activity of Thai medicinal plant
(Capparis micracantha)", AIP Publishing, 2018

Publication

% 1

11

Submitted to NCC Education

Student Paper

% 1

12

www.peertechz.com

Internet Source

% 1

13

Wendel, G.H.. "Antidiarrheal activity of dehydroleucodine isolated from *Artemisia douglasiana*", *Fitoterapia*, 200801

Publication

% 1

14

Shilpi, J.A.. "Preliminary pharmacological screening of *Bixa orellana* L. leaves", *Journal of Ethnopharmacology*, 20061124

Publication

% 1

15

jrkuiate.blogspot.com

Internet Source

% 1

16

Md. Al Foyjul Islam, Anindya Barua, Maksud Hassan Binoy, Md. Jahir Alam, Md. Ershad, M. S. K. Choudhuri, Runa Masuma. "Pre-clinical investigation of anti-diarrheal and CNS depressant effect of *Wedelia trilobata* in Swiss albino mice", *Oriental Pharmacy and Experimental Medicine*, 2019

Publication

% 1

17

Umukoro, S., and R.B. Ashorobi. "Effect of *Aframomum melegueta*. Seed Extract on Castor Oil–Induced Diarrhea", *Pharmaceutical Biology*, 2005.

% 1

18

www.dovepress.com

Internet Source

% 1

19

Ripa, FA, MTI Morshed, AA Sharmin, SB Papon, R Islam, and Z Sheikh. "Central Nervous System Depressant, Analgesic and Antidiarrheal Effects of the Seed Extracts of *Dimocarpus longan* Lour in Rats", *Tropical Journal of Pharmaceutical Research*, 2014.

Publication

% 1

20

healthy-synergies.com

Internet Source

<% 1

21

pubs.rsc.org

Internet Source

<% 1

EXCLUDE QUOTES ON

EXCLUDE ON

BIBLIOGRAPHY

EXCLUDE MATCHES < 10

WORDS