



The Joint Committee of  
**International Conference and Talk Show on Medicinal Plant**

“Effective, Safe and Qualified Herbal Medicine for Diabetes Mellitus Treatment”

hereby acknowledge the participation and valued contribution of

**DR. LANNIE HADISOEWIGNYO, M.SI, APT.**

**PARTICIPANT AND PRESENTER**

in Jakarta, 19<sup>th</sup> - 21<sup>st</sup> October 2010

**Dr. Rifatul Widjhati, M.Sc., Apt.**

Director  
Center for the Pharmaceutical and Medical Technology

**Indah Yuning Prapti, SKM, M. Kes**

General Secretary  
National Working Group on Indonesian Medicinal Plant

**dr. Hardhi Pranata Sp.S., MARS**

President  
Association of Indonesian Herbal Medical Doctors



**ORGANIZERS**

Agency for the Assessment and Application of Technology  
National Working Group on Indonesian Medicinal Plant  
Association of Indonesian Herbal Medical Doctors  
Deutscher Akademischer Austausch Dienst



39<sup>th</sup> Meeting of National Working Group on Indonesian Medicinal Plant

# PROCEEDING



## INTERNATIONAL CONFERENCE AND TALK SHOW ON MEDICINAL PLANT

*Effective, Safe and Qualified Herbal Medicine for Diabetes Mellitus Treatment*



**Jakarta, 19<sup>th</sup> – 21<sup>st</sup> of October 2010**

**Organized by :**

Agency for the Assessment and Application of Technology (BPPT)  
National Working Group on Indonesian Medicinal Plant (POKJANAS TOI)  
Association of Indonesian Herbal Medical Doctors (PDHMI)

**Supported by :**

Deutscher Akademischer Austausch Dienst (DAAD)

**PROCEEDING INTERNATIONAL CONFERENCE AND TALK SHOW ON  
MEDICINAL PLANT “Effective, Safe and Qualified Herbal  
Medicine for Diabetes Mellitus Treatment”**

**First Edition**, March 2011



**ISBN :**

ISBN 978-602-95911-1-8



9 786029 591118

**PUBLISHER**

Center for Pharmaceutical and Medical Technologies  
Agency for the Assessment and Application of Technology  
BPPT Building II, 15<sup>th</sup> Floor  
Jl. MH Thamrin No. 8, Jakarta 10340  
INDONESIA

**PROCEEDING INTERNATIONAL CONFERENCE AND TALK SHOW ON  
MEDICINAL PLANT “Effective, Safe and Qualified Herbal  
Medicine for Diabetes Mellitus Treatment”**

**First Edition**, March 2011

**ISBN :**



**PUBLISHER**

Center for Pharmaceutical and Medical Technologies  
Agency for the Assessment and Application of Technology  
BPPT Building II, 15<sup>th</sup> Floor  
Jl. MH Thamrin No. 8, Jakarta 10340  
INDONESIA

## **WELCOME MESSAGE**

Praise and gratitude towards Allah who has given His Grace so that the International Conference and Talk Show on Medicinal Plant could be conducted in BPPT Jakarta on the 19<sup>th</sup> – 21<sup>st</sup> October 2010.

This International Conference and Talk Show on Medicinal Plant which is also the 39<sup>th</sup> Meeting of National Working Group on Indonesian Medicinal Plant is organized by BPPT in collaboration with the National Working Group on Indonesian Medicinal Plants (POKJANAS TOI), Deutscher Akademischer Austausch Dienst (DAAD) and the Association of Indonesian Herbal Medical Doctors (PDHMI). This Conference is an effort to develop the national herbal medicine industry and as a venue for researchers, policy makers, clinicians, industrialists and observers of the herbal medicine to gather and discuss the issues regarding herbal medicine.

The theme of this Conference is “Effective, Safe and Qualified Herbal Medicine for Diabetes Mellitus Treatment”. This theme was chosen because diabetes mellitus is one of diseases that could become a serious threat to public health in the future and currently herbal medicine has not been optimally utilized for the treatment of various diseases, especially diabetes mellitus. This Conference is expected to provide an overview of recent advances in herbal medicine technology, herbal medicine role in the treatment of diabetes, the problems faced by the herbal medicine industry and alternative solutions.

This conference has brought together 88 research papers from researchers of various institutions with a variety of topics, ranging from the cultivation aspects to clinical trials which provide an overview of the research currently conducted by research institution in Indonesia. Through this Conference, it is expected that research institutions with similar interest could collaborate to accelerate research in Indonesian and also as a venue for facilitating collaboration between research institutes, universities and industries.

The committee realizes that there are still many shortcomings that may be less than satisfactory to many parties. Criticisms and suggestions are expected to improve this organization of seminars in the future.

On this occasion the committee expressed the highest appreciation to the guest speakers, papers contributors, participants and sponsors for their cooperation in organizing the International Conference and Talk Show on Medicinal Plant

Jakarta, 19<sup>th</sup> of October 2010

*Organizing Committee*

*International Conference and Talk Show on Medicinal Plant*





**The Agency for the Assessment and Application of Technology**  
**BADAN PENGKAJIAN DAN PENERAPAN TEKNOLOGI (BPPT)**  
**BPPT Chairman**

*His Excellency The Minister of Science and Technology, Mr. Suharna Surapranata*

*The Director of Deutscher Akademischer Austausch Dienst (DAAD), Indonesia  
Chairman of The Association of Indonesian Herbal Medical Doctors (PDHMI)  
Distinguished Speakers from China, Malaysia and Republic of Korea  
Chairmen of Pharmaceutical Industries,*

*My colleagues and all participants,*

*Ladies and Gentlemen,*

*Assalamu Alaykum wa rahmatullahi wa barakatuh.*

*Good morning, Selamat Pagi, Salam Sejahtera untuk kita semua.*

*May God Almighty blesses us in conducting this event*

Let us start by expressing our gratefulness to God almighty. His bless and mercy has given us opportunity to be here in the great and special event of **International Conference and Talkshow on Medicinal Plants** which focus on the theme of “Effective, Safe and Qualified Herbal Medicine for Diabetes Mellitus Treatment”.

I am delighted to welcome all distinguished guests, invited speakers and participants. To speakers coming from overseas, welcome to Jakarta and welcome to Indonesia. Jakarta is very wet at the moment, due to heavy rain in the last couple of months. I hope this will not discourage you to enjoy your visit to Jakarta.

In this very important moment, allow me to thank to the committee who has been working very hard to bring this event into reality. I also would like to express a special thank to the **Association of Indonesian Herbal Medical Doctors** (PDHMI) who has been actively involved and support to this event. And to **Deutscher Akademischer Austausch Dienst (DAAD)** or **German Agency for Academic Exchanges** as a main sponsor, gratitude thanks and most appreciation shall be addressed.

*Ladies and Gentlemen,*

I would like to draw your attention that BPPT as a research agency under coordination of the Ministry of Research and Technology, highly supports the development of herbal medicines as one of the BPPT program on health and pharmaceutical sector especially in fulfilling the national self-sufficiency on medicines. Beside Health and medicine, BPPT is conducting other focused program, namely food, energy, ICT, transportation, defence, environment, and manufacturing.

I would like to highlight here that herbal medicine development is strategic choice for developing country like Indonesia where mega biodiversity is one of its comparative advantages. Currently, chemical and semi-chemical drug industries has been dominated by global pharmaceutical industries in developed countries. For this reason, competition in this area is not beneficial for Indonesian pharmaceutical industries where its industrial structure in general, is still very weak and need to be further developed. The choice to promote program on herbal medicine development is the fact that this program has strong national capacity building from upstream to downstream processes by using locally available resources and involving various stakeholders.

*Ladies and gentlemen,*

This is our main mission to relate the most suitable technology to significantly increase the added value of our bioresources in order to support our national herbal medicine industries. Knowledge and technology would be a fundamental tools in the development of herbal medicine industries which enable to support national economic growth (known as knowledge based economy).

I do realise that to achieve the noble goal is not trouble-free. We need an active participation of all players not only from government institutions as facilitatory and regulatory bodies, but also from private sectors, academicians as well as researchers to work hand-in-hand toward the best



in the utilization of knowledge and local resources in natural product development.

*Ladies and gentlemen,*

Since the enthusiasm of back to nature have been gaining popularity and acceptance in recent decade, national herbal medicine industries should become one of the pillars of national economic development. This is also supported by the fact that Indonesia has a second largest biodiversity in the world, with more than 30,000 species of plants. The number of herbal medicine industries keep increasing, numbering to 1,500 companies, big and small corporation. According to Herbal Medicine Entrepreneur Association (**GP Jamu**) data in 2007, not less than 3 million people works on herbal medicine sector from upstream to downstream industries. The market share of herbal medicine has growing significantly, up to 15-20% yearly.

We all hope that herbal medicine will not only fulfil local and national market, but more than that will expand to other countries, as export commodity.

*Ladies and gentlemen,*

As I have just mentioned before, BPPT, through the **Centre of Pharmaceutical and Medical Technology** - fully supports the herbal medicine development. I hope throughout this great forum, all stakeholders involved in the utilization of medicinal plants will stumble on the best and most suitable ways to develop our biological resources as health products. We would like to utilize our biodiversity as source of natural products to prevent and cure diseases, especially degenerative diseases such as diabetes mellitus, and also to promote our well being. To PDHMI, the **Assosiation of Indonesian Herbal Medical Doctors** we would certainly rely on your share in using herbal medicine as part of formal health services.

At last, we should work closely with all experts and institutions who concern in this area not only from our national side but also from abroad. Therefore, BPPT is continually welcome to all collaborators from national as well as international institutions based on our mutual benefit for each parties involved.

*Ladies and gentlemen,*

Once again, I would like to thanks to the committee, sponshors, speakers and participants and wishing all a successful symposium.

Finally, by saying “*Bismillahirrohmanirrohim*”, in the name of GOD Almighty, the Most Gracious and the Most Merciful, I have the great pleasure to declare the **International Conference and Talkshow on Medicinal Plants** is officially opened.

Thank you very much for your passion and kind attention,

*Wassalamu’alaykum warahmatullahi wabarakatuhu*

BPPT Chairman,

Dr. Ir. Marzan Azis Iskandar

## CONTENT

Welcome Message	i
Opening Speech by BPPT Chairman	iii
Content	vii
RECENT ADVANCES OF NATURAL PRODUCT RESEARCH <i>Wahono Sumaryono</i>	1
DIURETIC EFFECT OF AQUEOUS EXTRACT OF LEAF OF <i>Averrhoa bilimbi</i> L. IN WISTAR RATS <i>Afifah B. Sutjiatmo, Elin Yulinah, Suswini K., Suci Narvikasari, Ita Nur Anisa</i>	22
HYPOGLYCAEMIC ACTIVITY OF AQUEOUS EXTRACT OF LEAVES OF <i>Strobilanthes crispus</i> L. IN MALE WISTAR RATS USING GLUCOSE TOLERANCE TEST METHOD <i>Afifah B. Sutjiatmo, Sri Wahyuningsih, Suci Nar Vikasari, Sri Wijayanti</i>	27
COMPARATIVE STUDY OF PRIMOJEL® 2%, 5%, & 8% TO PHYSICAL CHARACTERISTICS OF TABLET FROM DRIED EXTRACT <i>Murraya paniculata</i> (L.) Jack LEAVES <i>Agnes Nuniek Winantari, I. I. Panigoro, Ferrany</i>	32
IN VITRO BIOASSAY OF N-BUTHANOL ISOLATE OF <i>Acorus calamus</i> L. ON INHIBITORY OF ACTIVITY $\alpha$ -GLUCOSIDASE <i>Ahmad Najib, Sri Hartati, Berna Elya</i>	38
ISOLATION OF MAJOR COMPOUND HEXANE FRACTION FROM <i>Piper betle</i> L. LEAVE AND CHROMATOGRAM PROFILE <i>Ahwan, Haryoto, Haikal Sofyan Arif</i>	44
CLINICAL STUDY OF ETHANOL EXTRACT OF <i>Justicia gendarussa</i> Burm. f. LEAVES IN MALE FERTILITY <i>Bambang Prajogo E.W., Hudi Winarso, Kasmijati, Aucky Hinting; Dyan Pramesti, Maria Anggraeni, Achmad Radjaram, Dwi Setiawan, Sri Musta'ina, Noor Cholies Zaini</i>	49
THE MECHANISM OF ACTION INVESTIGATION OF ROCAGLAMIDES AS ANTIPROLIVERATIVE AGENTS IN CHRONIC MYELOID LEUKEMIA USING PATHWAY-BASED PREDICTION BY COMPUTATIONAL DOCKING <i>B.S. Ari Sudarmanto, Fajar S. Wibowo, Chaidir</i>	60

EFFECT OF <i>Curcuma xanthorrhiza</i> EXTRACT ON LYMPHOCYTE CELL PROLIFERATION <i>Churiyah, Tarwadi, Fery Azis Wijaya, Elrade Rofaani</i>	70
CYTOTOXIC ACTIVITY OF MARINE SPONGE <i>Petrosia</i> sp. FROM WEST SUMATERA, INDONESIA <i>Dian Handayani, Mega Yulia, Yohannes Allen</i>	77
INHIBITION EFFECT OF SIDAGURI ( <i>Sida rhombifolia</i> ) AND CELERY ( <i>Apium graveolens</i> ) EXTRACT FORMULA ON THE ACTIVITY OF XANTHINE OXIDASE AND ITS ANTI-INFLAMMATION EFFECT <i>Dyah Iswantini, Iskandar, Lukmanul Hakim.</i>	81
METHANOL EXTRACT DERIVED FROM NIAS'S AND MENTAWAI'S SPONGES AND SOFT CORALS AS ANTI ANGIOGENESIS <i>Dyah Iswantini, Min Rahminiwati, Latifah K Darusman, Rudi Heryanto</i>	91
SIMULTANEOUS ANALYSIS OF PARACETAMOL, PROPYPHENAZONE AND CAFFEINE MIXTURE IN TABLETS <i>Effendy De Lux Putra, Muchlisyam, Mustika Furi, Nerdy</i>	100
ANTIBACTERIAL POTENTIAL OF MARINE ALGAE COLLECTED FROM SOUTH SULAWESI COAST AGAINST HUMAN PATHOGENS <i>Elmi Nurhaidah Zainuddin</i>	115
PROGRESS RESEARCH OF TEMOELAWAK ( <i>Curcuma xanthorrhiza</i> Roxb.) TO PROVIDE HEALTHY PLANTING MATERIAL BY <i>IN VITRO</i> CULTURE <i>Endang Hadipoentyanti, Sitti Fatimah Syahid</i>	128
EVALUATION OF GROWTH, PRODUCTIVITY AND OIL QUALITY FOUR PROMISING NUMBERS OF MENTHA ( <i>Mentha arvensis</i> L.) <i>Endang Hadipoentyanti, Sri Suhesti, Amalia, Nursalam</i>	144
EFFECT OF GROWTH ENVIRONMENT AND CULTIVATION METHODS SMAL WHITE GINGER ( <i>Zingiber officinale</i> , L. <i>var. amarum</i> ) ON SEED PRODUCTION, FARMER INCOME AND FEASIBILITY OF FARMING SYSTEM <i>Ermianti, Sukarman</i>	152
ECONOMIC ANALYSIS OF FARMING SYSTEM WHITE BIG GINGER ( <i>Zingiber Officinale</i> L. <i>Var. Officinale</i> ) SEED PRODUCTION <i>Ermianti, Sukarman</i>	161

TABLET FORMULATION OF <i>Andrographis paniculata</i> Ness DRY EXTRACT BY DIRECT COMPRESSION METHOD <i>Etik Mardiyati, Hafizah, Farida Sulistiawati</i>	170
ANTIMICROBIAL OF SOME PARASITE PLANT IN THE FAMILY LORANTHACEAE <i>Fairus Zabadi, Haryoto, Peni Indrayudha</i>	177
STRUCTURE MODIFICATION OF THYMOQUINONE AND CYTOTOXICITY OF THE PRODUCTS AGAINST HEPATOCELLULAR CARCINOMA (HepG2) CELLS <i>Firdayani, Siswandono, Marcellino Rudyanto</i>	181
STUDY ON THE DPPH AND $\alpha$ -GLUCOSIDASE INHIBITORY POTENTIAL OF VARIOUS GAMBIER AVAILABLE IN INDONESIAN MARKET <i>Galuh Widiyarti, Andini Sundowo</i>	189
ANTIOXIDANT ACTIVITY OF TOTAL PHENOLIC COMPOUND IN HEXANE FRACTION FROM <i>Piper betle</i> L. LEAF WITH DPPH METHODE <i>Haryoto, Ahwan, Andika Punama Devi</i>	199
ACUTE TOXICITY TEST OF <i>Typhonium divaricatum</i> (KELADI TIKUS) ETHANOL EXTRACT IN MICE <i>Indah.D.Dewijanti, Marissa Angelina, Sofna D. Banjarnahor, Rina Andriyani, M. Hanafi</i>	206
ANTIPLASMODIAL EVALUATION OF ONE COMPOUND FROM <i>Calophyllum. depressinervosum</i> <i>Jamilah Abbas, Nina Artanti, Puspa Dewi, Puji Budi Setia Asih, Syafruddin</i>	212
CYTOTOXIC COMPOUND FROM <i>Calophllym</i> spp (GUTTIFERAE) <i>Jamilah Abbas</i>	222
FUCOXANTHIN IDENTIFICATION IN <i>Sargassum cinereum</i> , <i>Sargassum filipendula</i> , AND <i>Sargassum echinocarpum</i> BROWN ALGAE IN KECAMATAN TALANGO, SUMENEP, MADURA AT GPS (113.94444°BT – 7.08795°LS, 113.94231°BT – 7.08913°LS, 113.94548°BT – 7.08911°LS AND 113.94347°BT – 7.08999) <i>Kartini Zailanie, Sri Dayuti</i>	232
CORRELATION BETWEEN CYTOTOXIC ACTIVITY ON MCF-7 CELL AND TOTAL STEROID LEVEL ON <i>Trigonella foenum-graecum</i> .L <i>Kurnia Agustini, Wahono Sumaryono, Azri Fitria</i>	238

ROLE OF NATURAL POLYSACCHARIDES XANTHAN GUM- LOCUST BEAN GUM IN PER-ORAL DRUG DELIVERY SYSTEMS <i>Lannie Hadisoewignyo, Achmad Fudholi</i>	246
CYTOTOXIC EFFECT OF ETHANOL AND WATER EXTRACT OF <i>Centella asiatica</i> ON HUMAN BREAST CANCER T47D CELLS <i>Lucia Hendriati, Ferawati, Adrianta Surjadhana, Arijanto Jonosewojo, Elisabeth Catherina Widjajakusuma</i>	253
ACUTE TOXICITY STUDIES OF <i>Brucea javanica</i> Merril LEAVES EXTRACT ON MICE <i>Marissa A., Indah D. Dewijanti, Banjarnahor S.D.S, Megawati M. Hanafi</i>	258
PHYTOCHEMISTRY, TOXICITY AND ANTIOXIDANT ACTIVITIES OF <i>Kalanchoe pinnata</i> (Lam.) Pers. LEAVES EXTRACTS <i>Megawati, A. Darmawan, Sofa Fajriah, Lia Meilawati</i>	272
IN VITRO ACTIVITY OF CERTAIN PLANT EXTRACTS IN COMBINATION WITH VANCOMYCIN AGAINST METHYCYLIN RESISTANT <i>Staphylococcus aureus</i> <i>Murti, Y.B., Utami, D., Triutami, S., Cahayani, D.D., Mirayany, B.F., Ratnasari, D.</i>	265
POTENSI EKSTRAK ETANOL TUMBUHAN BABANDOTAN ( <i>Ageratum conyzoides</i> L) SEBAGAI BAHAN ANTIKANKER. <i>Pertamawati, Nuralih</i>	277
INDUKSI PEMBENTUKAN BATU GINJAL PADA TIKUS PUTIH JANTAN GALUR <i>Sprague Dawley</i> SECARA KIMIAWI UNTUK UJI EFIKASI HERBAL PELURUH BATU GINJAL (KALSIUM OKSALAT) <i>Pertamawati, Sriningsih, Kurnia Agustini,</i>	281
MELEAGRINE, A ISOLATE COMPOUND FROM SKY III.2.1 AN ENDOPHYTIC FUNGI FROM <i>Annona squamosa</i> . <i>Prasetyawan Yuniyanto, Yudi Rusman, Agus Supriyono, Wahyudi Priyono Suwarso<sup>3</sup></i>	296
CYTOTOXICITY COMPARATION OF ETHYL ACETATE EXTRACT AND THEIR ISOLATE FROM SKY III.2.1, AN ENDOPHYTIC FUNGI FROM SRIKAYA PLANTS ( <i>Annona squamosa</i> ) <i>Prasetyawan Yuniyanto, Fery Azis Wijaya, Nurhadi, Syofi Rosmalawati, Wahyudi Priyono Suwarso, Wahono Sumaryono</i>	304

ISOLATION AND IDENTIFICATION OF A CHEMICAL COMPOUND IN <i>n</i> - HEXANE PHASE FROM THE METHANOL EXTRACT OF BINAHONG LEAF ( <i>Anredera cordifolia</i> (Ten.) Steenis), BASELLACEAE <i>Ratna Djamil, Wahyudi PS, Wahono S, Wiwi Winarti, Evi Nurlativah</i>	310
ANTIDIABETIC PROPERTIES OF <i>Andrographis paniculata</i> Nees AND <i>Eugenia polyantha</i> Wight LEAVES IN WISTAR RATS BY ORAL GLUCOSE TOLERANCE TEST <i>Ratna Megawati Widharna, Ferawati, Lucia Hendriati<sup>1</sup>, Adrianta Surjadhana, Arijanto Jonosewojo, Elisabeth Catherina Widjajakusuma</i>	318
THE EFFECT OF SEED TREATMENT ON GINGER PLANT PARASITIC NEMATODE AND SCALE INSECT POPULATION DEVELOPMENT IN THE FIELD <i>S. Retno Djiwanti, Rodiah Balfas</i>	325
THE STUDY OF BAP AND IBA CONCENTRATION ON THE CALLUS FORMATION AND GROWTH OF <i>Artemisia annua</i> L. BY IN VITRO <i>Samanhudi</i>	332
EFFECT OF BENZYL ADENIN AND THIDIAZURON ON SHOOT MULTIPLICATION OF ST. JOHN'S WORT ( <i>Hypericum perforatum</i> ) IN VITRO <i>Sitti Fatimah Syahid</i>	341
DEVELOPMENT OF ACTIVE COMPOUNDS FROM ETHYL ACETATE EXTRACT OF RHIZOME <i>Acorus calamus</i> L AS ANTIDIABETIC <i>Sri Hartati, Rizna T. Dewi, A. Darmawan, Megawati</i>	346
THE INFLUENCE OF LONG TERM BLOOD GLUCOSE-LOWERING JAMU ADMINISTRATION ON RAT KIDNEY <i>Sri Ningsih, Rilianawati, Agung Eru Wibowo</i>	355
ANTIDIABETIC ACTIVITY OF MAHKOTA DEWA [ <i>Phaleria macrocarpa</i> (Scheff.) Boerl.] STEM EXTRACTS AS AN INHIBITOR OF ALPHA-GLUCOSIDASE <i>Sri Sugiwati, Siswati Setiasih</i>	365
ANTIDIABETIC AND ANTIOXIDANT ACTIVITIES FROM SEED EXTRACTS OF ROSELLA ( <i>Hibiscus Sabdariffa</i> L.) <i>Sri Sugiwati, Hani Mulyani, Megawati, Minarti</i>	374



UJI ANTIMIKROBA EKSTRAK ETANOL DAUN BINAHONG ( <i>Anredera Cardifolia</i> (Ten.) Steenis) DAN DALAM SEDIAAN SALEP DENGAN BASIS HIDROKARBON DIBANDINGKAN DENGAN BASIS ABSORPSI <i>Sudana Atmawidjaja, Ratna Djamil, Muhammad Yunus</i>	381
A PHYTOCHEMICAL AND PHARMACOLOGICAL STUDY OF <i>Averrhoa bilimbi</i> L. LEAVES <i>Suswini Kusmaningati, Afifah B. Sutjiatmo, Maria Immaculata Iwo</i>	404
INSECTS ASSOCIATED WITH HORSEWOOD PLANTS ( <i>Anis clausena</i> ) <i>Tri L. Mardiningsih, Dewi Sartiami</i>	408
PLANT DAMAGING ORGANISMS OF JAVA TURMERIC ( <i>Curcuma xanthorrhiza</i> ) <i>Tri L. Mardiningsih, Miftahurrohmah, Ripah Karyatiningsih, D. Anser Sianturi, Desmawati, Shinta Ramadhani, Lilik Sri Utami</i>	417
GROWTH AND YIELD OF <i>Artemisia annua</i> L. ON NITROGEN FERTILIZER AND CYTOKININ <i>Usman Siswanto, Ruri Kustia, Entang Inorih, Yuli Widiyastuti</i>	426
ISOLATION AND IDENTIFICATION OF A CHEMICAL COMPOUNDS IN <i>n</i> -HEXANE PHASE FROM THE METHANOL EXTRACT OF KELADI TIKUS LEAF ( <i>Typhonium flagelliforme</i> (Lodd.) Blume, Araceae. <i>Yunahara Farida, Resti Caroline. A, Wahyudi Priyono S., Wahono Sumaryono</i>	433
DETERMINATION OF EUGENOL IN VOLATILE OIL FROM CLOVE LEAF AND FLOWER ( <i>Eugenia aromatica</i> OK) BY TLC- DENSITOMETRY <i>Zuhelmi Aziz, Ratna Djamil, Desi Rahmi Suciati</i>	440
POTENCY OF SONGGA TREE ( <i>Strychnos lucida</i> R Brown) AS TRADITIONAL MEDICINAL PLANT : A CASE STUDY IN SUMBAWA ISLAND- INDONESIA <i>Zuraida</i>	448

## **ROLE OF NATURAL POLYSACCHARIDES XANTHAN GUM- LOCUST BEAN GUM IN PER-ORAL DRUG DELIVERY SYSTEMS**

***Lannie Hadisoewignyo<sup>1)</sup>, Achmad Fudholi<sup>2)</sup>***

<sup>1)</sup>Faculty of Pharmacy, Widya Mandala Catholic University  
Jl. Dinoyo 42-44, Surabaya, Phone(+62)031-5678478

<sup>2)</sup>Faculty of Pharmacy, Gadjah Mada University

Corresponding author: e-mail: lanhadi@yahoo.com

### **ABSTRACT**

Xanthan gum is a heteropolysaccharide derived from corn starch and produced by fermentation with the bacteria *Xanthomonas campestris*. Locust bean gum is a natural hydrocolloid obtained from the seed endosperm of the carob tree (*Ceratonia siliqua* L.). The structure of locust bean gum consists of a  $\beta$ -(1 $\rightarrow$ 4)-D-mannosa backbone to which D-galactose units are attached  $\alpha$ -(1 $\rightarrow$ 6). Locust bean gum is a non ionic polysaccharide, which can form gel if it is combined with xanthan gum. Locust bean gum is capable of synergistic interactions with xanthan gum.

This combination can be used for a matrix of a sustained release dosage form based on the ability of the combination of xanthan gum and locust bean gum to form gel. In this research, ibuprofen was used as an active ingredient formulated in control release tablet dosage form.

The results exhibited that the combination of xanthan gum and locust bean gum at a ratio of 1:1 gave the longest in vitro release. Kinetic drug release tend to zero order with dominant mechanism is erosion.

*Keywords:* xanthan gum, locust bean gum, matrix, sustained release.

### **INTRODUCTION**

Xanthan gum is a heteropolisakarida which has a large molecular weight and produced from the fermentation of carbohydrates by the microorganism *Xanthomonas campestris*. It has a double helical structure of antiparallel bimolekular. Glucan backbone structures protected by the side chains are along the sides, which makes xanthan gum is relatively stable on heating as well as in acid or alkaline conditions. Xanthan gum is easily hydrated in

systems containing water as it can dissolve in cold or hot water (Anonymous, 2005<sup>b</sup>).

Locust bean gum is often also called carob bean gum or a galaktomanan carubin is obtained from the seeds of *Ceratonia siliqua* extraction. Locust bean gum is a galaktomanan, containing chains (1→4) - are associated with β-D-mannopiranosas as the main chain, which in the sixth position associated with α-D-galactose. Locust bean gum is a non-ionic compound so that the ionic strength and pH has no effect on the viscosity, except at extreme pH and high temperature. Locust bean gum partially hydrated in cold water and perfect hydrated when heated (Anonymous, 2005<sup>a</sup>).

Xanthan gum when combined with locust bean gum to form a gel when liquid is in an environment (Baichwal, 2001). This behavior is called synergism. Medicine that is often used so frequently in a day that it can cause the patient to forget to take it and less effectiveness in the treatment. To overcome these events experts designing and developing pharmaceutical dosage forms which can release the drug gradually so that the availability of drugs in the blood can be maintained longer and the drug action can be extended. This dosage form is known as a sustained release dosage form (Ansel, 1989).

Ibuprofen [(±)-2 - (p-isobutilfenil) propionic acid) is a nonsteroidal anti-inflammatory (NSAID) that is often used. Ibuprofen has a short biological half-life of approximately two hours that causes need to be used repeatedly in a day. In the form of tablets are generally used in doses of 200 mg to 800 mg, three to four times daily or 600 mg to 3200 mg per day. This is why ibuprofen is formulated in a dosage suitable for slow release (Kurumaddali et al., 1994, Tehrani and shobeiri, 1995; Shehab and Richards, 1996).

Pembuatan suatu matriks dengan bahan obat berada di dalamnya atau tercampur homogen dengan bahan matriks merupakan salah satu cara yang dapat digunakan untuk membuat sediaan lepas lambat. Bahan pembentuk matriks dapat berupa bahan yang tidak larut dalam air maupun bahan yang larut dalam air dan membentuk gel. Salah satu polimer yang sekarang mulai banyak digunakan untuk matriks sediaan lepas lambat adalah kombinasi dari *xanthan gum* dan *locust bean gum*.

One way to make sustained released preparation use a matrix system. The matrix can be developing with water soluble material and form a gel when liquid is in environment. Combination of xanthan gum and locust bean gum can be consider to be use as matrix sustained release tablet (Anonim, 2005).

According to Higuchi (1963), the release of drug dispersed in a homogeneous solid and inert matrix can be describe following equations (1):

$$Q = [D(2A - Cs)Cs.t]^{1/2} \dots\dots\dots (1)$$

The rate of drug release at once (shortly) at time t obtained by differentiate equation (1), and obtained:

$$\frac{dQ}{dt} = \frac{1}{2} \left[ \frac{D(2A - C_s)C_s}{t} \right]^{1/2} \dots\dots\dots (2)$$

In general,  $A \gg C_s$ , so that equation (1) becomes

$$Q = (2ADC_s t)^{1/2} \dots\dots\dots (3)$$

and equation (2) becomes

$$\frac{dQ}{dt} = \left[ \frac{ADC_s}{2t} \right]^{1/2} \dots\dots\dots (4)$$

with Q is the amount of drug released after time t per unit area, D is the diffusion coefficient of drug in the matrix, A is the total amount of drug in the matrix per unit volume, and  $C_s$  is the solubility of drug in the matrix.

Type of matrix diffusion and development in general do not follow the Fickian diffusion mechanism. The simple equation that can describe the behavior of drug release from hydrophilic matrix systems are:

$$\frac{Mt}{M_\infty} = kt^n \dots\dots\dots (5)$$

$Mt / M_\infty$  is the fraction of drug release in time t, k is a constant and n is the exponential diffusion which indicatives of the mechanism of drug release (Wise, 2000).

## **METHODS**

### **Materials**

Ibuprofen (USP., BP., Hubei Biocause Helan Pharmaceutical CO., LTD.), *Xanthan gum* (degussa Texturant Systems, France), *Locust bean gum* (degussa Texturant Systems, France), Avicel® PH 101 (Asahi Kasei Chemical Corp., Tokyo, Japan), Polivinilpirolidon K-30 (ISP Technologies Inc., Wayne, New Jersey), Magnesium stearat (Peter Greven CV, Nederland), talk (Haichen, China), Natrium hidoksida (pa, Merck), Kalium dihidrogen fosfat (pa, Merck).

### **Equipments**

Single punch tableting machine (Type TDP, Shanghai, China), analytical balance (Sartorius), hardness tester (Schleuniger Hardness Tester, tipe 6D-30, Germany), friabilation tester (Erweka, type TA-3, Germany), dissolution tester (Erweka, type DT70, Germany), fluid bed dryer (Glatt, Germany), spectrophotometer UV-VIS (Hitachi, type U-1100, Japan).

### Tablet Manufacturing Method

Ibuprofen tablets made from a variety of formulas. The difference formula based on different combinations of the matrix. Combination matrix used is xanthan gum and locust bean gum. The concentration used for each combination were: xanthan gum, locust bean gum = 1: ½: 1:1, 1:1 ½. Composition formula ibuprofen tablets can be seen in Table I.

Table I. Composition (mg per tablet) of Sustained Release Tablet Ibuprofen (600 mg)

Ingredients	Amount (mg/tablet)		
	Formula I	Formula II	Formula III
Ibuprofen	400	400	400
Xanthan Gum	12.5	12.5	12.5
Locust Bean Gum	6.25	12.5	18.75
Avicel® PH 101	91.25	85	78.75
PVP K-30	30	30	30
Mg Stearat	6	6	6
Talk	54	54	54
Weight of one tablet	600	600	600

Formula I : Xanthan gum : Locust bean gum = 1 : ½

Formula II : Xanthan gum : Locust bean gum = 1 : 1

Formula III : Xanthan gum : Locust bean gum = 1 : 1½

The combination of matrix and Avicel® PH 101 is mixed until homogeneous and added water to form granules, then sifted with a sieve mesh 16. The granules are dried using a fluid bed dryer at 55 ° C for 30 minutes. Sifted dry granules by sieve mesh 18. Dried granules are mixed with ibuprofen and added to a solution of polyvinylpyrrolidone to form the mass of granules, and sieved with a sieve mesh 16. The granules are dried by fluid bed dryer at a temperature of 55 ° C for 30 minutes, then sieved with a sieve mesh 20. Added magnesium stearate and talk, then mixed and homogeneous granules were observed physical properties. Then 600 mg of the prepared granules was compressed using single punch tablet machine with the same hardness was 17-18 Kp.

### Characterization of Granules and Tablets

The flow characteristics, bulk and tapped density of granules were measured to verify the improvement in physical characteristics of granules for tableting. Physical properties of tablets include tablet weight uniformity, tablet hardness, tablet friability and the disintegration time of tablet, other than that conducted the assay of ibuprofen tablets.

### **Drug Release Testing**

The dissolution test was carried out using a dissolution apparatus II method USP over a 6-h period. The dissolution media was phosphate buffer pH 7.2. The temperature of medium and the rotation speed of the paddle were maintained at 37 °C (±0.5 °C) and 100 r/min, respectively. Six tablets were used for each test. Samples (at 5 mL) were withdrawn and replaced with 5 mL fresh phosphate buffer according to following time scheme: 30, 60, 90, 120, 180, 240, 300, and 360 min. The samples were filtered using 0.45 µm Milipore filter. The filtered sample was measured spectrophotometrically at 265 nm to determine the amount of released drug.

### **RESULTS AND DISCUSSION**

Table II shows that bulk density has a smaller value than the value of tapped density and has a significant difference between bulk density and tapped density. This indicates that these granules have large void between particles. From the data of Carr's index indicated that all formulas have a good flow properties because of Carr's index gained less than 20% and Hausner ratio values less than 1.25. Thus a mixture of granular flow is expected to be free flowing and to give homogeneity of content of ibuprofen in tablets.

Table II. Physical Properties of Granules

<b>Formula</b>	<b>Bulk density (g/mL)</b>	<b>Tapped density (g/mL)</b>	<b>Carr's Index (percent)</b>	<b>Hausner Ratio</b>
I	0.34	0.38	12	1.12
II	0.35	0.39	11	1.11
III	0.32	0.36	12	1.13

Percent friability of all formulas to meet the requirements of less than 0.8% showing resistance to the effects of tablet manufacturing process. Physical strength of the exterior of the tablet in the fight against mechanical shock is what determines the size of the fragility.

Table III. Physical Properties of Tablet

<b>Formula</b>	<b>Tablet Hardness (Kp)</b>	<b>Tablet Friability (percent)</b>	<b>Disintegrati on Time (min)</b>	<b>Content Uniformity (percent)</b>	<b>cv of Content Uniformity (percent)</b>
I	17.7	0.1	117	94.8	0.41
II	18.1	0.1	170	85.9	2.87
III	17.5	0.2	150	96.0	0.78

The disintegration time of all formulas using a matrix formula in which a combination of xanthan gum and locust bean gum is affected by formulation. With the same relative hardness tablet disintegration time obtained varies. The longest disintegration time of the formula II, this is probably due to the ratio of xanthan gum and locust bean gum = 1:1 causing the internal structure of the tablet will have a smaller porosity which causes the water will be difficult to penetrate and increase the disintegration time of tablets.

Table III shows that the content of ibuprofen in tablets of each formula has a cv of less than 5%, this means that content of ibuprofen in each tablet in each formula is uniform. This content uniformity data illustrate that tableting process goes well and produces tablets with active substances which approached the ideal levels.

Dissolution profiles was obtained by plotting the percent drug released *vs.* time as shown in Figs.1. The profile of drug released showed that the formula III has the percent of drug release of the most high despite the destruction of much longer time when compared with formula I. This is because formula III shows the mikrogranular while the formula I are makrogranular. While the profile of drug release on the formula II shows the release of the longest when compared with other formulas.




Figure 1. Release profile of ibuprofen sustained release tablet

Table IV. Mathematical Modeling and Drug Release Kinetics of Ibuprofen Sustained Release Tablet

Formula	Zero Order		Higuchi	First Order	Exponential Diffusion (n)
	k (mg/min)	r	r	r	
I	0.5190	0.9991	0.9922	0.9975	0.873
II	0.2679	0.9980	0.9933	0.9993	0.603
III	0.8130	0.9919	0.9830	0.9447	0.570



Table IV shows that both diffusion and erosion mechanisms applicable to the release of ibuprofen from matrix xanthan gum-locust bean gum in medium phosphate buffer pH 7.2, with the dominant mechanism is erosion. Characteristics of drug release, represented by the value of  $n$ . All formulas have the value of  $n$  greater than 0.45 and smaller than 0.89 means that the release characteristics of ibuprofen following the anomalous diffusion transport mechanism, where the rate of diffusion and polymer relaxation are comparable.  $N$  value above also indicates that the controlled drug release more than one process.

## **CONCLUSION**

Xanthan gum in combination with locust bean gum can be used as a slow release matrix tablet of ibuprofen with the kinetics of drug release tends to zero order and the dominant mechanism is erosion. The mechanism of drug transport that occurs is anomalous diffusion, which states that the controlled drug release more than one process. Levels of locust bean gum is used in combination matrix may affect the rate of dissolution, where the ratio of 1:1 between Xanthan gum and locust bean gum provides drug release profile of the longest with the lowest dissolution rate compared to other formulas.

## **REFERENCES**

- Anonim. 2005a. *Water Structure and Behavior : Locust Bean Gum*. Available at: <http://sbu.ac.uk/water/hyloc.html>. Accessed 28 January 2003.
- Anonim. 2005b. *Water Structure and Behavior : Xanthan Gum*. Available at: <http://www.sbu.ac.uk/water/hyxan.html>. Accessed 28 January 2003.
- Ansel, H.C. 1989. *Introduction to Pharmaceutical Dosage Forms*. 4<sup>th</sup> Ed., Lea&Febiger, Philadelphia.
- Baichwal, A, R. 2001. *Sustained Release Heterodisperse Systems-amorphous Drugs*. US Patent 9, 879, 296.
- Higuchi, W.I. 1963. Mechanism of Sustained Action Medication. Theoretical Analysis of Rate of Release of Solid Drugs Disperse in Solid Matrices. *J. Pharm. Sci.*, 52, 1145-1149.
- Kurumaddali, K.R., Ravis, W.R., and Betageri, G.V. 1994. Preparation and Evaluation of Sustained Release Ibuprofen Beads. *Drug Development and Industrial Pharmacy*. 20(17): 2659-2669.
- Tehrani, M.R., and Shobeiri, N. S. 1995. Effect of Various Polymers on formulation of Controlled Release (CR) Ibuprofen Tablets by Fluid Bed Technique,. *Drug Development and Industrial Pharmacy*. 21 (10): 1193-1202.
- Shehab, M. A., and Richards, J. H. 1996. Studies on the In Vitro Release of Ibuprofen from Polyethylene Glycol-Polyvinyl Acetat Mixtures Liquid Filled into Hard Gelatin Capsules. *Drug Development and Industrial Pharmacy*. 22(7): 645-651.
- Wise, D. L.,2000. *Handbook of Pharmaceutical Controlled Release Technology*. Marcel Dekker, Inc., New York.

**ISBN**

ISBN 978-602-95911-1-8



9 786029 591118

**CENTER FOR PHARMACEUTICAL AND MEDICAL TECHNOLOGIES  
DEPUTY FOR AGROINDUSTRY AND BIOTECHNOLOGY  
AGENCY FOR THE ASSESSMENT AND APPLICATION OF TECHNOLOGY  
BPPT Gedung II, Lantai 15  
Jalan M. H. Thamrin No 8, JAKARTA 10340  
Telp. / Fax : (021) 316 9505**

**2010**