



University of Jenderal
Soedirman



National Working Group for
Indonesian Medicinal Plant

INTERNATIONAL CONFERENCE ON MEDICINAL PLANTS 2012

Horison Hotel Purwokerto, 11-13 October 2012

PROCEEDING BOOK



Scientification of Jamu (Evidence-based Jamu Development): A Breakthrough Program from Plant to Medicine for Health Care

The 43rd Meeting of National Working Group
on Indonesia Medicinal Plant
"Exploration, Conservation, Development,
and Utilization of Indonesian Medicinal Plant"

Penerbit :
UNIVERSITAS JENDERAL SOEDIRMAN
PURWOKERTO

PROCEEDINGS OF INTERNATIONAL CONFERENCE ON MEDICINAL PLANTS

Scientification of Jamu (Evidence-based Jamu Development): A Breakthrough Program from
Plant to Medicine for Health Care

In occasion of

The 43th National Meeting of National Working Group on Indonesia

Medicinal Plant

11-13 October 2012

Purwokerto, Indonesia

Editors:

dr. Agung Saprasetya Dwi Laksana, M.Sc., PH.

Heny Ekowati, M.Sc., Apt.

Dr. Agus Nuryanto, M.Si.

Dr. Puji Widodo, M.Sc.

Rehana, M.Si., Apt.

Lay-out:

Sarmoko

Prima Andika Citra

Revo Almando Harris

Published by.

Universitas Jenderal Soedirman

Perpustakaan Nasional RI: Katalog Dalam Terbitan

PROCEEDINGS OF INTERNATIONAL CONFERENCE ON MEDICINAL PLANTS

Scientification of Jamu (Evidence-based Jamu Development): A Breakthrough Program from Plant to Medicine for Health Care

© Universitas Jenderal Soedirman

Cetakan Pertama, 2012

Hak Cipta dilindungi Undang-undang

All Right Reserved

- Editor : dr. Agung Saprasetya Dwi Laksana, M.Sc., PH.
Heny Ekowati, M.Sc., Apt.
Dr. Agus Nuryanto, M.Si.
Dr. Puji Widodo, M.Sc.
Rehana, M.Si., Apt.
- Perancang Sampul : Panitia
- Penata Letak : Sarmoko
Prima Andika Citra
Revo Almando Harris
- Pracetak dan Produksi : Tim UPT. Percetakan dan Penerbitan Unsoed

Penerbit



UNIVERSITAS JENDERAL SOEDIRMAN

Jalan Prof. Dr. H.R. Boenyamin 708 Purwokerto

Kode Pos 53122 Kotak Pos 115

Telepon 635292 (Hunting) 638337, 638795

Faksimile 631802

www.unsoed.ac.id

ISBN: 978-979-9204-80-6

vii + 271 hal, 29.7 cm x 21 cm

Dilarang keras memfotokopi atau memperbanyak sebagian atau seluruh buku ini tanpa seizin tertulis dari penerbit

Committee

Steering committee:

Dr. Hj. Retno Widiastuti, MS

Indah Yuning Prapti, SKM., M.Kes.

Prof. Ir. Totok Agung DH, M.P., Ph.D.

Prof. Suwijiyo Pramono, DEA., Apt.

Prof. Dr. Ir. Suwarto, MS.

Ir. Agus Margiwiyatno, MS., Ph.D

Dr. Bambang Heru Budiarto, MS.

Dr. Ir. Akhmad Sodiq, M.Sc., Agr.

Sugiyono, Ph.D

Tuti Sri Suehesti, M.Sc.

Dr. Lesley Braun

Assoc. Prof. Wandee Gritsanapan

Chariperson : Dra. Warsinah, M.Si., Apt.

Contents

The Effect of Aqueous Extract of <i>Scrophularia Striata</i> on The Growth of <i>Leishmania Major</i> In Vitro and In Vivo Conditions <i>Adolhossein Dalimi, Saeid Soflaei, Fatemeh Ghaffarifar</i>	1
Antifungal Activity and Stability of Long Term Storage of Turmeric Cream Preparations <i>Mukda Jankasem, Wuthi-udomlert, Wandee Gritsanapan</i>	7
Antibacterial Activity of Ethanolic Extract of Mangosteen Skin Fruit (<i>Garcinia mangostana</i>) Against <i>Gardberella vaginalis</i> <i>Fuji Faujiah, Iskandar Sobri, Sunarto</i>	11
Potency of Methanol Extract of <i>Artocarpus chamededen</i> Stem Bark to Inhibit the Growth of <i>Plasmodium falciparum</i> <i>Maximus Taek, Aty Widyawaruyanti Maria Nindatu</i>	17
The Most Four Utilise Medicinal Plants For Anti-Malarial Phytomedicine in Papua and Their Main Chemical Constituents <i>Wahyudi</i>	21
Antifungal Effect of Virgin Coconut Oil (VCO) to <i>Candida albicans</i> : in Vitro Study <i>Thianthy Silviningrum, Fitranto Arjadi</i>	29
Antibacterial Activity Test of Sambiloto Herbs Extracts Against <i>Staphylococcus aureus</i> ATCC 29213 and <i>Escherichia coli</i> ATCC 25922 <i>Yonas B.A., Martha Ervina, Dien Limyati, Lisa Soegianto, Hendrawati</i>	33
The Effect of <i>Hibiscus sabdariffa</i> Linn. Calyx Water Extract on Increasing of Lymphocyte Proliferation In Vitro <i>Nurkhasanah, Puti Ratna Puri</i>	37
Free Radical Scavenging and Antifungal Activities of Juice and Ethanolic Extracts Of <i>Garcinia mangostana</i> L Leaves <i>Elza Sundhani, Diniatik, Muhammad Taufiq Nugroho</i>	41
Antioxidant Activity by DPPH Method and Total Phenolic Content of Water Fraction of Methanolic Extract of <i>Piper betle</i> L. Leaf <i>C.J. Soegihardjo, Rollando</i>	47
In vitro antioxidant activity of <i>Amomum cardamomum</i> leaves and stem <i>Hery Winarsi, Nurtjahjo D. Sasongko, Agus Purwanto, I Gede Arinton, Indah Nuraeni</i>	53
Antioxidant Activity of Extracts and Fractions of Sambiloto Herbs and Salam Leaves <i>Hendra Kurniawan, Lannie Hadisoewignyo, Martha Ervina, Lisa Soegianto</i>	60
Xanthone Derivate Compound Of Mangosteen (<i>Garcinia mangostana</i>) in Suppressing VEGF Expression of SP-C1 Tongue Cancer Cell Line <i>Arlette Setiawan, Roosje R. Owen, Supriatno, Willyanti Soewondo, Unang Supratman, HR Sidik</i>	65
Potency Extract of <i>Morus alba</i> L. As Chemoprevention Agent in Vitro and in Silico <i>Sarmoko, Nia K. Sholihat, Revo A. Harris, Guruh Pratomo, Desiainy S. Sahara, Firster Nugroho</i>	71
Cytotoxic Effects of Methanol Extracts of Soursop (<i>Annona muricata</i>) Leaves on MCF-7 Cell Line and Its Effect on Expression of Bcl-2 Protein <i>Annisa Nur Jannah, Eka Prasasti Nur Rachmani, Iskandar Sobri</i>	76
Identification and Cytotoxic Activity from n-Hexane Fraction of Red Mangrove (<i>Bruguiera gymnorhiza</i>) Against to HeLa Cells <i>Warsinah, Sismindari, Ratna Asmah Susidarti</i>	81
Petroleum Ether, Ethyl Acetate, and Methanol Extracts of <i>Pseudocalymma allicium</i> (Lam.) Sandwith Leaves and Their Antiviral Activities Against Newcastle Disease Virus <i>Nuning Rahmawati, M. Kuswandi, Ratna Asmah Susidarti</i>	86
Hepatoprotective Effect of Curcumin and Piperine against Iron Overload in Rats <i>Heny Ekowati, Sarmoko, Joko Setyono</i>	89
Effect of Green Tea on Hematological Profile in Lead-Exposed Rat <i>Hernayanti, Sukarti Moeljopawiro, Ahmad Hamim Sadewa, Bambang Hariono, Subagus Wahyuono</i>	93

Potential Combination of Temulawak (<i>Curcuma xanthorrhiza</i>), Turmeric (<i>Curcuma domestica</i>) and Red Ginger (<i>Zingiber officinale</i> cv. Rubrum) As Preventive of Nephrotoxicity Due to the Use of Doxorubicin <i>Soraya Diliwiyani, Ulya Rahmawati, Ayu Mayungsari, Retna Pancawati, Dina Arumi, Sarmoko</i>	97
Combination Extract <i>Curcuma domestica</i> , <i>Zingiber officinale</i> cv. Rubrum and <i>Curcuma xanthorrhiza</i> As Anticardiototoxicity of Doxorubicin <i>Yohan Budi Alim, Wulan P. Rahayu, Agustin Damayanti, Ning Uswiyatun, Rika Triyanaputri, Heny Ekowati</i>	101
Protective Effect of <i>Curcuma xanthorrhiza</i> , <i>Curcuma domestica</i> , and <i>Zingiber officinale</i> Combination Against Doxorubicin: A Hematological Study <i>Revo Almando Harris, Sarmoko, Heny Ekowati</i>	106
Potential Ethanol Extracts of Red Fruit Guava (<i>Psidium guajava</i>) on Acute Toxicity Rat Liver Tissue that Induced Glucose <i>Kristianto, Anisyah</i>	111
Safety Evaluation of Slimming Herbal X In Liver Function of Female Wistar Rat <i>Giva Oliviana Yudhista, Hanif Nasiatul Baroroh, Vitis Vini Fera</i>	118
Acute and Subchronic Toxicity Study of Extract Secang (<i>Caesalpinia sappan</i> L) <i>Agus Triyono, Sari Haryati</i>	127
Effect of Antidiabetic Oleum Azadirachtac Semen Using Glucose Tolerance Test in Albino Male Rats <i>Monica Widayawati Setyawan, Irwan Setyabudi, Yohanes Hermawan Widjaya</i>	131
Bioactivity Study of the Larvae of Myrmelcontidae Species (Myrmelcon sp.) As an Oral Hypoglycemic Agent <i>Rahajeng P. Aryani, Sherliyana Novita, Nadia Ayunita, Harwoko</i>	138
Naphtoquinone Derivativesas Potential Inhibitors of α -Glucosidase Activity <i>Mutia Hardhiyuna, Rilianawati, Firdayani, Susi Kusumaningrum, Siska Andrina K.</i>	142
Effect Of Hydroxy Group Addition on 1,4 Naphtoquinon to Inhibit Alpha Glycoside Enzyme <i>Siska Andrina Kusumastuti, Rilianawati, Mutia Hardiyuna, Susi Kusumaningrum, Firdayani</i>	147
Diuresis Effect Comparison Between Ethanolic Leaf Extract of <i>Carica Papaya</i> L. with Hydroclorotiazid to Male White Rat <i>Ratih Febrina Puspitasari, Rendy Primananda Zilmi, Samigun</i>	153
Evaluation of Formulation of Antidiabetic Herbal Tablet of <i>Andrographis paniculata</i> Ness <i>Endang Darmawan, Sri Mulyaningsih</i>	157
Formulation and Evaluation Metformin Hydrochloride Beads Using Matrix of Chitosan-Sodium Alginate Crosslinkage <i>Dhadhang Wahyu Kurniawan, Vitis Vini Fera, Prisci Permanasari</i>	162
Formulation and Evaluation of Preparations Lotion Antioxidant Ethanolic Extract of Soursop Leaves (<i>Annona muricata</i> L.) <i>Shinta Angresti, Iskandar Sobri, Tuti Sri Suhesti</i>	169
Formulation of Enteric Coated Tablets Containing Pineapple (<i>Ananas Comosus</i> Merr.) Fruit Juice <i>Teguh Widodo, Alisyahbana, Ferry Irawan</i>	177
Influence of Various Concentrations of PVP K-30 As a Binder on The Physical Quality of Cerme (<i>Phyllanthus acidus</i>) Leaves Extract Tablet <i>Kuncoro Foe, Teguh Widodo, Juleiny Selfiani</i>	184
Standardization and Formulation of Antipyretic Tablet of Holy Basil Leaves Extract (<i>Ocimum sanctum</i> L.) <i>Farida Lanawati Darsono, Yuliani, Sri Harti</i>	190
Effectiveness of Antifungi Galanga Essential Oil (<i>Alpinia galanga</i>) Preparations in Ointment <i>Tri Ayu Apriyani, Banu Aji Wijayanto, Ardhea Fredyantara, Testy Dwi Septiandini, Sarmoko</i>	197
Potential and Analysis Larvacide Fraction of Interest on Rumput Laut (<i>Eryngium foetidum</i>) Against Mosquito Larva <i>Aedes aegypti</i> <i>Budi Untari, Herlina</i>	202
Application of High Performance Liquid Chromatography in the Determination of Antifungal Drug in Samples <i>Dadan Hermawan, Nur Atiqah binti Abu Bakar, Wan Aini Wan Ibrahim, Mohd Marsin Sanagi</i>	210
Separation and Enrichment of Triterpenoid and Flavonoid from <i>Centella asiatica</i> Leaf Extracts <i>Harwoko, Esy Nansy, Suwijiyono Pramono</i>	219

Isolation and Identification of Triterpenoid Glycoside of Ethanolic Extract of <i>Centella asiatica</i> Herbs by Using Thin Layer Chromatography and High Performance Liquid Chromatography <i>Meli Anita Sari, Diniatik, Indri Hapsari</i>	223
Efficacy Herbs Tempuyung and Kejibeling as Kidney Stones Reduced (preclinical testing on experimental animals white male rats of <i>Sprague dawley</i> strain with kidney stones /calcium oxalate) <i>Pertamawati, Sriningsih, Nuralih, Julham Efendi, Fachry Fachrudin</i>	230
Effect of Cerme (<i>Phyllanthus acidus</i> (L.) Skeels) Leaves Extract on the Decrease of Body Weight and Epididymal Fat As Well As Changes of Triglyceride Levels in Albino Male Rat <i>Idajani Hadinoto, Ening Pujiharsih</i>	238
<i>Ziziphus mauritiana</i> Seed Extract Potentiates Pentobarbital- Induced Sleep in Mice <i>Aye Moh Moh San, Suchitra Thongpranditchote, Wandee Gritsanapan</i>	246
Hair Growth Activity of Ethanolic Extract of Bitter Gourd (<i>Momordica charantia</i>) Leaves <i>Ratnawati, Hanif Nasiatul Baroroh, Iskandar Sobri</i>	250
Hair Growth Activity of Petroleum Ether Extract of <i>Hibiscus rosa-sinensis</i> Leaves Gel <i>Sabrina Nur Fitriisca, Hanif Nasiatul Baroroh, Dhadhang Kurniawan</i>	256
Correlation between Nutritional Status and Depressive Tendencies on The Medical Students Faculty of Medicine and Health Sciences Jenderal Soedirman University <i>Diah Krisnansari, Zaenuri Syamsu Hidayat, Redho Afriando</i>	262
The Effect of Light and Water Stress Intensity on the Growth and Reserpine Content of Pule Pandak (<i>Rauvolfia serpentina</i> Benth.) <i>Edi Purwanto, Samanhuji, Hadiwiyono, Yovita C. Haryza</i>	266

Antioxidant Activity of Extracts and Fractions of Sambiloto Herbs and Salam Leaves

Lannie Hadisoewignyo*, Martha Ervina, Lisa Soegianto, Hendra Kurniawan

*Faculty of Pharmacy, Widya Mandala Catholic University Surabaya
Dinoyo 42-44 Surabaya, Indonesia, telp 6231-5678478 ext. 179,
fax. 6231-5630169, email: lanhadi@yahoo.co

Abstract

Antioxidants are substance which able to scavenge free radicals. Sambiloto herbs (*Andrographis paniculata* Nees) and salam leaves (*Syzygium polyanthum*) extracts have potential of antioxidant activity, so the combination is expected not only reduce blood sugar levels, but also prevent further complications of chronic diseases. In the case of chronic hyperglycemia will increase oxidative stress which is resulting in a reduced number of glucose transporters. This mechanism will increase insulin resistance, lack of insulin signaling and impair insulin secretion by pancreatic β cells. The aims of this research was to determine antioxidant activity of extracts and fractions of sambiloto herbs and salam leaves. As a screen by TLC autography method of extracts and fractions were shown that aqueous fraction of the leaves and herbs ethanol extract of sambiloto and salam have strongest scavenger capacity of DPPH on Rf 0.58; 0.73 and 0.84 for sambiloto aqueous fraction; while salam aqueous fraction on Rf 0.21; 0.34; 0.40; 0.44 and quercetin as antioxidant reference on Rf 0.86. Quantitative antioxidant capacity of aqueous fraction of the extracts were assessed by inhibition of DPPH (1,1-diphenyl-2-picryl-hydrazyl) free radical scavenging method which assay by spectrophotometry at wavelenght 515 nm. Antioxidant capacity of the fractions was showed by IC_{50} . As result was measured that IC_{50} sambiloto herbs aqueous fraction was 46.15 $\mu\text{g/mL}$; while IC_{50} salam leaves aqueous fraction was 18.74 $\mu\text{g/mL}$. These results was suggested that the antioxidant capacity of aqueous fraction of sambiloto herbs and salam leaves higher than its extracts and it might be have therapeutic value in prevent advanced complication in chronic diseases.

Keywords: sambiloto herbs, salam leaves, antioxidant capacity, DPPH

Background

There have been many studies conducted to determine the effects of antidiabetic and antioxidant extracts of sambiloto herbs. Infusion of sambiloto herbs 20% were given a dose of 12.5 mL/kg and 37.5 mL/kg orally in rabbits, hypoglycemic effect at both doses, for comparison used tolbutamide 50 mg /kg (Puslitbang, Jakarta). Research conducted by Soetarno *et al.*, 1999, stating that the use of sambiloto herbs in traditional form, with crude brew sambiloto herbs with water, is appropriate, as the sambiloto herb extracts obtained with organic solvents showed no antidiabetic effects.

Dandu and Inamdar, 2009, conducted a study to determine the antioxidant properties of water extract of sambiloto herbs on mice made diabetic with streptozotocin induced. His study states that a dose of 400 mg /kg, administered by mouth, significantly lowers blood glucose levels and increases the activity of superoxide dismutase (SOD) and catalase enzymes. Hypothetically sambiloto herb can lower blood glucose levels through antioxidant properties. Borhanuddin *et al.*, 1994, noted that there is a significant hypoglycemic effect of aqueous extract of sambiloto herb at a dose of 10 mg /kg, in a study conducted in rabbits. Verma and Vinayak, 2007, stating that the water extract of sambiloto herbs have antioxidant defense system in lymphoma in the liver of mice. Tripathi and Kamat, 2007, stating that the water extract of sambiloto herbs showed the potent of effects anti-radical, conducted on rat liver.

While research on salam leaf extract states that salam leaf extract has antidiabetic and antioxidant effects. Studiawan and Santoso, 2005, stated in his research that extract ethanol salam leaves can lower blood glucose alloxan induced mice, at doses of 2.62 and 5.24 mg/20 g body weigh. Lelono *et al.*, 2009, stating that the methanol-water extract of salam leaf has free radical scavenging activity was high, with ED50 0.18 mg /mL, and protection from the effects of β -carotene was 85.7% at a concentration of 100 $\mu\text{g/mL}$.

Sambiloto herbs have a working mechanism accelerates the release of glucose by increasing metabolism, while the salam leaves can serve as an astringent that can inhibit glucose intake. Seeing both the mechanism of action of these two natural ingredients, it is possible to combine of to obtain a synergistic effect. According to research conducted by Widjajakusuma *et al.*, 2009, the specific combinations and doses of water extract of sambiloto and salam leaves can provide a synergistic effect as antidiabetic, and has a better effect in lowering blood sugar than the use of a single extract, as well as to metformin.

Based on research conducted by Widjajakusuma et al, 2009, the purpose of this study was to investigate the antioxidant activity of extract combination of sambiloto herbs and salam leaf that have been shown to have antidiabetic effects.

Methods

Equipments: percolator, rotavab, UV-VIS spectrophotometer, UV lamps, TLC plates, and filter paper.

Materials: sambiloto herbs and salam leaves obtained from Materia Medika Indonesia, DPPH (1,1-diphenyl-2-picrylhydrazyl), methanol, n-hexane, ethyl acetate, 96% ethanol, chloroform, and quersetin.

Standardization of simplisia including the determination of ash content, the assay of water extrateable, and assay of ethanol extractable (Anonim, 1980). Preparation of extract were done by maceration. At the end of the extraction of total extract obtained from sambiloto herbs and salam leaves. Standardization of extracts had been conducted were moisture content, ash content, extract yield, and chromatogram profiles of active substances (Anonim, 1980).

Preparation of crude plant extract

The dried plant materials were powdered using a grinder. The extraction was done at room temperature. About 5 kg of dried, ground plant materials were soaked in methanol (37.5 L) for 24 h while stirring occasionally. The final extracts were passed through Whatman filter paper (maserat I) and the residue were soaked with 96% ethanol (20 L), and after 24 hours, the ethanol extract filtered (maserat II). Maserat I and II combined and concentrated by using a evaporator, followed by heating in a porcelain dish on a water bath until a thick extract. The viscous extract weighed and then diluted with 90% methanol-water as much as 100 mL. After that, a gradual fractionation in a separating funnel with 100 mL of n-hexane in a way shaken for 5 minutes and separated fractions of n-hexane. Fractionation done several times until the n-hexane fraction colorless again. All the n-hexane fraction obtained put together and concentrated in the evaporator. Methanol-water fraction was evaporated in a porcelain dish over a water bath until methanol evaporated and the fraction of water. Do fractionation of water by using ethyl acetate in a separating funnel as the fraction of n-hexane. All ethyl acetate fraction obtained and collected into one concentrated in the evaporator. Water fraction derived from the fractionation of the ethyl acetate remainder concentrated in porcelain dish over a water bath.

At the end of the extract obtained by extraction of the total, the fraction of hexane, ethyl acetate fraction and water fraction of sambiloto herbs and salam leaves.

In-Vitro determination of antioxidant capacity with DPPH (1,1-diphenyl-2-picrylhydrazyl) (Cavin, dkk., 1988)

a. TLC Autography technique

Each fractions were impregnated with the mobile phase n-hexane: ethyl acetate (9: 1); methanol: chloroform (1:2), and butanol: acetic acid: water (3:1:1) , which is then saturated in each chamber. Further stages as made stock solution of each fractions test extract: 20 mg dissolved in 5 ml methanol pa. After spotting the extracts on the TLC plates, the spot sprayed with DPPH 1% and observed the formation of yellow on spot stain solution is then compared with the benchmark being used.

b. Determination of Inhibition Concentration (IC₅₀)

On the quantitative determination of antioxidant activity, determined IC₅₀ values, with the following stages of work: preparing the test solution of the water fraction of extract with a concentration of the mother liquor (10000 ppm) and 0.13% DPPH solution. Created 5 different concentration of the mother liquor, pipetted 85 µL DPPH solution and add 3 mL of methanol pa 0.13% and then shaken and allowed to stand for 30 minutes; then added with methanol, and determined the maximum absorption wavelength and absorbance examined. Each test preparation was taken as much as 3 ml, placed in vials, 85 µL so lution of 0.13%, DPPH mixed and allowed to stand for 30 minutes, after which the preparation was observed absorbance at a wavelength of maximum absorption. IC₅₀ values, calculated from a linear regression curve between% inhibition of uptake versus In concentration of the extract, as in equation (1).

$$\% \text{ inhibition DPPH activity} = (A \text{ DPPH} - A \text{ sampel}) / A \text{ DPPH} \times 100\% \dots\dots\dots (1)$$

IC₅₀ is a number that indicates the concentration of extract (mg /mL) were able to inhibit 50% oxidation. The smaller the IC₅₀ values higher antioxidant activity. An antioxidant compound is said to be very powerful if IC₅₀ values less than 50, a strong (50-100), moderate (100-150), and weak (151-200).

Results

The results for the determination of ash content of sambiloto herbs and salam leaves in the form of crude extract, can be seen in Table 1.

Table 1. Standarization of Simplisia of Sambiloto Herbs and Salam Leaves (Anonim, 1978)

No.	Parameter test	Requirements	Test Result	
			Sambiloto Herbs	Salam Leaves
1	Ash content	< 5%	11,13 ± 0,35	5,69 ± 0,09
2	Assay of water extractable	> 15,6%	17,03 ± 0,38	5,45 ± 0,24
3	Assay of ethanol extractable	> 4,3%	13,53 ± 0,10	11,55 ± 0,03

Preparation of extract of sambiloto herbs and salam leaves using maceration method. The final result is obtained that the sambiloto herbs extract yield as much as 245 g (24.50%), while the salam leaves extract as much as 141.3 g (14.15%) yield. Extract and fractions were screened using silica gel stationary phase and mobile phase butanol: acetic acid: water = 3: 1: 1, to determine the class of active compounds from the extracts and the fractions.

TLC Autography technique

Antioxidant activity assays carried to all factions, with KLT autography, using a quersetin as a standard. The TLC results contained in Figure 2 and Figure 3. Based on test free radical activity in TLC autography, it is known that which has the greatest antioxidant activity, both sambiloto herbs and salam leaves is the fraction of water.




Figure 1. TLC profile of extract and fraction of sambiloto herbs and salam leaves with stationary phase is silica gel and mobile phase is Diam butanol :acetic acid : water = 3 : 1 : 1, (a) UV 254 nm, (b) UV 366 nm, (c) observations with spot spray vaniline-sulfuric acid

Description: 1= crude extract of sambiloto herbs 5 = crude extract of salam leaves
 2 = n-hexane fraction of sambiloto 6 = n-hexane fraction of salam leaves
 3 = ethyl acetate fraction of sambiloto 7 = ethyl acetate fraction of salam leaves
 4 = water fraction of sambiloto 8 = water extract of salam leaves
 9 = rutine




Figure 2. TLC profile of antioxidant of sambiloto herbs with the stationary phase is silica gel and mobile phase is butanol: acetic acid: water = 3: 1: 1
Description: (1) crude extract, (2) the fraction of non-polar n-hexane, (3) semi-polar ethyl acetate fraction, (4) the fraction of water, (5) quersetin




Figure 3. TLC profile of antioxidant of salam leaves with the stationary phase is silica gel and mobile phase is butanol: acetic acid: water = 3: 1: 1
Description: (1) crude extract, (2) the fraction of non-polar n-hexane, (3) semi-polar ethyl acetate fraction, (4) the fraction of water, (5) quersetin

Determination of Inhibition Concentration (IC₅₀)

The qualitative antioxidants activities had been test by determining the IC₅₀ values (Table 3).

Table 3. IC₅₀ values of Extracts and their Fractions

Sample	IC ₅₀ (µg/mL)
DPPH 0,13% + metanol p.a	-
Water fraction of Sambiloto herbs	46.15 ± 10.46
Water fraction of Salam leaves	18.74 ± 0.25
Water fraction of salam leaves : sambiloto herbs = 1 : 2	15.78 ± 1.14
Water fraction of salam leaves : sambiloto herbs = 2 : 1	30.02 ± 3.22
Water fraction of salam leaves : sambiloto herbs = 1 : 6	37.60 ± 1.37
Water fraction of salam leaves : sambiloto herbs = 6 : 1	29.09 ± 3.08
Quercetin	7.00 ± 0.12

Discussion

Sambiloto herbs and salam leaves in the form of extract and fraction both have antioxidant activity, based on TLC autography assay and IC₅₀ determination. Salam leaves has a greater antioxidant activity (18.74 µg/mL) than sambiloto (46.15 µg/mL) based on IC₅₀ values (Table 3). Combination with the higher concentrations of salam leaves indicates greater antioxidant activity, but when combined with sambiloto herbs the antioxidant activity will increase up to a certain value ratio (in this study up to 1:2), while increasing the sambiloto herbs concentrations up to six times compared to the salam leaves showed a decrease in antioxidant activity.

Conclusion

Antioxidant capacity of the fractions was showed by IC₅₀. As result was measured that IC₅₀ sambiloto herbs aqueous fraction was 46.15 µg/mL; while IC₅₀ salam leaves aqueous fraction was 18.74 µg/mL. These results was suggested that the antioxidant capacity of aqueous fraction of sambiloto herbs and salam leaves higher than its extracts and it might be have therapeutic value in prevent advanced complication in chronic diseases.

References

- Anonim, 1980, *Materia Medika Indonesia*, Jilid IV, Departemen Kesehatan Republik Indonesia, Jakarta.
- Borhanuddin, M., Shamsuzzoha, M., dan Hussain, A.H., 1994, Hypoglycaemic Effects of *Andrographis paniculata* Nees on Non-Diabetic Rabbits, *Bangladesh Med Res Counc Bull*, 20, 24-26.
- Dandu, A.M, dan Inamdar, N.M., 2009, Evaluation of Beneficial Effects Antioxidant Properties of Aqueous Leaf Extract of *Andrographis paniculata* in Stz-Induced Diabetes, *Pak.J.Pharm,Sci*, 22(1), 49-52.
- Lelono, R.A.A., Tachibana, S., Itoh, K., 2009, In Vitro Antioxidant Activities and Polyphenol content of *Eugenia polyantha* Wight in Indonesia, *Pakistan Journal of Biological Sciences*, 12(24), 1564-1570.
- Puslitbang Farmasi, Badan Litbang Kesehatan, Tinjauan Hasil Penelitian Tanaman Obat di Berbagai Industri (1), Jakarta.
- Soetarno S., Sukandar, E.Y., Sukrasno, dan Yuwono, A., 1999, Aktivitas Hipoglikemik Ekstrak Herba Sambiloto (*Andrographis paniculata* Nees, Acanthaceae), *JMS*, 4 (2), 62-69.
- Studiawan, H, dan Santosa, M.H., 2005, Uji Aktivitas Penurunan Kadar Glukosa Darah Ekstrak Daun *Eugenia Polyantha* pada Mencit yang Diinduksi Aloksan, *Media Kedokteran Hewan*, 21(2), 62-65.
- Tripathi, R, dan Kamat, J.P., 2007, Free Radical Induced Damages to Rat Liver Subcellular Organelles: Inhibition by *Andrographis paniculata* Extract, *Indian J Exp Biol*, 45, 959-967.
- Verma, N, dan Vinayak, M., 2007, Antioxidant Action of *Andrographis paniculata* on Lymphoma, *Mol Biol Rep*.
- Widjajakusuma, E.C., Hendriati, L, Ferawati, dan Surjadhana, A., 2009, Uji Toksisitas dan Farmakologi *Andrographis paniculata* dan Antikanker dari *Centella asiatica*, Laporan Penelitian PPOT, Universitas Katolik Widya Mandala, 2009.