

← → ↻ <https://mail.google.com/mail/u/1/#search/editor%40ejpam.com/FMfcgzGrfBkdmHqpVFDwtgmPKwVSH> ☆

editor@ejpam.com

Active

Google

Compose

Inbox 132

Starred

Snoozed

Sent

Drafts 7

More

Labels +

[EJPAM] Submission Acknowledgement External Inbox x

2 of 5

Editorial Office, EJPAM <editor@ejpam.com> to me

Sun, Jan 22, 10:29 PM

Dear Gregoria Ariyanti,

Thank you for submitting the manuscript, "The Discrete Lyapunov Equation of The Orthogonal Matrix in Semiring" to European Journal of Pure and Applied Mathematics. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URI: <https://ejpam.com/index.php/iejpam/author/3aashboard/submission/4712>
 Manuscript ID: 4712
 Username: gregoria

You might suggest at least 5 potential reviewers including their names, emails, departments and universities if you have not submitted yet. You can send/submit the potential reviewer list by email

This will certainly make the review process faster.

NOTE: We have a fast track review program which is not compulsory, and an optional program to accelerate the editorial and review processes. In order to take your manuscript into the fast track review program you should pay Fast Track Review Fee of 400USD with the link provided below.

Fast track review fee link: <https://buy.stripe.com/dR64jb6Zcds0M05K>

After your payment you should contact the editorial office.

If you have any questions, please contact us.

Editorial Office, EJPAM

← → ↻ <https://mail.google.com/mail/u/1/#search/eycetin%40gmail.com/FMfcgzGrJgDKXlZnfwMoxGmQWTDs> ☆

eycetin@gmail.com

Active

Google

Compose

Inbox 132

Starred

Snoozed

Sent

Drafts 7

More

Labels +

Eyup Cetin
Editor, EJPAM

3 of 3

Review #1 (also attached)

The results presented in the article is new.

Nevertheless, because in conclusion it is written "Furthermore, this condition can be used to find other characteristics of a linear system over a semiring", so an example or explanation need to be added.

Examples or explanations are added in the results and discussion sections.

Review #2

Regarding the article
The Discrete Lyapunov Equation of The Orthogonal Matrix in Semiring

- 1- The title of the article is well chosen.
- 2- The topic of the article is good and usable by researchers.

But:

- 3- The introductions of the article are too much compared to the obtained results.
- 4- The number of results obtained is very small.

Editor's Suggestion

← → ↻ <https://mail.google.com/mail/u/1/#search/eycetin%40gmail.com/FMfcqzGsmDtkbJKNcdIxChRbfcqQZTh> ☆

☰ Gmail × Active ? ⚙️ ☰ Google 

☰  Compose

☰ Mail  Inbox 132

☰ Chat  Starred

☰ Spaces  Snoozed

☰ Meet  Sent

 Drafts 7

 More

Labels +

← 🗑️ 🕒 📧 📧 📧 📧 📧 1 of 3 < >  

Proof / 9 - 4712 External 

 **Eyüp Çetin** <eycetin@gmail.com>
to me, Barış ▾ 📧 Fri, Apr 21, 2:35 PM ☆ ↶ ⋮

Dear Professor,

I hope this email finds you well.

I have attached a proof copy of your research article prepared for publication by *European Journal of Pure and Applied Mathematics*. Please review the document- as soon as possible- for any final corrections, modifications, or inadvertent copy- editing errors, and inform me of your findings by either annotating the proof PDF directly, or listing any corrections by page number and line number. Please note that the page numbers may be different once the paper is published.

We will consider that you approved the publication of your paper as it is unless you do not reply **within 2 days**.

Also, please be sure that you have completed correctly the information of all authors (orders of authors, affiliations and emails) and of title, keywords and abstract on the online system as in the pdf version of your paper.

I appreciate your prompt attention to this matter; don't hesitate to let me know if you have any questions.

Thank you in advance,
Kind regards,

Eyüp Çetin
Editor, EJPAM

The Discrete Lyapunov Equation of The Orthogonal Matrix in Semiring

Gregoria Ariyanti^{1,*}, Ana Easti Rahayu Maya Sari¹

¹ *Department of Mathematics Education, Widya Mandala Surabaya Catholic University, Indonesia*

Abstract. Semiring is an algebraic structure of $(S, +, \times)$, similar to a ring, but without the need to add a reciprocal to each element. The forms $(S, +)$ and (S, \times) are semigroups that satisfy the distributive law of multiplication and addition. In matrix theory, there is a term known as the Kronecker product. This operation transforms two matrices into a larger matrix containing all possible products of the entries in the two matrices. This Kronecker product has several properties often used to solve the complex problems of linear algebra and its applications. The Kronecker product is related to the Lyapunov equation of a linear system. Based on previous research in the Lyapunov equation in conventional linear algebra, this paper will describe the characteristics of the Lyapunov equation in a semiring linear system in terms of the Kronecker product.

2010 Mathematics Subject Classifications: 16Y60, 34D08, 37M25

Key Words and Phrases: linear system, the Lyapunov equation, semiring, the Kronecker product.

1. Introduction

A non-empty set G with a binary operation $*$ is known as Group if it has the following properties: associative, a zero element of binary operations $*$, and every element, not a zero element, has an inverse. Meanwhile, a non-empty set R with two binary operations, particularly $*$ and \circ is known as Ring if it has the following properties: $(R, *)$ is a commutative group, closed \circ , associative to binary operation \circ , and each binary operations $*$ and \circ is distributive. If the Ring has the following properties: commutative to binary operation \circ , it has a unit element of binary operation \circ , and every element, not a zero element, has an inverse to binary operation \circ , it is called Field. Different algebraic systems will appear if Group and Ring conditions are weakened, particularly Semigrup and Semiring. If a few Group or Ring characteristics are removed, the algebraic structure formed is

*Corresponding author.

Email addresses: gregoria.ariyanti@ukwms.ac.id (G. Ariyanti), ana.easti.rahayu@ukwms.ac.id (A.E.R.M. Sari)

Semigrup, after which Semiring. One of the problems and applications often encountered in mathematics is completing the Linear Equations System.

The Kronecker product is a binary matrix operator that maps two arbitrarily dimensioned matrices into a larger matrix with a particular block structure ([13]). The Kronecker product discussed by Zhou et al. (1996) and Whitcomb (2020) applies to matrices whose elements are real numbers ([4],[12]). The set of real numbers is a field.

The linear equations system that researchers have developed is a system of linear equations over Field, which include real numbers \mathbb{R} or complex numbers \mathbb{C} ([9],[4]). In other studies, the object of research is extended not to Field anymore but to commutative Ring and linear equations system Commutative over ring have been discussed by Brown ([4]). Likewise, assuming an extension from the Ring to Ring commutative does not generally change the definition.

This paper presents the characteristics of the discrete Lyapunov equation of a matrix. And the scope of the topic is in a system of linear equations in semiring in terms of the Kronecker product. First, section 2 will review some basic facts about the semiring, the Kronecker product of the matrix in semiring, and the Lyapunov equation of the matrix in semiring. Then, in Section 3, we show a necessary or sufficient condition of the Lyapunov equation in a linear system over semiring in terms of the Kronecker product.

2. Materials and literature review

2.1. Semiring and Matrices in Semiring

Definition 1. *Semigroup S is an empty set that is equipped with an associative binary $*$ operation, that is $x * (y * z) = (x * y) * z$ for every $x, y, z \in S$.*

Poplin defines a semiring and its properties as follows([3],[10],[9],[12]).

Definition 2. *Semiring is a non-empty set S with two binary operations, addition $(+)$ and multiplication (\times) , which have the following properties: commutative and associative properties of $+$, associative property of \times , distributive property of \times to $+$, the set S has a zero element $0 \in S$ so that $0 + a = a + 0 = a$ and $0 \times a = a \times 0 = 0$ for every $a \in S$. This zero element is called the absorbent element (absorption), and the set S has a unit element e , $e \times a = a \times e = a$ for every $a \in S$.*

The commutative and idempotent characteristics in Group and Ring also apply in Semiring ([10]). Let $M_{n \times 1}(S)$ be the set of all vectors $n \times 1$ with the elements of semiring S . And, let $M_{n \times n}(S)$ be the set of all $n \times n$ matrices with the elements of semiring S ([4],[2]). The $+$ and \times operation for matrices over semiring is defined as in Definition 3.

Definition 3. *Let S semiring, a positive integer n , and $M_n(S)$ is the set of all $n \times n$ matrices over S . For every $A, B \in M_n(S)$, $+$ and \times operations over semiring S are defined $C = A + B$ as $c_{ij} = a_{ij} + b_{ij}$ and $C = A \times B$ as $c_{ij} = \sum_l a_{il} \times b_{lj}$.*

The semiring S has 0 as a zero element and 1 as an identity element, as in the matrix of conventional algebra. We can form a zero matrix and an identity matrix based on the zero element and the identity element ([3]). The zero matrix $n \times n$ over semiring S is 0_n is defined as a matrix with all elements equal to the 0–element, that is $(0_n)_{ij} = 0$. The identity matrix $n \times n$ over S is defined as the matrix with all elements equal to the e –element, that is $[I_n]_{ij} = \begin{cases} e, & \text{if } i = j \\ 0, & \text{if } i \neq j \end{cases}$.

In semiring, the element of semiring has an inverse operation on $+$ to determine a determinant of a matrix in semiring S . Thus, a determinant of a matrix over a semiring S is characterized by a permutation.

2.2. Kronecker Product

Kronecker's product is related to the stack operator. The stack operator maps an $n \times m$ matrix to a $nm \times 1$ vector ([13]). The stack of the $n \times m$ matrix A is represented by $vec(A)$ which is a vector formed by stacking the columns of A on the vector $nm \times 1$.

Example 1. Let A is a matrix with $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then its stack form is

$$vec(A) = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}.$$

If C is an $n \times m$ matrix comprising m column vectors $c_1, c_2, c_3, \dots, c_m$, where each c_i is an $n \times 1$ vector $C = [c_1, c_2, c_3, \dots, c_m]_{n \times m}$, then

$$vec(C) = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix}_{nm \times 1}.$$

Let $X \in M_{m \times n}(S)$, the form $vec(X)$ denote the vector formed by stacking the columns of X into one long vector:

$$vec(X) = [x_{11} \ x_{21} \ \dots \ x_{m1} \ x_{12} \ x_{22} \ \dots \ x_{1n} \ x_{2n} \ \dots \ x_{mn}]^T.$$

Kronecker product is an operation on two matrices that do not require size ([7], [8]). The notation \otimes denotes Kronecker products. With S semiring, let $A \in M_{m \times n}(S)$ and $B \in M_{p \times q}(S)$ then the Kronecker product of A and B is defined as

$$A \otimes B := \begin{pmatrix} a_{11}B & \dots & a_{1n}B \\ \vdots & \ddots & \vdots \\ a_{m1}B & \dots & a_{mn}B \end{pmatrix} \in M_{mp \times nq}(S).$$

Furthermore, if A and B are square matrices with $A \in M_{n \times n}(S)$ and $B \in M_{m \times m}(S)$ then the Kronecker sum of A and B is defined as

$$A \oplus B := (A \otimes I_m + I_n \otimes B) \in M_{nm \times nm}(S).$$

Properties of Kronecker product is given the following theorem ([7], [13]).

Theorem 1. *For $A \in M_{m \times n}(S)$ and $B \in M_{p \times q}(S)$ with S semiring, we have the following properties: the Kronecker product is associative, not in general commutative, i.e., usually $(A \otimes B) \neq (B \otimes A)$, and transpose distributes over the Kronecker product (do not reverse order) $(A \otimes B)^T = A^T \otimes B^T$.*

Let $A \in M_{n \times n}(S)$ and $B \in M_{m \times m}(S)$, and λ_i with $i = 1, 2, \dots, n$ be the eigenvalues of A and μ_j with $j = 1, 2, \dots, m$ be the eigenvalues of B . Then we have the following properties: the eigenvalues of $A \otimes B$ are the mn numbers $\lambda_i \mu_j$ and the eigenvalues of $A \oplus B = (A \otimes I_m) + (I_n \otimes B)$ are the mn numbers $\lambda_i + \mu_j$, with $i = 1, 2, \dots, n, j = 1, 2, \dots, m$.

2.3. Discrete Lyapunov equation

The linear system is closely related to stability, which can be observed using the eigenvalue criterion of matrix A . Furthermore, the stability of the linear system is closely related to the existence of a solution to the Lyapunov equation ([1]). Therefore, this method can determine the stability of the system without knowing the solution to the system. Lyapunov's equation for a linear system over a field was given by Zhou ([6]). In this study, the definition of the Lyapunov equation is given for a discrete linear system on a semiring, adopting the meaning of the Lyapunov equation for a linear system on a plane. The discrete Lyapunov equation for the linear system over semiring is defined as follows ([11], [5]).

Definition 4. *Given a matrix $A, X, Q \in M_n(S)$. The Lyapunov equation for a linear system over a semiring is defined as $AXA^T - X + Q = 0$.*

For linear systems over the field, the existence of solutions to the Lyapunov equations is associated with asymptotic stability. That is, the system is asymptotically stable if a solution to the Lyapunov equation exists. On the other hand, if the system is asymptotically stable, a solution to the Lyapunov equation exists. ([1]).

3. Results and discussion

The problem of discrete Lyapunov equations over semirings is limited to orthogonal matrices. These are because of the limited nature of the semiring.

Theorem 2. *Let S is semiring. Then for any matrices $A \in M_{k \times m}(S)$, $B \in M_{n \times l}(S)$, and $X \in M_{m \times n}(S)$, we have $\text{vec}(AXB) = (B^T \otimes A)\text{vec}(X)$.*

Proof. We have $(AXB)_{.k} = \sum_j b_{jk}AX_{.j} = (b_{1k}A \ b_{2k}A \ \dots \ b_{nk}A)$ and

$$\begin{bmatrix} X_{.1} \\ X_{.2} \\ \vdots \\ X_{.n} \end{bmatrix} = [B_{.k}^T \otimes A] \text{vec}(X) = [(B^T)_{.k}^T \otimes A] \text{vec}(X).$$

Furthermore, we conclude $\text{vec}(AXB) = [B^T \otimes A]\text{vec}(X)$ since the transpose of the k th column of B is the k th row of B^T .

In the following, another property of the Kronecker product for matrix over a semiring is given.

Theorem 3. *Let $A \in M_{m \times m}(S)$, $B \in M_{n \times n}(S)$, and $X \in M_{m \times n}(S)$, we have $\text{vec}(AX + XB) = (B^T \oplus A)\text{vec}(X)$.*

Proof. From the definition of Kronecker sum, we have

$$(B^T \oplus A)\text{vec}(X) = (B^T \otimes I_m + I_n \otimes A)\text{vec}(X) = (B^T \otimes I_m)\text{vec}(X) + (I_n \otimes A)\text{vec}(X).$$

According to Theorem 2, we have

$$(B^T \oplus A)\text{vec}(X) = \text{vec}(I_m XB) + \text{vec}(AX I_n^T) = \text{vec}(XB) + \text{vec}(AX).$$

Due to the limitations of semirings, the existence of discrete solutions to the Lyapunov equations on semirings depends on the orthogonal matrix. This statement is given in the following theorem.

Theorem 4. *Let discrete Lyapunov equation over semiring $AXA^T - X + Q = 0$ with $A \in M_{n \times n}(S)$, $Q \in M_{n \times n}(S)$. For A is orthogonal matrix, there is a unique solution $X \in M_{n \times n}(S)$ if and only if $\lambda_i(A) + \lambda_j(-A) \neq 0$.*

Proof. A matrix A is orthogonal if $A^T = A^{-1}$, hence we have $AXA^T - X + Q = 0$ and then $(AXA^T)A - XA = -QA$. Because A is orthogonal, so we have $AX - XA = -QA$. Therefore, $AX + X(-A) = Q(-A)$ and $\text{vec}(AX + X(-A)) = ((-A)^T \oplus A)\text{vec}(X)$. Finally, we have $((-A)^T \oplus A)\text{vec}(X) = \text{vec}(-QA)$. Based on conditions, we have a unique solution if and only if $(-A)^T \oplus A$ non-singular.



4. Conclusion

We have shown that the solutions of the discrete Lyapunov equations for matrices over semirings are also valid and given in Theorem 3 and Theorem 4. Furthermore, this condition can be used to find other characteristics of a linear system over semiring.

Acknowledgements

We gratefully thank the Directorate of Research and Community Service, Deputy for Research and Development Strengthening Ministry of Education and Culture, Research and Technology, Indonesia.

References

- [1] I Ahmed MJ Ibrahim F Jarad A Ibrahim, SI Bala. Numerical construction of lyapunov functions using homotopy continuation method. *Advances in the Theory of Nonlinear Analysis and its Applications*.
- [2] H Anton and C Rorres. *Elementary Linear Algebra: Applications Version, 11th edition*. John Willey and Sons, The United States of America, 2013.
- [3] G Ariyanti. A Note of The Linear Equation $AX=B$ with Multiplicatively-Reguler Matrix A in Semiring. *International Conference on Applied & Industrial Mathematics and Statistics (ICoAIMS), Journal of Physics: Conference Series*, 2019.
- [4] WC Brown. *Matrices over Commutative Rings*. New York : Marcel Dekker, Inc., 1993.
- [5] AS Hodel and B Tenison. *Linear Algebra and Its Applications*. Elsevier Science Inc., 1996.
- [6] K Glover K Zhou, JC Doyle. *Robust and Optimal Control*. Prentice Hall, New York, 1996.
- [7] S Kathrin. *On the Kronecker Product*. PhD thesis, University of London, 2013.
- [8] CFV Loan. The ubiquitous kronecker product. *Journal of Computational and Applied Mathematics*, 123:85–100, 2000.
- [9] P Narendran. Solving Linear Equations Over Polynomial Semirings. "11 th Annual IEEE Symposium on Logic in Computer Science", IEEE Computer Society Press, Los Alamitos, CA, pages 466–472, 1996.
- [10] PL Poplin. *The Semiring of Multisets*. PhD thesis, Faculty of North Caroline State University, 2000.
- [11] D Palitta S Massei and Robol. Solving rank structured sylvester and lyapunov equations. *SIAM Journal on Matrix Analysis and Applications*.
- [12] A Wasanawichit W Mora and Y Kemprasit. Invertible matrices over idempotent semirings. *Chamcuri Journal of Mathematics*.
- [13] LL Whitcomb. *Notes on Kronecker Products*. Johns Hopkins University, 2020.