



[VIEW ARTICLES](#)

[ISSUES](#)

[ABOUT](#)

[FOR AUTHORS](#)

[FOR REFEREES](#)

EDITORIAL BOARD

Chief Editor

Son Radu

Malaysia

Editor

Vivian New Chia Yeung

Malaysia

Assistant Editor

Kimberley Rinai

Malaysia

Associate Editors

Prem Narayan Paudel

Kathmandu University
Nepal

Associate Editors

Borarin Buntong

Royal University of Agriculture
Cambodia

Associate Editors

Yoshitsugu Nakaguchi

Ishikawa Prefectural University
Japan

Nasreldin Elhadi

University of Dammam
Saudi Arabia

Dwaipayan Sikdar

University of Chittagong
Bangladesh

Rungsinee Sothornvit

Kasetsart University,
Thailand

Nikos Tzortzakis

Cyprus University of Technology

Saisamorn Lumyong

Chiang Mai University

Ekachai Chukeatirote

Mae Fah Luang University

Cyprus

Pin Kar Yong

Forest Research Institute
Malaysia (FRIM)
Malaysia

Chim Chay

University of Agriculture
Cambodia

Le Van Viet Man

Ho Chi Minh City University of
Technology
Viet Nam

Oscar Herrera-Calderon

Universidad Nacional Mayor de
San Marcos
Peru

Hasmadi Mamat

University Malaysia Sabah
Malaysia

Regiane Ribeiro dos Santos

Federal Rural University of Rio de
Janeiro
Brazil

Koh Soo Peng

Malaysian Agricultural Research
and Development Institute
(MARDI)
Malaysia

Dinko Hristov Dinko

Trakia University

Thailand

Hanifah Nuryani Lioe

Bogor Agricultural University
Indonesia

Michelle Galindo de Oliveira

Federal University of
Pernambuco
Brazil

Alper Sener

Onzekiz Mart University
Turkey

Angelo Maria Giuffrè

Mediterranea University of
Reggio Calabria
Italy

Javier Garcia-Lomillo

University of Burgos
Spain

Laura Marcela Zárate Polanco

National University of Colombia-
Bogotá
Colombia

Swapan K. Banerjee

HPFB Health Canada,
Canada

Adel Abu Bakr Abd El-Hamid**Shatta**

Suez Canal University
Egypt

Thailand

Saroat Rawdkuen

Mae Fah Luang University
Thailand

Giovanna Giacalone

Università di Torino
Italy

Lorina Acilo Galvez

Visayas State University
Philippines

Sergio Soto Simental

Universidad Autónoma del
Estado de Hidalgo
Mexico

Fernando Cebola Lidon

Universidade Nova de Lisboa
Portugal

Linda Lim Biaw Leng

Universiti Brunei Darussalar
Brunei Darussalam

J.M. Krishanti Jayarukshi Kur**Premarathne**

Wayamba University of Sri Lar
Sri Lanka

Katherine Ann T. Castillo-Isra

University of the Philippines L
Banos
Philippines

Steve Flint

Bulgaria

Giuseppe Sortino

University of Palermo

Italy

Iddya Karunsagar

Nitte University

India

Massey University

New Zealand

COPYRIGHT © 2016 - 2023 RYNNYE LYAN RESOURCES

[VIEW ARTICLES](#)[ISSUES](#)[ABOUT](#)[FOR AUTHORS](#)[FOR REFEREES](#)

FOOD RESEARCH

Volume 7, Supplementary 1, 2023

International Food Conference 2021

Effects of frozen storage duration on the physicochemical and sensory properties of cassava sticks

Trisnawati, C.Y., Sutedja, A.M. and Kaharso, V.C.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).5](https://doi.org/10.26656/fr.2017.7(S1).5)

The effects of frozen storage duration on the physicochemical and sensory properties of cassava sticks was studied by Trisnawati *et al.*

Re-formulating *Monascus* fermented durian seeds yogurt with strawberry (*Fragaria x ananassa*) puree to enhance its microbiological, physicochemical and organoleptic properties

Kuswardani, I., Yuwono, F., Dharmawan, N.E., Nugerahani, I., Ristiarini, S., Srianta I. and Tewfik, I.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).11](https://doi.org/10.26656/fr.2017.7(S1).11)

Kuswardani *et al.* re-formulated *Monascus* fermented durian seeds yogurt with strawberry (*Fragaria x ananassa*) puree for better enhancement of its microbiological, physicochemical and organoleptic properties.

The effects of tempe protein isolate from non-germinated and germinated soybean on oxidative stress in diabetic rats

Rachmawati, N.A., Astawan, M., Wresdiyati, T. and Yoshari, R.M.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).19](https://doi.org/10.26656/fr.2017.7(S1).19)

The effects of tempe protein isolate from non-germinated and germinated soybean on oxidative stress in diabetic rats was studied by Rachmawati *et al.*

Galangal and ginger essential oils exerted microbial growth inhibitory activity and preservation potential on tofu

Hamad, A., Djalil, A.D. and Hartanti, D.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).7](https://doi.org/10.26656/fr.2017.7(S1).7)

The microbial growth inhibitory activity on tofu preservation by galangal and ginger essential oils was evaluated by Hamad *et al.*

Omega-3 profiles and chemical substances of chicken meat fed diets containing purslane (*Portulaca oleraceae*) meal rich in omega-3 fats

Kartikasari, L.R., Hertanto, B.S. and Nuhriawangsa, A.M.P.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).15](https://doi.org/10.26656/fr.2017.7(S1).15)

The omega-3 profiles and chemical substances of chicken meat fed diets containing purslane (*Portulaca oleraceae*) meal was evaluated by Kartikasari *et al.*

Antioxidant activity and isoflavone content of overripe Indonesian tempe

Astawan, M., Cahyani, A.P. and Wresdiyati, T.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).16](https://doi.org/10.26656/fr.2017.7(S1).16)

Astawan *et al.* evaluated the antioxidant activity and isoflavone content of overripe Indonesia tempe.

Nutritional, physical, and sensory properties of fish crackers produced from the head of catfish (*Clarias gariepinus*)

Canti, M., Kurniady, F., Hutagalung, R.A. and Prasetya, W.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).17](https://doi.org/10.26656/fr.2017.7(S1).17)

The nutritional, physical and sensory properties of fish crackers produced from the head of catfish (*Clarias gariepinus*) were studied by Canti *et al.*

Physicochemical and microbiological characteristics of fruit-based kombucha

Zubaidah, E., Cahyadi, A.B., Srianta, I. and Tewfik, I.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).8](https://doi.org/10.26656/fr.2017.7(S1).8)

Zubaidah *et al.* evaluated the physicochemical and microbiological characteristics of fruit-based kombucha.

The production process of tempe protein isolate from germinated soybeans and its potential as an antidiabetic

Yoshari, R.M., Astawan, M., Prangdimurti, E. and Wresdiyati, T.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).23](https://doi.org/10.26656/fr.2017.7(S1).23)

Yoshari *et al.* produced tempe protein isolate from germinated soybeans for its antidiabetic potential.

Amylase, protease, and lipase-producing microbes of local origin as potential starter cultures for low-salt *moromi* fermentation

Devanthi, P.V.P., Wardhana, Y.R., Pratiwi, G. and Surjawan.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).22](https://doi.org/10.26656/fr.2017.7(S1).22)

Devanthi *et al.* studied on the amylase, protease and lipase-producing microbes of local origin as a potential starter cultures for low-salt *moromi* fermentation.

The effect of rising *Monascus* fermented durian seed concentration on physicochemical and organoleptic properties of meat analog consisted of sweet potato flour and gluten

Srianta, I., Soejanta, B.R., Natanael, J., Ristiarini, S., Nugerahani, I. and Tewfik, I.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).4](https://doi.org/10.26656/fr.2017.7(S1).4)

Srianta *et al.* evaluated the effect of rising *Monascus* fermented durian seed concentration on physicochemical and organoleptic properties of meat analog consisted of sweet potato flour and gluten.

Starter cultures inoculation procedure changes microbial community structure during low-salt *moromi* fermentation

Pramanda, I.T., Saputro, M.N.B., Naidu, N.C. and Devanthi, P.V.P.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).18](https://doi.org/10.26656/fr.2017.7(S1).18)

Pramanda *et al.* evaluated the changes in the microbial community structure via the starter cultures inoculation procedure during low-salt *moromi* fermentation.

Potency of pluchea (*Pluchea indica* Less) leaves to increase functional value of wet noodles: a review

Widyawati, P.S., Darmoatmodjo, L.M.Y.D., Wibisono, D.A.S., Putra, E.W. and Dharma, A.W.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).21](https://doi.org/10.26656/fr.2017.7(S1).21)

Widyawati *et al.* reviewed the potency of pluchea (*Pluchea indica* Less) leaves to increase functional value of wet noodles.

Efficacy of soy biopeptide on serum iron, serum ferritin and hemoglobin levels of adolescent girls in Pandeglang district in Indonesia

Laily, N., Aji, G.K., Sukarti, I., Susanti, I., Illaningtyas, F. and Pranamuda, H.

Available Online: 27 APRIL 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).20](https://doi.org/10.26656/fr.2017.7(S1).20)

The efficacy of soy biopeptide on serum iron, serum ferritin and hemoglobin levels of adolescent girls in Pandeglang district, Indonesia was evaluated by Laily *et al.*

Hypocholesterolemic effects of noodles prepared from sago, sorghum and mung bean flours in hyperglycemic rats

Wahjuningsih, S.B. and Azkia, M.N.

Available Online: 20 MAY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).25](https://doi.org/10.26656/fr.2017.7(S1).25)

The hypocholesterolemic effects of noodles prepared from sago, sorghum and mung bean flours in hyperglycemic rats was studied by Wahjuningsih and Azkia.

Mold characterization in "RAPRIMA" tempeh yeast from LIPI and over-fermented *Koro Pedang* (Jack Beans) tempeh

Yarlina, V.P., Nabilah, F., Djali, M., Andoyo, R. and Lani, M.N.

Available Online: 20 MAY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).27](https://doi.org/10.26656/fr.2017.7(S1).27)

Yarlina *et al.* characterized mold in "RAPRIMA" tempeh yeast from LIPI and over-fermented *Koro Pedang* (Jack beans) tempeh.

Optimization of temperature and reaction influence on ultrasound-modified sweet potato starch

Ulfa, G.M., Putri, W.D.R., Fibrianto, K. and Widjanarko, S.B.

Available Online: 20 MAY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).12](https://doi.org/10.26656/fr.2017.7(S1).12)

Ulfa *et al.* optimized the temperature and reaction influence on ultrasound-modified sweet potato starch.

3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase inhibitory activity of Indonesian *Cajanus cajan* leaves and *Zingiber officinale* extracts

Wresdiyati, T., Papilaya, M.C., Laila, S.R., Darawati, M., Sadiyah, S. and Astawan, M.

Available Online: 10 JUNE 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).26](https://doi.org/10.26656/fr.2017.7(S1).26)

Wresdiyati *et al.* evaluated the 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase inhibitory activity of the extracts of the Indonesian *Cajanus cajan* leaves and *Zingiber officinale*.

Identification of lactic acid bacteria isolated from ethnic fermented bamboo shoot "*Lemea*" in Bengkulu, Indonesia

Okfrianti, Y., Herison, C., Fahrurrozi and Budiyanto

Available Online: 10 JUNE 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).29](https://doi.org/10.26656/fr.2017.7(S1).29)

Okfrianti *et al.* identified and isolated lactic acid bacteria from ethnic fermented bamboo shoot "*Lemea*" in Bengkulu, Indonesia.

Effect of rice bran and spent soymilk on the dough rheological properties and quality of bread

Wang, C.C.R., Nugrahedi, P.Y. and Johan, Y.

Available Online: 2 JULY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).30](https://doi.org/10.26656/fr.2017.7(S1).30)

The effect of rice bran and spent soymilk on the dough rheological properties and quality of bread was studied by Wang *et al.*

Red kidney bean (*Phaseolus vulgaris* L.) instant porridge: effect of isomalto-oligosaccharides and Fibercreme as sucrose replacement on lipid profile improvement in hypercholesterol-induced rats

Marsono, Y., Putri, R.G., Gunawan, H. and Indrawanto, R.

Available Online: 2 JULY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).24](https://doi.org/10.26656/fr.2017.7(S1).24)

Marsono *et al.* developed red kidney bean (*Phaseolus vulgaris* L.) instant porridge and studied the effect of isomalto-oligosaccharides and Fibercreme as sucrose replacement in hypercholesterol-induced rats.

Effect of using different clarifying agents and temperature on physicochemical and sensory properties of sweet sorghum syrup extract

Harlen, W.C. and Ristiarini, S.

Available Online: 2 JULY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).28](https://doi.org/10.26656/fr.2017.7(S1).28)

The effect of using different clarifying agents and temperature on physicochemical and sensory properties of sweet sorghum syrup extract was studied by Harlen and Ristiarini.

Synergistic effect of kappa-carrageenan and konjac flour in enhancing physicochemical and organoleptic properties of wheat-based edible straw

Jati, I.R.A.P., Natasha, L., Nugraha, D.T., Virly and Setijawaty, E.

Available Online: 2 JULY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).35](https://doi.org/10.26656/fr.2017.7(S1).35)

Jati *et al.* evaluated the synergistic effect of kappa carrageenan and konjac flour in enhancing physicochemical and organoleptic properties of wheat-based edible straw.

The food industry supply chain in the light of COVID-19: the constraint and development of measures to ensure food safety and quality control

Nanfack, D.C.V., Kaharso, V.C., Hlaing, K.S.S., Phaviphu, K., Sun, J., Ji, J. and Sun, X.

Available Online: 19 JULY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).34](https://doi.org/10.26656/fr.2017.7(S1).34)

Nanfack *et al.* reviewed the food industry supply chain in the light of COVID-19.

Effect of bee pollen on the characteristic of bread incorporated with *Monascus*-fermented durian seeds and rice bran

Goberto, M.A., Trisnawati, C.Y., Nugrahani, I., Srianta, I. and Marsono, Y.

Available Online: 19 JULY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).36](https://doi.org/10.26656/fr.2017.7(S1).36)

The effect of bee pollen on the characteristic of bread incorporated with *Monascus*-fermented durian seeds and rice bran was studied by Goberto *et al.*

Effect of addition of texturizer on the proximate content and texture properties of geblek, a cassava starch-based traditional food from Indonesia

Ratnaningsih, N., Sugati, D., Handayani, T.H.W., Handayani, S. and Devi, M.K.

Available Online: 19 JULY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).33](https://doi.org/10.26656/fr.2017.7(S1).33)

Ratnaningsih *et al.* studied on the effect of addition of texturizer on the proximate content and texture properties of geblek, a cassava starch-based Indonesian traditional food.

Evaluation of hypoglycaemic potency in tempe with soybean germination process and extended fermentation time

Abdurrasyid, Z., Astawan, M., Lioe, H.N. and Wresdiyati, T.

Available Online: 19 JULY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).32](https://doi.org/10.26656/fr.2017.7(S1).32)

The evaluation of hypoglycaemic potency in tempe with soybean germination process and extended fermentation time was studied by Abdurrasyid *et al.*

Quality of purula (rice seasoning for anemia from soy protein hydrolysate and seaweed) during pilot plant scale development using drum drying process

Kahfi, J., Laily, N., Pangestu, A., Muhammaludin, and Rachman, D.

Available Online: 19 JULY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).14](https://doi.org/10.26656/fr.2017.7(S1).14)

Kahfi *et al.* evaluated the quality of purula during pilot plant scale development using drum drying process.

In silico anticholesterol of Monacolin from *Monascus* sp. on HMG-CoA protein receptor

Singgih, M., Kurniati, N.F., Permana, B., Amelia, D.R. and Yuliana, A.

Available Online: 19 JULY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).1](https://doi.org/10.26656/fr.2017.7(S1).1)

Singgih *et al.* conducted an in silico study on the anticholesterol of Monacolin from *Monascus* sp. on HMG-CoA protein receptor.

Application of ultrasound in germinated soybean tempe protein concentrate production with various types of solvents

Prayudani, A.P.G., Winarsih, W., Subarna, Syamsir, E. and Astawan, M.

Available Online: 19 JULY 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).2](https://doi.org/10.26656/fr.2017.7(S1).2)

Prayudani *et al.* applied ultrasound to extract and produce protein concentrate from germinated soybean tempe using various types of solvents.

Flavonoid and main aroma-active compounds identification of Taiwan *Citrus depressa* Hayata's peels

Saputri, D.S., Yanti, S., Lin, H-Y., Agrawal, D.C. and Chien, W-J.

Available Online: 15 AUGUST 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).38](https://doi.org/10.26656/fr.2017.7(S1).38)

Saputri *et al.* identified flavonoid and main aroma-active compounds extracted from Taiwan *Citrus depressa* Hayata's peels.

Antioxidant activities and physical properties of milk chocolate enriched with plant-based functional ingredients

Muhammad, D.R.A., Zulfa, F., Purnomo, D., Widiatmoko, C. and Fibri, D.L.N.

Available Online: 15 AUGUST 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).6](https://doi.org/10.26656/fr.2017.7(S1).6)

The antioxidant activities and physical properties of milk chocolate enriched with plant-based functional ingredients were studied by Muhammad *et al.*

Isolation and identification of gastric acid-tolerant yeast from tapai

Wahyuni, I., Subagio, A., Nurhayati, N. and Purwandari, U.

Available Online: 15 AUGUST 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).13](https://doi.org/10.26656/fr.2017.7(S1).13)

Wahyuni *et al.* isolated and identified gastric acid-tolerant yeast from tapai.

The differences in the results of haemoglobin level examinations in female adolescents using digital and automated hematology analyzer methods

Illaningtyas, F., Laily, N., Sukarti, I. and Widarsih, W.

Available Online: 10 SEPTEMBER 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).10](https://doi.org/10.26656/fr.2017.7(S1).10)

Illaningtyas *et al.* evaluated the differences in the results of haemoglobin level examinations in female adolescents using digital and automated hematology analyzer methods.

The isolation and characterization of indigenous microbes with probiotic potential from gatotan

Nurhayati, N., Subagio, A., Purwandari, U. and Zamzami, F.Y.

Available Online: 10 SEPTEMBER 2023 | [https://doi.org/10.26656/fr.2017.7\(S1\).9](https://doi.org/10.26656/fr.2017.7(S1).9)

Nurhayati *et al.* isolated and characterized the indigenous microbes with probiotic potential from gatotan.

VOL. 7 | ISSUE 4

Physicochemical and microbiological characteristics of fruit-based kombucha

^{1,*}Zubaidah, E., ¹Cahyadi, A.B., ²Srianta, I. and ³Tewfik, I.

¹Department of Food Science and Technology, Faculty of Agricultural Technology, Brawijaya University, Jalan Veteran Malang, Indonesia 65145

²Department of Food Technology, Faculty of Agricultural Technology, Widya Mandala Catholic University Surabaya, Jalan Dinoyo 42-44 Surabaya, 60265, Indonesia

³School of Life Sciences, University of Westminster, 115 New Cavendish Street, London, W1W 6UW, United Kingdom

Article history:

Received: 5 February 2022

Received in revised form: 28 March 2022

Accepted: 4 October 2022

Available Online: 27 April 2023

Keywords:

Kombucha,
Fruit,
Physical,
Chemical,
Microbiology

DOI:

[https://doi.org/10.26656/fr.2017.7\(S1\).8](https://doi.org/10.26656/fr.2017.7(S1).8)

Abstract

Kombucha is a fermented drink made out of tea leaves, although recently other alternatives were considered as substitutes, such as fruits. Using different types of fruit could affect the characteristics of fruit-based kombucha. This study investigated the physicochemical and microbiological characteristics of fruit-based kombucha through a Randomized Block Design with the type of fruit (red dragon fruit, apple, snake fruit, strawberries, grapes, pear, red guava, and citrus) as the factor. The produced kombucha drinks were analyzed and compared for pH, total sugar, total phenolic compounds, total flavonoids, antioxidant activity, and total microbes. Statistical tests such as Analysis of Variance (ANOVA) and Least Significance Different (LSD) were employed ($\alpha = 5\%$). Results revealed that physicochemical and microbiological characteristics of kombucha were significantly related to the type of fruit with snake fruit kombucha showing the optimal results of each characteristics as follows: total acetic acid bacteria and yeasts of 1.53×10^9 CFU/mL, pH of 3.07, total sugar of 2.41%, total phenol of 1006.85 $\mu\text{g/mL}$ GAE, total flavonoids of 1.75 mg QE/mL, and IC_{50} of DPPH scavenging activity of 5.46 $\mu\text{g/mL}$. Fermented fruit-based kombuchas are regarded as healthy substitutes to traditional kombucha as they offer rich source of nutrients that enhance human well-being.

1. Introduction

Kombucha is one of many traditional healthy drinks from China, its produced by fermenting tea with starter kombucha called SCOBY (Symbiotic Culture of Bacteria and Yeast) (Naland, 2008). The antioxidant properties of kombucha comes from organic acid, such as lactic acid, acetic acid, ascorbic acid, and gallic acid. Kombucha also contains polyphenol compounds that act as antioxidant such as flavonoids and tannins. Furthermore, the antimicrobial properties of kombucha improve the microflora in the intestine. In general, kombucha is made from tea, but recently fruit-based kombuchas have begun to be developed. Using fruits as the raw material of kombucha would produce kombucha with various functional benefits and appeal to the market.

Indonesia is a country with an abundance of fruits which can be employed to formulate beneficial functional kombucha drinks, including dragon fruit, apple, snake fruit, strawberry, grape, pear, guava, and citrus. Dragon fruit is rich in vitamins e.g., thiamine,

riboflavin, niacin, carotene, and ascorbic acid which play a role as antioxidants (USDA, 2017). Rome beauty apple (*Malus sylvestris Mill*) is a fruit with high phenolic compounds, e.g., quercetin, glycosidic, procyanidin B2, chlorogenic acid, and epicatechin (Kurniawan, 2014). Suwaru snake fruit is rich in flavonoid, alkaloid, terpenoid, catechin, tannin, quinone, and ascorbic acid (Priyatno, 2006). Strawberry is cultivated worldwide and is known for its high content of vitamin C. Also, it is high in phytochemical compounds such as anthocyanin, flavanol, catechin, and epicatechin (Seeram, 2007). Grapefruit is high in vitamin C, K, A, and flavonoid compounds (Tepe and Hoover, 2015). Pear contains chlorogenic acid which is a hydroxyl cinnamic acid derivative that can prevent the formation of cancer cells (Gui *et al.*, 2016). Guava contains high phenolic and flavonoid compounds such as quercetin, guaijaverin, galic acid, leucocyanidin, and elagat acid (Javed *et al.*, 2017). Citrus contains D-limonene that prevents skin cancer, β -cryptoxanthin that improves respiratory system, and many other flavonoids and phytochemicals

*Corresponding author.

Email: elok@ub.ac.id

(Makni et al., 2018).

As the raw material of kombucha affects the fermentation products such as organic acid, phenol, and flavonoid, using different types of fruit as raw material can affect the characteristics of fruit-based kombucha. Therefore, this study was aimed to investigate the physicochemical and microbiological characteristics of fruit-based kombucha.

2. Materials and methods

2.1 Materials

Red dragon fruit (*Hylocereus* sp.) was resourced from a local store in Malang. Rome Beauty apple (*Malus sylvestris* Mill) and strawberry (*Fragiara x ananassa*) were collected from local farmer in Bumiaji, Batu. Snake fruit (*Sallaca zallaca*) was purchased from a local farmer in Suwaru, Pagelaran, Malang. Grape (*Vitis vinifera*), pear (*Pyrus communis*), guava (*Psidium guajava*), and citrus (*Citrus reticulate*) were bought from Blimbing Market in Malang. Kombucha (control) was made from “Naga” black tea. Sugar, commercial kombucha starter, and water were resourced from the University laboratories.

2.2 Preparation of fruit juice

The preparation of fruit juice was conducted according to Esti and Sediadi (2000) with minor modifications. The fruits were washed with clean water and drained. They were then peeled, weighed in 250 g, cut into small pieces, and blended for 30 mins (water: fruit = 1:1). After that, the fruit pulp was filtered and the fruit juice was analyzed as raw material.

2.3 Preparation of fruit-based kombucha

Fruit juice (400 mL) and 10% sugar were mixed and pasteurized (Waterbath Memmert, Germany) at 85°C for 15 mins, was let cool to 25±2°C, then poured into a sterile glass jar. After that, 10% of kombucha starter (SCOBY) was inoculated, covered with cloth, and incubated at 25±2°C for 14 days. At the end of the fermentation, the kombucha was filtered (using cheese cloth) to separate it from the SCOBY.

2.4 Determination of pH and total sugar

The pH of kombucha was measured with a pH meter (Hanna, Thermo Fischer Scientific, USA). Total sugar of kombucha analysis was conducted according to Apriyantono et al. (1989). Samples (1 mL) were transferred to test tubes and mixed with 5 mL of anthrone reagent (0.05 g of anthrone in 50 mL of concentrated H₂SO₄). The test tubes were covered with plastic, homogenized, and heated at 100°C for 12 mins.

After the samples cooled, the absorbance was measured at 630 nm in a spectrophotometer (Unico, UV-2100 Spectrophotometer) with a glucose solution as the standard.

2.5 Determination of total phenolic compounds and total flavonoid

The total phenolic compounds were determined with Follin – Ciocalteu reagent (1:10) according to Prangdimurti (2009). Samples (0.5 mL) were placed into test tubes with 2.5 mL Follin-Ciocalteu reagent, vortexed, and were put aside for 5 mins. After that, 4 mL of Na₂CO₃ 7.5% (w/v) (7.5 g Na₂CO₃ in 100 mL distilled water) was added. The mixture was vortexed and put aside for 1 hr in a dark place. Absorbance was measured at 765 nm with gallic acid as standard in Unico, UV-2100 Spectrophotometer. The total phenolic compounds were expressed as mg GAE/mL (Gallic Acid Equivalent/mL).

Total flavonoid was measured according to Atanassova et al. (2011) with slight modification. Samples (1 mL) were placed in test tubes with 4 mL of distilled water. Then, 0.3 mL NaNO₂ 5% was added and put aside for 5 mins, before 0.3 mL of AlCl₃ 10%, 2 mL of NaOH 1M, and 2.4 mL of distilled water were added. The mixtures were vortexed and put aside for 5 mins. Absorbance was measured at 510 nm with quercetin as the standard.

2.6 Determination of antioxidant activity

The antioxidant activity was measured with DPPH method referenced from Huang et al. (2005). Samples (1 mL) were dissolved with distilled water in 10 mL volumetric flask. There were four series of dilutions e.g. 2000, 1500, 1000, and 500 ppm from the main solution. Then, 1 mL of 0.2 mM DPPH was added, homogenized, and put aside for 30 mins. Absorbance was measured using a spectrophotometer (Unico, UV-2100 Spectrophotometer) at 517 nm.

2.7 Determination of total microbes

The determination of total microbes was done using the standard plate count procedure according to Fardiaz (1992). Samples (1 mL) were mixed with 9 mL of sterile 0.1% peptone water and further diluted to have a serial of decimal dilutions. Each dilution (1 mL) was transferred into petri dishes and poured with sterile plate count agar (50°C), mixed, and left to solidify before incubation at 37°C for 24 hrs. Colony counts were calculated with the following equation expressed in log CFU/mL: [total colony per mL = total colony×(1/diluted factor)].

Table 1. pH of fruit-based kombucha during the fermentation process

Types of Fruit	pH			Decrease (%)	LSD 5%
	Day-0	Day-7	Day-14		
Apple	3.43	3.13	2.83	0.17	0.36
Black Tea	4.10	3.33	3.10	0.24	
Citrus	4.27	3.63	3.23	0.24	
Dragon Fruit	4.17	3.17	2.80	0.33	
Grape	4.07	2.80	2.77	0.32	
Guava	4.07	3.30	2.90	0.29	
Pear	4.50	3.20	3.07	0.32	
Snake Fruit	4.10	3.30	3.07	0.25	
Strawberry	3.33	3.00	2.93	0.12	

2.8 Statistical analysis

The data was analyzed using Minitab 17. ANOVA (Analysis of Variance) and LSD (Least Significant Difference) with $\alpha = 5\%$ were used to differentiate between the mean values.

3. Results and discussion

3.1 pH of fruit-based kombucha

The pH of fruit-based kombucha ranged from 3.33 – 4.50 on day-0, 2.80 – 3.63 on day-7, and 2.77 – 3.23 on day-14. According to Table 1, the pH values gradually decreased from day-0 to day-14. Based on statistical analysis, there was no significant effect ($\alpha = 0.05$) of different types of fruit on the kombucha pH average. The pH of fruit-based kombucha decreased during the fermentation process. The decrease of pH during fermentation may be caused by the accumulation of organic acids as a result of yeast and bacterial metabolism. During fermentation, the yeasts involved produce ethanol and carbon dioxide. Ethanol is oxidized by acetic acid bacteria to become acetaldehyde, then oxidized to acetic acid. Acetic acid bacteria also produce gallic acid during kombucha fermentation. In addition, there are also other organic acids such as malic acid, oxalic acid, and lactic acid (Júnior *et al.*, 2009). On the last day of fermentation, the lowest pH of kombucha was recorded.

Table 2. Total sugar of fruit-based kombucha during the fermentation process

Types of Fruit	Total Sugar (%)			Decrease (%)	LSD 5%
	Day-0	Day-7	Day-14		
Apple	13.72	7.03	2.38	0.83	0.55
Black Tea	10.55	5.28	1.81	0.83	
Citrus	10.59	5.35	1.83	0.83	
Dragon Fruit	14.74	7.57	2.59	0.82	
Grape	13.82	6.95	2.23	0.84	
Guava	12.47	6.18	2.09	0.83	
Pear	13.66	6.81	2.31	0.83	
Snake Fruit	14.17	7.41	2.41	0.83	
Strawberry	11.42	5.74	1.99	0.83	

3.2 Total sugar of fruit-based kombucha

Based on analysis, total sugar of kombucha with different types of fruit ranged from 8.84% to 12.96% on day-0, 4.05% to 6.47% on day-7, and 2.43% to 3.86% on day-14. The total sugar of fruit-based kombucha was shown on Table 2. Based on LSD analysis, kombucha made from different types of fruit had no significant difference on total sugar. The decrease of total sugar in fruit-based kombucha may be caused by the synergistic metabolic activity of yeast and bacteria. In such metabolism, sugar was used as a substrate in the metabolic reaction. According to Napitupulu *et al.* (2015), the sugar contained in kombucha is generally sucrose sugar. Sucrose is converted into glucose and fructose with invertase enzyme. Glucose will then be used in the Krebs's Cycle.

3.3 Total phenol of fruit-based kombucha

The average of total phenol in fruit-based kombucha ranged from 82.61 $\mu\text{g/mL}$ GAE to 895.03 $\mu\text{g/mL}$ GAE on day-0, 98.97 $\mu\text{g/mL}$ GAE to 954.73 $\mu\text{g/mL}$ GAE on day-7, and 198.67 $\mu\text{g/mL}$ GAE to 1006.85 $\mu\text{g/mL}$ GAE on day-14. Figure 1 demonstrates an increase of total phenol in fruit-based kombucha from day-0 to day-14. Based on LSD analysis, different types of fruit resulted in significant differences on the total phenol of fruit-based kombucha. Figure 1 shows the increase of total phenol during the fermentation period. This occurred due to the activity of microorganisms that carry out fermentation. During fermentation, microorganisms produced enzymes that can break down phenolic compounds, causing a change in complex phenolic compounds into simpler phenolic compounds, thereby increasing the average total phenolic content. According to Suhardini and Zubaidah (2016), the increase in phenolic compounds during fermentation is caused by the involvement of various types of microbes, bacteria and yeasts that can be metabolized to produce phenolic compounds through enzymatic reactions. *Saccharomyces cerevisiae*, a common yeast found in kombucha, is able to produce β -glucosidase enzymes that can break down glycoside bonds so that phenolic compounds can be released into fermentation medium. *S. cerevisiae* is also able to synthesize free phenolic compounds, namely 4-ethylphenol and 4-ethylguaiacol derived from hydroxybenzoic acid and ferulic acid. *S. cerevisiae* is able to produce cinnamic decarboxylase and vinylphenol reductase enzymes. Cinnamic decarboxylase enzyme catalyzes the decarboxylation process of cinnamic acid and vinylphenol reductase enzyme catalyzes the reduction of 4-vinyl phenol and 4-vinylguaiacol into 4-ethyl phenol and 4-ethylguaiacol. However, the increase in phenolic compounds is not always significant.

Nonetheless, phenols which also act as an antimicrobial compounds can also cause a decrease in the activity of microorganisms in kombucha.




Figure 1. Total phenol of fruit-based kombucha during the fermentation process

Based on Figure 1, the kombucha with the highest total phenol on day-14 was snake fruit kombucha, while the one with the lowest total phenol was kombucha pear. The difference in total phenol between kombuchas was based on their phenolic compounds and the types of phenols they contained in each raw material (Essawet *et al.*, 2015). According to Jayabalan *et al.* (2007), the content of phenolic compounds is influenced by the flavonoid content, where flavonoid compounds are influenced by cultivation conditions and seasonality (the availability of sunlight, nutrients, pH and humidity).

3.5 Total flavonoid of fruit-based kombucha

The fermentation process from day-0 to day-14 indicated an increase of total flavonoid content in fruit-based kombucha. Based on Figure 2, the total flavonoids in fruit-based kombucha were 1.12 – 4.01 µg/mL GAE on day-0, 1.50 – 5.26 µg/mL GAE on day-7, and 1.75 – 6.05 µg/mL GAE on day-14. According to LSD statistical analysis, different types of fruits in kombucha had no significant difference on total flavonoid. Microorganisms produced enzymes that break down complex flavonoid into simple flavonoid compounds. According to Chu and Chen (2006), the total flavonoid compounds increase linearly with the length of fermentation time. Four types of catechins commonly found in kombucha such as epicatechin and epigallocatechin were converted into simpler ones by enzymes produced from kombucha fermentation. Yeast type *C. tropicalis* can degrade several types of polyphenols and flavonoids (Ettayebi *et al.*, 2003). While yeast *S. cerevisiae* has the ability to convert several types of organic acids into flavonoid compounds, such as cinnamic acid to pinocembrin, p-coumaric to naringenin, and caffeic acid to eriodictyol (Yan *et al.*, 2005).

Based on Figure 2, the highest mean of total flavonoid on day-14 was from the strawberry kombucha. Meanwhile, the one with the lowest average was snake

fruit kombucha. The average total flavonoid was highest in strawberry kombucha while the lowest increase was in citrus kombucha. In Figure 2, strawberry had the highest total flavonoid while snake fruit had the lowest total flavonoid. The content of flavonoids in fruit was influenced by various factors such as growing conditions, sunlight for photosynthesis, nutrients, moisture and soil pH (Jayabalan *et al.*, 2007). According to Koopman *et al.* (2012), another factor that can affect the total flavonoids of kombucha is the sugar content in the raw material as the main carbon source in the formation of flavonoid compounds.




Figure 2. Total flavonoid of fruit-based kombucha during the fermentation process

3.6 Antioxidant IC_{50} of fruit-based kombucha

In this study, antioxidant activity was expressed using IC_{50} value. IC_{50} is the sample concentration which reduces the DPPH activity by 50%. The smaller the IC_{50} value, the higher the antioxidant activity. IC_{50} value of fruit-based kombucha ranged from 8.35 to 47.50 mg/mL on day-0, 6.11 to 34.76 mg/mL on day-7, and 5.46 to 31.04 mg/mL on day-14. The chart of IC_{50} value is shown in Figure 3. Figure 3 indicates the decrease of IC_{50} value of fruit-based kombucha. The antioxidant activity of kombucha was influenced by organic acids, total phenols, and total flavonoids it contained. In addition, the sugar content in kombucha as a source of nutrients for metabolism also affected antioxidant activity where the increased sugar would increase the in production of organic acids and enzymes that could degrade complex phenolic compounds into simple




Figure 3. Antioxidant activity of fruit-based kombucha during the fermentation process

phenolic compounds.

Apple kombucha had the highest decrease in IC_{50} value while snake fruit kombucha had the lowest decrease. The decrease of IC_{50} value was caused by an increase in phenolic compounds as antioxidants due to a biotransformation process that utilized certain enzymes in a plant cell to change the functional group of a chemical compound found in plants (Jayabalan *et al.*, 2008). Furthermore, the decrease of IC_{50} on day-7 was not significant. Based on Ayu *et al.* (2013), acidic conditions caused phenolic compounds to become more stable and difficult to release protons that can bind to DPPH.

3.7 Total microbes of fruit-based kombucha

The total microbes of fruit-based kombucha showed varied results. The microbes grown were most likely acetic acid bacteria and yeasts which are commonly found in kombucha. The average total microbes ranged from 5.6×10^8 CFU/mL to 9.3×10^{10} CFU/mL on day-0, 2.0×10^9 CFU/mL to 1.4×10^{11} CFU/mL on day-7, and 3.1×10^8 CFU/mL to 2.2×10^{10} CFU/mL. There was a gradual increase of total microbes from day-0 to day-7 and a decrease of total microbes from day-7 to day-14 as shown on Figure 3. The difference in the total microbes between types of kombucha was influenced by the characteristics of the fruit. The average of total microbes is shown on Table 3. During the first week of fermentation, microbe cells would multiply first and cause an increase of total microbes. The sucrose in kombucha was then converted to glucose and fructose. The glucose broke down to organic acids and energy. The organic acid induced the optimum environment for acetic acid bacteria growth with pH around 5.0 – 6.5 up to 3 – 4. Data presented in Table 3 showed that apple kombucha has the highest level of total microbes on day-14 and citrus kombucha has the lowest value. Lončar *et al.* (2014) reported that sugar is a convenient substrate for bacteria metabolism. Organic acids and energy were the results of metabolism and were used for cell multiplication so that the higher the sugar concentration added, the higher the total microbes obtained. According to Caldwell (2000), excessive amounts of nutrients can

actually have an impact on the accumulation of inhibitory compounds. The difference in total microbes between fruit-based kombucha was caused by various influencing factors such as the total sugar acts as a carbon source and energy provider, natural antioxidants in fruits, and the total content of phenolic compounds and flavonoids in kombucha.

4. Conclusion

Based on the findings, different types of fruit had no significant difference on pH, total sugar, total flavonoid in fruit-based kombucha. In contrast, different types of fruit significantly affected the total phenol and antioxidant activity. Snake fruit kombucha showed the optimal characteristics as a probiotic drink and a substitute of tea kombucha as follows: total microbes of 1.53×10^9 CFU/mL, the pH of 3.07, total sugar of 2.41%, total phenol of 1006.85 μ g/mL GAE, total flavonoids of 1.75 mg QE/mL, and IC_{50} of DPPH scavenging activity of 5.46 μ g/mL.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

This research was financially supported by Universitas Brawijaya.

References

- Aditiwati, P. and Kusnadi. (2003). Mixed Culture and Environmental Factors of Microorganisms Playing a Role in "Tea-Cider" Fermentation. *Proceeding ITB Sains dan Teknologi*, 35(2), 147-162. <https://doi.org/10.5614/itbj.sci.2003.35.2.5>
- Apriyantono, A.D., Fardiaz, D., Puspitasari, N.L., Sudarmadji, Sedamawati and Budiyanto, S. (1989). *Food Analysis*. Bogor, Indonesia: Pusat Antar Universitas Pangan dan Gizi IPB.
- Atanassova, M., Georgieva, S. and Ivancheva, K. (2011). Total Phenolic and Total Flavonoid Contents, Antioxidant Capacity and Biological Contaminants

Table 3. Total microbes of fruit-based kombucha during the fermentation process

Types of Fruit	Total Microbes (CFU/mL)			
	Day-0	Day-7	Day-14	Difference
Apple	9.29×10^{10}	1.20×10^{11}	2.17×10^{10}	-7.12×10^{10}
Black Tea	4.74×10^9	1.41×10^{11}	2.23×10^9	-2.51×10^9
Citrus	9.26×10^8	7.83×10^9	3.13×10^8	-6.13×10^8
Dragon Fruit	9.16×10^9	5.97×10^{10}	7.3×10^8	-8.43×10^9
Grape	1.26×10^9	3.59×10^{10}	4.46×10^9	3.20×10^9
Guava	5.56×10^8	2.01×10^9	4.86×10^8	-7.00×10^7
Pear	1.22×10^{10}	6.49×10^9	2.5×10^9	-9.70×10^9
Snake Fruit	7.3×10^8	6.53×10^{10}	1.53×10^9	8.00×10^8
Strawberry	3.5×10^8	7.15×10^{10}	2.43×10^9	2.08×10^9

- in Medicinal Herbs. *Journal of The University of Chemical Technology and Metallurgy*, 46(1), 81-88.
- Ayu, S., Yan, R. and Eka, L. (2013). Determination of Antioxidants in Local Kombucha Black Tea in Bali with Fermentation Time. Bali, Indonesia: Udayana.
- Caldwell, D.R. (2000). *Microbial Physiology and Metabolism*. 2nd ed. USA: Star Publishing Company.
- Chu, C.S. and Chen. (2006). Effect of Origin and Fermentation Time on The Antioxidant Activities of Kombucha. *Food Chemistry Journal*, 98, 502-507. <https://doi.org/10.1016/j.foodchem.2005.05.080>
- Ettayebi, K., Errachidi, F., Jamai, L., Tahri-Jouti, M.A., Sendide, K. and Ettayebi, M. (2003). Biodegradation of Polyphenols with Immobilized *Candida tropicalis* Under Metabolic Induction. *FEMS Microbiology Letters*, 233(2), 215-219. [https://doi.org/10.1016/S0378-1097\(03\)00380-X](https://doi.org/10.1016/S0378-1097(03)00380-X)
- Esti and Sediadi, A. (2000). *Food Processing*. Jakarta, Indonesia: BPPT.
- Essawet, N.A., Cvetkovic, D., Velicnaski, A.J., Brunet, C., Vulic, J., Maksimovic, V. and Markov, S. (2015). Polyphenols and Antioxidant Activities of Kombucha Beverage Enriched with Coffeeberry Extract. *Chemical Industry & Chemical Engineering Quarterly*, 21(3), 399-409. <https://doi.org/10.2298/CICEQ140528042E>
- Fardiaz. (1992). *Food Microbiological Analysis*. Jakarta, Indonesia: Grafindo.
- Gui, H.J., Sun, H.Y. and Jong, B.E. (2016). Physicochemical Characteristics and Antioxidant Activities of New Asian Pear Cultivars. *Journal of Applied Biological Chemistry*, 59(4), 337-343. <https://doi.org/10.3839/jabc.2016.057>
- Huang, Y.C., Yung H.C. and Yi-Yuan S. (2005). Effects of Genotype and Treatment on the Antioxidant Activity of Sweet Potato in Taiwan. *Food Chemistry*, 98, 529-538. <https://doi.org/10.1016/j.foodchem.2005.05.083>
- Javed, M.S., Randhawa, M.A., Ahmad, Z. and Tariq, M.R. (2017). Effect of CaCl₂ and Controlled Atmosphere Storage on Phytochemical Attributes of Guava. *Food Science and Technology*, 38(2), 356-362. <https://doi.org/10.1590/1678-457x.05517>
- Jayabalan, R., Marimuthu, S. and Swaminathan, K. (2007). Changes in Content of Organic Acids and Tea Polyphenol During Kombucha Tea Fermentation. *Food Chemistry Journal*, 102(1), 392-398. <https://doi.org/10.1016/j.foodchem.2006.05.032>
- Jayabalan, R., Subathradevi P., Marimuthu S., Sathiskumar, M. and Swaminathan, K. (2008). Changes in Free-Radical Scavenging Ability of Kombucha Tea during Fermentation. *Food Chemistry*, 109(1), 227-234. <https://doi.org/10.1016/j.foodchem.2007.12.037>
- Júnior, R.J.S., Batista, R.A., Rodrigues, S.A., Filho, L.X. and Lima, Á.S. (2009). Antimicrobial Activity of Broth Fermented with Kombucha Colonies. *Journal of Microbial and Biochemical Technology*, 1(1), 72-78. <https://doi.org/10.4172/1948-5948.1000014>
- Koopman, F., Beekwilder, J., Crimi, B., van Houwelingen, A., Hall, R.D., Bosch, D., van Maris, A.J.A., Pronk, J.T. and Daran, J.M. (2012). De Novo Production of the Flavonoid Naringenin in Engineered *Saccharomyces cerevisiae*. *Microbial Cell Factories*, 11(1), 1-15. <https://doi.org/10.1186/1475-2859-11-155>
- Kurniawan, R.F. (2014). *Efficacy and Benefits of Apple Skin*. Bandung, Indonesia: Lembar Langit Indonesia.
- Lončar, E.S., Kanuric, K.G., Malbasa, R.V., Duric, M.S. and Milanovic, S.D. (2014). Kinetics of Saccharose Fermentation by Kombucha. *Chemical Industry and Chemical Engineering Quarterly*, 20(3), 345-352. <https://doi.org/10.2298/CICEQ121113016L>
- Makni, M., Jemai, R., Kriaa, W., Chtourou, Y. and Fetoui, H. (2018). Citrus Limon from Tunisia: Phytochemical and Physicochemical Properties and Biological Activities. *BioMed Research International*, 2018, 6251546. <https://doi.org/10.1155/2018/6251546>
- Malbasa, R.V., Loncar, E.S., Djuric, M.S., Kolarov, L.A. and Klasnja, M.T. (2005). Batch Fermentation of Black Tea by Kombucha: A Contribution to Scale-Up. *APTEFF*, 36, 221-229. <https://doi.org/10.2298/APT0536221M>
- Napitupulu, M.O.W., Setyohadi and Lubis, L.M. (2015). The Effect of Sugar Concentration and the Fermentation Time of Kombucha Coffee. *Jurnal Rekayasa Pangan dan Pertanian*, 3(3), 316-322.
- Naland, H. (2008). *Kombucha Tea with a Thousand Benefits*. Jakarta, Indonesia: Agromedia.
- Prangdimurti, E. (2001). *Probiotics and their Protective Effects Against Colon Cancer*. Bogor, Indonesia: Insitut Pertanian Bogor, MSc. Thesis.
- Priyatno, L.H.A, Ibrahim, S., Adnyana, I.K. and Sukandar, E.Y. (2006). Antioxidant Activity of Fruit Flesh Extract of Bangkok Varieties of Salak (*Salacca edulis* Reinw). *Acta Pharmaceutica Indonesia*, 1, 24-27.
- Seeram, N.P. (2007). Strawberry Phytochemicals and Human Health: A Review. *Critical Reviews in Food Science and Nutrition*, 44(1), 1-17.
- Suhardini, P.N. and Zubaidah, E. (2016). Study of Antioxidant Activity on Various Kombucha Leaves During Fermentation. *Jurnal Pangan dan*

- Agroindustri*, 4(1), 221-229.
- Tepe, E.S. and Hoover, E.E. (2015). Growing grapes for home use. Retrieved on January 18, 2018 from website <http://www.extension.umn.edu/garden/yard-garden/fruit/growing-grapes-for-home-use/>
- USDA Branded Food Products. (2017). Full Report (All Nutrients): 45136474, Dragon Fruit, UPC: 609207618690. Retrieved on January 17, 2018 from <https://ndb.nal.usda.gov/ndb/foods/show/95261>
- Yan, Y., Kohli, A. and Koffas, M.A.G. (2005). Biosynthesis of Natural Flavanones in *Saccharomyces cerevisiae*. *Applied and Environmental Microbiology*, 71(9), 5610-5613. <https://doi.org/10.1128/AEM.71.9.5610-5613.2005>