Potential of coconut milk and mung bean extract combination as foam stabilizer in non-dairy ice cream

by Anna Ingani Widjajaseputra

FILE TIME SUBMITTED SUBMISSION ID 2-POTENTIAL_OF_COCONUT_MILK....PDF (241.48K) 28-AUG-2020 12:12PM (UTC+0700) WORE 1375317759 CHAR

WORD COUNT3303CHARACTER COUNT17615

International Food Research Journal 24(3): 1199-1203 (June 2017)

Journal homepage: http://www.ifrj.upm.edu.my



C All Rights Reserved

Potential of coconut milk and mung bean extract combination as foam stabilizer in non-dairy ice cream

*Widjajaseputra, A.I. and Widyastuti, T.E.W.

Faculty of Agricultural Technology, Widya Mandala Catholic University Surabaya, Surabaya 60265, East Java, Indonesia

Non-dairy ice cream is one of many food products that are currently developed especially for

vegetarians. The use of coconut milk in the manufacture of non-dairy ice cream made from

extracts of mung bean (Vigna radiata) is expected to increase foam stabilization of ice cream

mixture. Coconut milk can serve as fat and protein sources that are needed in the formation of emulsion and stabilization of the foam system that will determine the characteristic of non-dairy ice cream. The aims of this research was to study the potential of a combination of coconut milk

and mung bean extract as a foam stabilizer on a non-dairy ice cream mixture. A single factor randomized block design was used and the ratios of mung bean extract : coconut milk (w/w)

consisted of four levels, namely P1, P2, P3 and P4, respectively for 90: 10, 85: 15, 80: 20 and

75: 25. The experiments were carried out with three replications. The viscosity of the mixture

was affected by the mixture components such as starch and albumin that are capable of trapping

water in gel formation during heating process. Globulin and albumin enveloped the air bubbles and stabilized the emulsion by increasing the foam stability. Combination of mung bean extract and coconut milk in the ratio of 85 : 15 gave the maximum effect on foam stability. This is related to the proportion of the total solids and viscosity that were relatively high compared to



Abstract 19

Received: 2 June 2016 Received in revised form: 28 June 2016 Accepted: 30 June 2016

Keywords

Vigna radiate Coconut milk Foam stability Non-dairy ice cream

Introduction

*Corresponding author. Email: ingani9456@yahoo.com

Nowadays people are concerned about the health aspects of food choices, for example, some consumers prefer low-fat ice cream products and vegetable-oriented material. Sometimes, mung bean is also used as raw material, of non-dairy ice cream, but scientific studies on the use of mung bean is relatively limited. Some of the issues with non-dairy ice cream are high solids non-fat, low fat content and adequate protein component (Ahsan *et al.*, 2015). These conditions affect the texture and sensory three main structural components of air bubbles, ice crystals and fat globules.

the other combinations.

Characterization of the 16 varieties of mung bean (*Vigna radiata*) produced a wide range of proximate composition protein, 24.26 to 28.50%; crude fiber, 3.21 to 4.18%; fat, 0.57 to 1.86%; ash, 3.64 to 4.24%; water, 7.49 to 8.45%; carbohydrates, 54.25 to 58.69% (Li *et al.*, 2010). Mung bean flour is able to stabilize emulsion (37.95%), similar to the rice flour (37.31%), wheat flour (38.38%) and potato starch (41.92%) (Chandra and Samsher, 2013). The capacity of the foam formation from mung bean flour is highest (24.23%) than other flour in the order of

wheat and potatoes. At high enough concentrations (18 g/100 mL), mung bean flour can also form gels (Kaushal et al., 2012). Mung bean starch (MBS) has the ability to form gel due to a high amylose content (33%) (Keum-Il Jang et al., 2009). Mung bean starch can act as stabilizer in the manufacture of ice cream, but the functional properties (2) some proteins are required for quality ice cream. Globulin and albumin are the main storage proteins in mung bean seeds and make up over 60% and 25% of the total mung bean protein respectively (Tang et al., 2014). Studies evaluating the functional properties of mung bean protein isolate (MPI) revealed the water absorption capacity to be 1.03 to 2.78 g/g, foaming capacity as 33.00 to 67.50%, foam stability as 20.00 to 56.00%, and emulsifier activity as 1.77 to 3.30 m2/g (Li et al., 2010).

3.52%, 12.92% and 6.84%, respectively for rice flour,

Protein content in coconut milk was reported to be 5-10% 15 y basis, and based on the solubility properties, 80% of the proteins from coconut endosperm was 13 assified as albumin and globulin (Soler, 2005). Coconut milk has a high surface protein load (7 mg/m²), of which the major component is cocosin (Tangsuph 12 n and Coupland, 2009). Some proteins that exist in the liquid phase of coconut milk interact with fat globules and act as emulsifiers around its gurface (Peamprasart and Chiewchan, 2006). The emulsion formed from the aqueous extract of coconut solid endosperm is relatively unstable because of the large droplet size (Tangsuphoom and Coupland, 2008a). Coconut milk emulsions are negatively charged and not flocculate at pH 6 (Tangsuphoom and Coupland, 2008b). Coconut proteins coagulate at 80.9°C (Tipvarakamkoon, 2009), but fat droplet aggregation would occur if proteins were denatured by heat. Small oil droplets will interact and coalesence into larger droplets and caused an emulsion destabilization (Raghavendra and Raghavarao, 2010).

During the process of making ice cream, fat globules envelop air bubbles to stabilize the foam. The fat globules will also affect the size of ice crystals. Dispersed small fat globules will inhibit the development of ice crystals into large crystals. Proteins are needed to form a surface layer that surrounds the fat globules during the homogenization operation and during stirring as air bubbles are incorporated. The combination of coconut milk and mung bean extract in a proper proportice 22 vill produce expected non-dairy ice cream texture. Therefore, the purpose of this study was to evaluate the potential of combining coconut milk and mung bean extract in a non-dairy ice cream mixture.

Materials and Methods

Materials

Dehulled local mung bean, sugar, coconut, and drinking water were obtained from one supermarket in Surabaya, Indonesia. Chemicals for analysis consisted of distilled water, K₂S₂O₄, HgO, H₂SO₄, Zn plate, K₂S, NaOH, HCl, methyl red indicator, n-hexane, and were of analytical grade.

Mung bean extraction

The mungbean was washed and then soaked with water (1: 4) for 8 hr, and drained before milling in hot water (100°C) at a ratio mungbean : water of 1 : 4. Milling was done with blender (Miyako, Japan), and the milled bean suspension was filtered with a cheese cloth.

4

Manufacture of coconut milk

Coconut milk was prepared according to refers to milk-making pasta of The Indonesian Academy of Science (2000). This involved extracting the milk from coconut without adding water in a food processor.

Fable	 Formul 	ation of	of the	mung	bean	non-d	lairy	ice
		crea	um m	ixtures				

Component				
-	P1	P2	P3	P4
Mung bean extract g)	450	425	400	375
Coconut milk (g)	50	75	100	125
Sucrose (g)	60	60	60	60
Total (g)	560	560	560	560

Preparation of non-dairy ice cream mixture

Mung bean-coconut mixtures, shown in Table 1, were pasteurized at 85°C for 25 sec. The mixtures were homogenized (Ultra-Turrax T25, Janke and Kenkel IKA Labortechnik) after the temperature reached 72°C. The rotation speed of the homogenizer was gradually increased from 8,000 rpm for 2 min, followed by 10,000 rpm for 2 min. The formulation (Table 1) followed a single factor study with a randomized block design at four levels (P1-P4) of the mung bean extract : coconut milk; 90:10; 85:15; 80:20, and 75:25% (w/w). Three replicates of each formulation were prepared and studied.

Experimental design

The single factor study with a randomized block design was applied. The factor was the proportion of mung bean extract : coconut milk (P) consisting of four level of factor, namely P1, P2, P3 and P4, respectively 90: 10; 85: 15; 80: 20 and 75: 25% (w/w). The effective was carried out with three replications. Analysis of Variance (ANOVA) was performed and differences in mean values were determined using Least Significant Difference (LSD) test at $\alpha = 0.05$ by employing SPSS version 12.

Chemical analysis

Proximate composition of the mung bean extract, and chemical characteristics of the coconut milk and the non-dairy ice cream mixture were determined according to standard procedures (AOAC ,2005).

Physicochemical analysis

Viscosity, foaming capacity and foam stability were the tested physicochemical properties. Viscosity of the homogenized non-dairy ice cream mixtures was determined by viscometer (Brookfield Viscosimeter model LVDVE, USA) at $30 \pm 1^{\circ}$ C. Foaming capacity was determined by recording the volumes of the partonized mixtures before and after homogenization. The percentage ratio of volume increase to that of the original volume of pasteurized mixture was catellated and expressed as the foaming capacity. The change in volume of foam every 30 minutes until 150 minutes of standing at room temperature was recorded as the foam stability. Table 2. Composition of mung bean extract and coconut milk

	Content ± SD (%)			
Component	Mung bean extract	coconut milk*)		
Water	91.78 ± 0.746	49.58 ± 1.213		
Protein	2.41 ± 0.067	4.33 ± 0.548		
Fat	0.01 ± 0.004	24.00 ± 0.401		
Starch	7.81 ± 0.345	-		
Reducing Sugar	0.10 ± 0.008	3.33 ± 0.004		
Ash	0.04 ± 0.014	0.55 ± 0.026		

*) extraction of coconut milk by food processor without added water

Result and Discussion

Composition of mungbean extract and coconut milk

Table 2 shows the composition of the mung bean extract and coconut milk. Total solid of mung bean extract is 6.07%, while the fat content is 0.01% (wb) or 0.16% (db). Protein from coconut milk is needed in emulsions stabilization and is expected to stabilize the foam system, which will improve the characteristics of non-dairy ice cream. In addition, levels of the protein in coconut milk is higher (4.33%) compared to extracts of mung bean (2.41%).

Physicochemical characteristic of non-dairy ice cream mixture

The water content of the non-dairy ice cream mixture 23 h various proportions of mung bean extracts: coconut milk, as shown in Table 3.The water content of coconut milk (49.58%) was much lower than mung bean extract juice (91.78%). It caused the water content of ice cream mixture to drop significantly with increasing proportion of coconut milk.

Protein can form a gel. Gel formation by 20 tein conformational changes or preceded by partial denaturation of the protein molecules by heat, followed by aggregation of denatured protein molecules gradually to increase the viscosity and form a continuous network. During the preparation of mung bean extracts, heating and pasteurization process carried out at 85-90°C caused protein denaturation. Those temperature was needed to form a stable gel network due to the formation of covalent bond that converts into a thiol group of disulfide bonds and hydrophobic interactions. Intermolecular (13)Ifide bonds increase the stability of the gel matrix (Phillips *et al.*, 1994).

Functional properties of proteins in a food product can be igluenced by the size, shape, composition and amino acid sequence, charge, hydrophobicity, structure, molecular rigidity in response to environmental conditions (pH,

temperature, salt concentrati (3) or interaction with other food components. Some functional properties of proteins in food products depend on the exposure of hydrophobias roups on the molecular surface and its interaction with oil (emulan), air (foam) or another protein molecule. The residues of hydrophobic amino acids are generally located inside globular protein molecules. If the natural structure of proteins is open due to the stages of food processing, such as homogenization, liquidification and heating, it can be necessary to allow these hydrophobic groups to participate in intermolecular interactions. Flexibility molecular of proteins can be formed by the strength of hydrophobic and electrostatic interactions that maintain internal natural structure, or by the presence of covalent disulfide intramolecular bonds (Alleoni, 2006).

The results of LSD test ($\alpha = 5\%$) shown in Table 3, indicated that all treatments resulted in significantly different of viscosities. The viscosity of the mixture formed by the result of the mixture components that are capable of trapping water, such as starch and albumin, thereby reducing the mobility of the water. Prapasuwannakul *et al.* (2014) reported that the viscosity of germinated brown rice ice cream is high because of high starch content. Increasing viscosity with the higher proportion of mung bean extract showed that the viscosity of mixture formula is determined by the high content of the starch in mung bean extract. Too high viscosity of ice cream mixture was undesirable because it allowed low air incorporation in the mixture and lead to low over run.

The data in Table 3 showed the mixture of total solid is higher with increasing proportion of coconut milk, as the total solid of mung bean extract was only 6.07% compared to total solid of coconut milk (50.42%). The addition of coconut milk on mung bean extract improved protein and fat content of the ice cream mixture. Coconut milk protein that has globulin and albumin determined the characteristics of ice cream. Protein enveloped particles and stabilized fat emulsions in homogenized mixture.

Foaming capacity

Foam systems in ice cream mixture play an important role in determining the texture. Product capability in maintaining the texture and structure during and after the freezing process depends on the ability of the dough to trap air. Protein will protect and maintain the air bubbles so that it will improve the texture according to the desired character. Various proteins can be used as an emulsifying agent because its surface properties as a major factor in the expansion of ice cream mixture. Development

Treatment	Water	Viscosity	Total	Foaming
(mung bean extract :	Content	(cP)	Solid	Capacity
coconut milk)	(%)		(%)	(mL foam/mL
				total dough)
P1 (90:10)	79 ^d	10.212 ^d	20 ^a	0.85 ^b
P2 (85:15)	77°	9.670 ^c	22 ^{ab}	0.83 ^b
P3 (80:20)	76 ^b	5.670 ^b	23 ^b	0.76 ^a
P4 (75:25)	73 ^a	5.248 ^a	25°	0.74 ^a

Table 3. Water content, viscosity, total solid and foaming capacity of non-dairy ice cream mixture*)

*) For each column, values with different letters are significantly different ($\alpha = 5\%$)

of film formation of proteins associated with the protein's ability to lower the surface tension between air bubbles and solution.

The data in Table 3 showed that foam capacity decreased with increasing proportion of coconut in the mixture. This indicated that the pain proteins storage, globulin and albumin were 60% and 25% of the total mung bean protein respectively, were the important materials that have the capacity as the foaming formation. The role of mung bean protein as foam forming also shown in research of Butt and Batool (2010) who found the highest foaming capacity (110 \pm 6.78%) of mung bean protein isolates compared to cowpea and pigeon pea. High capacity of foam formation can occur because the protein was dissolved and has sufficient electrostatic repulsion that caused decreasing of protein-protein interactions. A similar case was reported by Chaiseri and Chonhenchob (2005) who studied the effect of chemical reagents on functional properties of mung 7 an protein product. The low foaming capacity could be due to inadequate electrostatic repulsions, and hence, excessive protein-protein interaction to form aggregates that were detrimental to foam formation.

Foam stability

Foam stability of the non-dairy ice cream mixture as shown in Figure 1 indicated similar pattern of foam stability with a variety of treatments. Decreasing of foam volume during 150 minutes of observation was affected by protein's ability to lower the surface tension between the surface layer of air and water system. Coconut milk proportion affected foam stability because of its fat content.

Mung bean extract used in this research is potentially a good candidate as a stabilizer to foam forming and gel system. Those potential are due to the content of albumin (25% of total protein) in the mung bean, considering that extraction was carried out using water as the solvent at neutral pH.



Figure 1. Foam stability of non-dairy ice cream mixture. P1 (mung bean extract:coconut milk=90:10); P2 (mung bean extract:coconut milk=85:15); P3 (mung bean extract:coconut milk=80:20): P4 (mung bean extract:coconut milk=75:25

Conclusion

Mixing mung bean and coconut milk displayed emulsion characteristics that are suitable for nondairy ice cream. The present study showed that the the of coconut milk with the proportion of 15:85 to mung bean extract has potential foam stabilizer in non-dairy ice cream. The starch content of the mung bean extract was about 8%, and this was thought to be mainly responsible for the increase in the viscosity of the mixtures, although both the albumin of mung bean and coconut milk also contributed to the viscosity by gelation during the heating.

Acknowledgement

Authors would like to thank to the Faculty of Agricultural Technology, Widya Mandala Catholic University Surabaya, Indonesia for funding of the research. The technical support of the staffs of the Food Chemistry-Biochemistry and Nutrition Laboratory is acknowledged research.

References

Ahsan, S., Zahoora, T., Hussain, M., Khalida, N. Khaliqa,

A. and Umar, M. 2015. Preparation and quality characterization of soy milk based non-dairy ice cream. International Journal of Food and Allied Sciences 1(1): 25-31.

- Alleoni, A.C.C. and Antunes, A.J. 2004. Albumen foam stability and s-ovalbumin contents in eggs coated with whey protein concentrate. Revista Brasileira Ciência Avícola 6(2):105-110.
- Alleoni, A.C.C. 2006. Albumen protein and functional properties of gelation and foaming. Scientia Agricola 63(3): 291-298.
- AOAC. 2005. Official Methods of Analysis. The Association of Official Analytical Chemists, Washington DC.
- Butt, M.S. and Batool, R. 2010. Nutritional and functional properties of some promising legumes protein isolates. Pakistan Journal of Nutrition 9(4): 373-379.
- Chaiseri, N.W.T.P.S. and Chonhenchob, V. 2005. Effect of chemical reagents on functional properties of mungbean protein products. Kasetsart Journal (Natural Science) 39: 109- 118.
- Chandra, S. and Samsher. 2013. Assessment of functional properties of different flours. African Journal Agricultural Research 8(38): 4849-4852.
- Keum-IlJang, Han, H.J., Lee, K.Y., Bae, I.Y., Lee, J.Y., Kim, M.K., and Lee, H.G. 2009. Physicochemical properties of mung bean starch paste, a main ingredient of omija-eui. Food Science and Biotechnology 18(4): 991-995.
- Kaushal, P., Kumar, V. and Sharma, H.K. 2012. Comparative study of physicochemical, functional, anti-nutritional and pasting properties of taro (*Colocasia esculenta*), rice (*Oryza sativa*), pigeon pea (*Cajanus cajan*) flour and their blends. LWT-Food Science and Technology 48: 59-68.
- Li, W., Shu, C., Yan, S., and Shen, Q. 2010. Characteristics of sixteen mung bean cultivars and their protein isolates. International Journal of Food Science and Technology 45(6): 1205-1211.
- Peamprasart, T. and Chiewchan, N. 2006. Effect of fat content and preheat treatment on the apparent viscosity of coconut milk after homogenization. Journal of Food Engineering 77: 653-658.
- Phillips, L.G., Whitehead, D.M. and Kinsella, J.E. 1994. Structure-function properties of food proteins. San Diego: Academic Press, 1994, 271p (Food Science and Technology International Series).
- Prapasuwannakul, N., Boonchai, S. and Pengpengpit, N. 2014. Use of green coconut pulp as cream, milk, stabilizer and emulsifier replacer in germinated brown rice ice cream. International Journal of Biological, Veterinary, Agricultural and Food Engineering 8(5): 443-446.
- Raghavendra, S.N. and Raghavarao, K.S.M.S. 2010. Effect of different treatments for the destabilization of coconut milk emulsion. Journal of Food Engineering 97: 341-347.
- Soler, L. 2005. Development of non-dairy frozen dessert containing soy protein and coconut milk. Baton Rouge, United States: The Department of Food Science,

The Louisiana State University and Agricultural and Mechanical College, MS thesis

- Tang, D., Dong, Y., Ren, H., Li, L. and He, C. 2014. A review of phytochemistry, metabolite changes, and medicinal uses of the common food mung bean and its sprouts (Vignaradiata). Chemistry Central Journal 8: 4-16.
- Tangsuphoom, N. and Coupland, J.N. 2008a. Effect of surface-active stabilizers on the microstructure and stability of coconut milk emulsions. Food Hydrocolloids 22: 1233-1242.
- Tangsuphoom, N. and Coupland, J.N. 2008b. Effect of pH and ionic strength on the physicochemical properties of coconut milk emulsions. Journal of Food Science 73(6): E274-80.
- Tangsuphoom, N. and Coupland, J.N. 2009. Effect of thermal treatments on the properties of coconut milk emulsions prepared with surface-active stabilizers. Food Hydrocolloids 23: 1792–1800.
- The Indonesian Academy of Science. 2000. Milk Coconut Pasta. Centre for Documentation and Scientific Information, Indonesia.
- Tipvarakarnkoon, T. 2009. Material science properties of coconut milk, cheese, and emulsion. Berlin: Institute of Food Rheology, Technology University of Berlin, dissertation.

Potential of coconut milk and mung bean extract combination as foam stabilizer in non-dairy ice cream

ORIGIN	ALITY REPORT			
% SIMILA	9 ARITY INDEX	% 13 INTERNET SOURCES	% 14 PUBLICATIONS	% 6 STUDENT PAPERS
PRIMAR	Y SOURCES			
1	ifrj.upm.e	edu.my		%3
2	journal.cl	nemistrycentral.o	com	%
3	Li-Chan, spectroso dithiothre Journal o 1991. Publication	Eunice, and Shu copic study of th aitol induced gela of Agricultural an	uryo Nakai. "Ra ermally and/or ation of lysozyn d Food Chemis	ne", stry,
4	WWW.SCIE	elo.br		%1
5	Kisambira Byaruhar Tumwega "Compos Bean (Pa Nutrition	a, Abbas, John H nga, Phinehas T amire, and Wolfg sition and Function achyrhizus spp.) Sciences, 2015.	H. Muyonga, Yu ukamuhabwa, S gang J. Grüneb onal Properties Seed Flour", F	usuf B. % 1 Silver erg. of Yam ood and

6	Ignasius Radix A.P. Jati. "Black Soybean Seed", Elsevier BV, 2020 Publication	% 1
7	Chavan, U "Functional properties of protein isolates from beach pea (Lathyrus maritimus L.)", Food Chemistry, 200108 Publication	% 1
8	Tangsuphoom, N "Effect of thermal treatments on the properties of coconut milk emulsions prepared with surface-active stabilizers", Food Hydrocolloids, 200910 Publication	% 1
9	Submitted to Higher Education Commission Pakistan Student Paper	%1
10	Arogundade, L.A "Effect of NaCI and its partial or complete replacement with KCI on some functional properties of defatted Colocynthis citrullus L. seed flour", Food Chemistry, 200402 Publication	% 1
11	Shu-Lai Liu. "Original article: Super-chilling maintains freshness of modified atmosphere- packaged Lateolabrax japonicus : Super-chilling maintains freshness of fish", International Journal of Food Science & Technology, 09/2010 Publication	% 1



13	Vipa Surojanametakul, Hirotoshi Doi, Haruki Shibata, Tasuku Mizumura et al. "Reliable Enzyme-Linked Immunosorbent Assay for the Determination of Coconut Milk Proteins in Processed Foods", Journal of Agricultural and Food Chemistry, 2011 Publication	% 1
14	depositonce.tu-berlin.de	%
15	Chee C. Seow, Choon N. Gwee. "Coconut milk: chemistry and technology", International Journal of Food Science & Technology, 2003 Publication	% 1
16	WWW.rjm-cs.ro Internet Source	% 1
17	psasir.upm.edu.my Internet Source	% 1
18	Ana Cláudia Carraro Alleoni. "Albumen protein and functional properties of gelation and foaming", Scientia Agricola, 2006 Publication	%1
19	Submitted to University of Queensland Student Paper	%

Joshi, M., P. Aldred, J.F. Panozzo, S. Kasapis, and B. Adhikari. "Rheological and microstructural characteristics of lentil starchlentil protein composite pastes and gels", Food Hydrocolloids, 2013.

<%1

Publication

20

21	id.123dok.com Internet Source	<%1
22	worldwidescience.org	<%1
23	www.j3.jstage.jst.go.jp	<%1
24	Nebahat Sule Ustun, Sadettin Turhan. "Chapter 10 Antifreeze Proteins in Foods", Springer Science and Business Media LLC, 2020 Publication	<% 1

EXCLUDE QUOTES	ON	EXCLUDE MATCHES	< 10
EXCLUDE BIBLIOGRAPHY	ON		WORDS