

Side Effects of Antipsychotics on Schizophrenia Patients : A Literature Review

by Yufita Ratnasari Wilianto

FILE	1-SIDE_EFFECTS_OF_ANTIPSYCHOTICS__BU_YUFI-FF.PDF (332.88K)		
TIME SUBMITTED	27-JAN-2021 05:53AM (UTC+0700)	WORD COUNT	5636
SUBMISSION ID	1494954758	CHARACTER COUNT	36005



Side Effects of Antipsychotics on Schizophrenia Patients : A Literature Review

Yufita Ratnasari Wilianto^{*1}, Yulistiani²

¹Student of Master Clinical Pharmacy, Faculty of Pharmacy, Airlangga University, Surabaya, Indonesia

²Department of Clinical Pharmacy, Faculty of Pharmacy, Airlangga University, Surabaya, Indonesia

ARTICLE HISTORY

Manuscript submission: April 15th 2019

Manuscript acceptance for review: July 2nd 2019

Approval for publication: July 24th 2019

Keywords :

Atypical Antipsychotics; Mental Disorder; Schizophrenia; Side Effects; Typical Antipsychotics.

ABSTRACT

Schizophrenia is a chronic mental disorder that is characterized by distracted mind, behavioral changes, including delusions and hallucinations. The incidence of schizophrenia is 1.25% of the 2.47% of the psychotic disorder population. Pharmacological therapy given to schizophrenia patient is antipsychotic agents that work on the brain neurotransmitters including dopamine, serotonin, and glutamate. Atypical antipsychotic agents are the first-line therapy for schizophrenia, whereas typical antipsychotics are used as the second line when there is no improvement in symptoms. However antipsychotics agents have frequent side effects such as weight gain and extrapyramidal symptoms. Other reported side effects are including increased prolactin level (sexual dysfunction and menstrual irregularity), cardiovascular system disturbance (hypotension and prolonged QT interval), and lipid changes (elevated triglyceride and cholesterol). Consequences of these side effects can lead to poor adherence of patients, disease progressiveness and increase of physical morbidity and mortality. This article uses 17 literatures with literature parameters used over the past ten years. The purpose of this article is to provide a concise review of schizophrenia and discusses the side effects of atypical and typical antipsychotics based on literature reviews.

* Corresponding author: Yufita Ratnasari Wilianto, Fakultas Farmasi Universitas Airlangga, Surabaya, Indonesia. E-mail: yufita18@gmail.com

1. Introduction

Schizophrenia is one of the most frequent mental health problems to be encountered. According to the study by Chang *et al.* in 2017, the study results showed that lifetime prevalence for schizophrenia is 1.25% of total psychotic disorder population.¹ Schizophrenia disorder is characterized by positive symptoms (for instance, delusion, hallucination, uncontrolled mumbling), negative symptoms (such as withdrawal from social life and decreased life motivation) and cognitive impairments.² Additionally, patients are often difficult to differentiate between reality and their thoughts.³

Eugen Bleuler created a term that is called "schizophrenia" in 1991. According to him, annual schizophrenia incidence was 15-20/100.000 with lifetime risk as many as 0.7%, and higher risk was found more in men than in women (1.4:1).³ Study in the United States with sample based on households showed that prevalence of schizophrenia and other psychotic disorders ranged between 0.25% and 0.64%; while estimated prevalence of schizophrenia on non-institutionalized sample ranged between 0.33% and 0.75%.⁴

Pharmacological therapy that was administered on schizophrenia patients was antipsychotic agents that targeted on the change in neurotransmitter in the brain such as dopamine, serotonin, norepinephrine, and glutamate. Antipsychotic agents that were used as the first line treatment is the atypical group, whereas typical antipsychotic agents were used if there is no improvement in symptoms. The most frequent side effects of antipsychotic agents including increased body weight and extrapyramidal syndrome. Other side effects that had also been reported including increased prolactin level (sexual dysfunction and irregular menstruation), cardiovascular system disturbance (hypotension and prolonged QT interval), and alteration of lipid level (increased triglyceride and cholesterol level). The impact of these side effects could lead to patients' disobedience, the progressiveness of other diseases and increased physical morbidity and mortality.⁵

2. Discussion

Clinical Implications of Schizophrenia

Schizophrenia is a complex mental disorder that could impact significantly to an individual and their family.² It is because schizophrenia is a chronic mental disorder that could lead the patient to have a distracted mind, including delusion and hallucination. Additionally, patients often difficult to differentiate between reality and their minds.^{3,6,7} Schizophrenia consists of 5 subtypes as follows:³

- Paranoid schizophrenia, the most common schizophrenia, in which delusion and auditory hallucination are evident.
- Catatonic schizophrenia, which is less frequent type (about 7% cases). Typical symptoms include psychomotor disorder, catatonic, change of behaviour, echolalia (repetition of speech)

and echopraxia (repetition of action).

- Hebephrenic schizophrenia (unorganized), which is early onset of schizophrenia followed by a bad prognosis. Irresponsive behaviour and unpredictable; influenced by inappropriate mood, including exaggerated laugh, inappropriate behaviour and joke are symptoms that can reveal. Additionally, this subtype is often accompanied by delusion and hallucination.
- Residual Schizophrenia (chronic), which is schizophrenia with a history of one of schizophrenia that has been explained above, yet the disorder is currently showing "negative" symptoms and often dominated by cognitive symptoms.
- Simple schizophrenia, is uncommon schizophrenia; whereas negative symptom without exhibit psychotic trait.

Etiopathophysiology of Schizophrenia

Causes of Schizophrenia are as follows:³

- Genetic
Studies showed that schizophrenia is caused by family inheritance (genetic). Relative risk of schizophrenia inheritance from first line biological family is 10%. If both parents have schizophrenia, then the risk to be schizophrenia on their child will be 40%. Schizophrenia concordance is 10% for dizygotic twins and 40-50% on monozygotic twins.⁷
- Neurochemical alteration
Some neurotransmitters have been associated with this disorder. Dopamine, serotonin, norepinephrine, GABA, and glutamate are some neurotransmitters that have shown involvement in schizophrenia pathogenesis (Figure 1).^{7,8}

Figure 1a explains about glutamatergic, GABAergic, and dopaminergic from cortical area and substantia nigra to accumbens nucleus and striatum, and hippocampus. Whereas figure 1b reveals about the effects of NMDA antagonist receptor on this cortex; reduced GABAergic innervations, reduced tonic dopamine and increased of phasic dopamine release in the striatum.

Dopamine role is based on two dopamine hypotheses. First, there is a group of agents that block dopamine function, known as phenothiazine and could reduce psychotic symptoms. Second, amphetamine, which increases dopamine release inducing paranoid psychotic and aggravate schizophrenia, and disulfiram that inhibits hydroxylase of dopamine and exacerbation of schizophrenia.^{6,7}

Glutamate role in schizophrenia is majorly based on a hypothesis that glutamate reduces the function of NMDA glutamate receptors involved in schizophrenia pathophysiology. Besides, excessive serotonin could lead to the positive and negative symptoms of schizophrenia.

Norepinephrine as a neurotransmitter could also be included to cause schizophrenia. Selective neural degeneration in norepinephrine reward system explains anhedonia that occurs in schizophrenia patient. Other neurotransmitters that are involved in the occurrence of schizophrenia are G-Aminobutyric Acid (GABA) which has a regulation effect on dopamine activity and the loss of inhibitory GABAergic neurons could lead to the hyperactivity of dopaminergic neurons.^{6,7}




Figure 1. Role of Dopamine, Glutamate, GABA in Schizophrenia Pathogenesis⁸

Disorder in neurotransmission has yielded fundamental of pathophysiology theory of schizophrenia. This theory mainly focused on excessive or deficiency of neurotransmitters in the brain including dopamine, serotonin, norepinephrine, and glutamate. Other theories stated that the pathophysiology of schizophrenia involves aspartate, glycine, and gamma-aminobutyric acid as part of the neurochemical imbalance in schizophrenia.⁶

Abnormal activities on dopamine receptor particularly D_2 dopamine receptor is suspected to be associated in many of schizophrenia symptoms. Four dopaminergic pathways have been implicated (see Figure 2). The first pathway is the nigrostriatal pathway that is originated from substantianigra and ended in the caudate nucleus. Low dopamine level in this pathway is considered affecting the extrapyramidal system that leads to motoric symptoms. The second pathway is the mesolimbic pathway that spread from ventral tegmental area (VTA) to limbic area, also has a role in the positive symptom of schizophrenia to excessive dopamine level. The third pathway is the mesocortical pathway which is spread from the VTA to the cortex. Negative symptoms and cognitive deficits in schizophrenia are suggested to be caused by low mesocortical dopamine level. The fourth pathway is the tuberoinfundibular pathway from the hypothalamus to the pituitary gland, where decreased or blockade on tuberoinfundibular dopamine increases prolactin and as a

result, may cause galactorrhea, amenorrhea, and reduced libido.^{6,9,10}

Therapeutic Management of Schizophrenia

The therapy used to treat schizophrenia is an antipsychotic agents. Antipsychotic agents are very effective to treat any positive symptoms in acute episodes including hallucination, delusion, and also very useful to prevent relapse. Antipsychotic therapy algorithm for schizophrenia is shown in Figure 3.¹⁰

Antipsychotic consisted of 2 groups including typical antipsychotic agents (conventional/first generation) and atypical antipsychotic agents (second generation). Antipsychotic grouping is shown in Table 1.^{3,7}

Table 1. Atypical and Typical Antipsychotics³

Atypical and typical antipsychotics		
Atypical antipsychotics	Typical antipsychotics	
Clozapine	Phenothiazines	Chlorpromazine,
Risperidone		Fluphenazine,
Quetiapine		Trifluoperazine
Olanzapine	Butyrophenones	Haloperidol
Aripiprazole	Thioxanthenes	Flupentixol,
Ziprasidone		Zuclopenthixol
Amisulpride	Diphenylbutylpiperidines	Pimozide
Paliperidone	Substituted benzamides	Sulpiride

Side Effects of Antipsychotic & Therapy Management

Atypical group of antipsychotic agents has a lot more metabolic side effects such as increased body weight, cardiac arrhythmia, hyperlipidemia, and diabetes which is caused by insulin resistance; whereas typical group of antipsychotic agents has more significant side effects such as extrapyramidal syndromes (EPS), hyperprolactinemia, and tardive dyskinesia, because of stronger dopaminergic effects of the these agents.^{9,3,12,5,10} Table 2 shows a comparison among the side effects of typical and atypical group of antipsychotic agents in causing body weight and extrapyramidal syndromes. Side effect comparison among antipsychotic agents, FGA (first-generation antipsychotics) and SGA (second-generation antipsychotic¹² is showed on Table 3. Prevalence side effects in the first and second generation of antipsychotic drug that occurred in Indonesia is shown on Table 4.

Side Effects of Antipsychotic Agents on Endocrine System

Hyperprolactinemia has been reported up to 87% patients that were administered with antipsychotic agents including risperidone, haloperidol, and paliperidone.^{9,16,6,17} Hyperprolactinemia is clinically defined as plasma prolactin level > 20 ng/ml for male and > 25 ng/ml for female. Hyperprolactinemia could lead to sexual dysfunction, and decrease libido, cause irregular

menstruation and gynecomastia.^{18,19} It is suggested that these side effects are




Figure 2. Pathophysiology of Schizophrenia⁵

caused by antipsychotic agents that block D₂ dopamine receptor in tuberoinfundibular thus remove inhibition of prolactin secretion and increase prolactin plasma level. Aripiprazole or ziprasidone is preferred for patients with increased prolactin level because this antipsychotic agent has a relatively low sexual dysfunction compared to some antipsychotic agents including clozapine, olanzapine, risperidone, and other conventional antipsychotic agents.^{22,23,17}

Therapeutic management to resolve sexual dysfunction side effects is by using dopamine agonist (e.g. bromocriptine and cabergoline) and by using dopamine release agents such as amantadine, although the usage of the agents needs further study. Additionally, these side effects could also be done by sildenafil therapy. Then, recent study showed that more erection that was enough to penetrate is found more in the sildenafil group than placebo group ($p=0.001$).^{16,24}

According to study by Takachiet *al.* in 2017, a study of 83 women, mean prolactin serum level was higher in patients with irregular menstruation than patients without menstruation disorder (89.3 ± 36.9 vs 42.5 ± 54.5 ; 95% confidence interval (CI): 46.8 [19.3 – 74.3]).²⁵ How antipsychotics promote menstruation disorder or stopped menstruation is not fully elucidated. However, dopamine blockade induced hyperprolactinemia is believed to become a major cause. Increased prolactin level promotes hormonal disorder that releases gonadotropin, which in the end causes irregular menstruation and amenorrhea.²⁶

Increased body weight is another significant side effect on patients with antipsychotic agents. This side effect remains to happen in the early therapy of schizophrenia patients with antipsychotic agents and could lead to patient disobedience to consume their antipsychotic agents. Then, these antipsychotic agents could promote some side effects that have been related to increased risk of

diabetes mellitus and cardiovascular and cerebrovascular mortality.^{8,5,27} Clozapine and olanzapine have shown to have the biggest risk of increasing body weight and the highest risk of diabetes mellitus, followed by risperidone and quetiapine. However, risperidone and quetiapine had a minimal risk in increasing body weight.^{9,12,28,6,29}

Table 2. Side Effects Comparison Between Typical and Atypical Antipsychotic in Causing Body Weight and Extrapyramidal Syndromes⁵

Drug	Weight Gain	Extrapyramidal Syndromes
Typical Antipsychotics (First-generation Antipsychotics)		
Chlorpromazine (Thorazine)	++	+++
Fluphenazine (Prolixin)	+	++++
Haloperidol (Haldol)	+	++++
Perphenazine (Trilafon)	+	++++
Thioridazine (Mellaril)	+	+++
Thiothixene (Navane)	+	++++
Atypical Antipsychotics (Second-Generation Antipsychotics)		
Aripiprazole (Abilify)	+	+
Anesapine (Saphris)	+	++
Clozapine (Clozaril)	++++	+
Iloperidone (Fanapt)	++	±
Lurasidone (Latuda)	±	+
Olanzapine (Zyprexa)	++++	++
Paliperidone (Invega)	++	++
Quetiapine (Seroquel)	++	+
Risperidone (Risperdal)	++	++
Ziprasidone (Geodon)	+	++

± = negligible risk;
 + = low risk;
 ++ = moderate risk;
 +++ = moderately high risk;
 ++++ = high risk

Prevalence of metabolic syndrome in schizophrenia is exceptionally high, supported by a meta-analysis of 77 publications that reported the prevalence was

as high as 32,5% (95% CI = 30,1-35%). Based on Adult Treatment Panel III of the National Cholesterol Education Program Criteria estimated that this worrying proportion as many as 49,4% was observed.^{16, 27} This side effect, increased body weight, has been associated with the antipsychotic mechanism of action on 5-HT_{2A} and 5-HT_{2C} serotonin, dopamine D₂ and D₃, H₁ Histamine and M₃ muscarinic receptors. Antipsychotic agents also affect neuropeptide that controls appetite and energy metabolism. Other mechanisms that involve in increasing body weight are

increased leptin level and reduced adiponectin level that was produced in adipose tissue, and alteration ghrelin level which acts on the arcuate nucleus of the hypothalamus to increase appetite and adipose tissue deposition.^{29,27}

Therapeutic management to overcome this increasing body weight as a comorbid of diabetes, can be considered by administering metformin therapy. There is convincing evidence that supported metformin use in antipsychotic side effects management on increasing body weights. Based on two randomized controlled trials




Figure 3. Antipsychotic Therapeutic Algorithm for Schizophrenia¹⁰

Side Effects of Antipsychotic Agents on Central Nervous System

The main extrapyramidal syndromes that are caused by antipsychotic agents are acute dystonia, akathisia, pseudoparkinsonism, and tardive dyskinesia.³² These symptoms could lead to inconvenience, social stigma, and disobedience for patients. Dystonia often leads to disobedience and could threat patient's life. A dystonic reaction usually occurred in patients that were treated with FGA (first generation antipsychotics/typical) and often seen in the younger male. Dystonia can be minimized by using SGA (second generation of antipsychotics/atypical) or by using FGA with lower dose.^{9,6}

Akathisia includes 50% of extrapyramidal syndrome. Akathisia often concurred with dysphoria, which was seen 20% up to 40% patients that were treated with high potent FGA such as haloperidol and

fluphenazine; whereas quetiapine and clozapine are seemed to have the lowest akathisia side effect. Based on a study that was conducted in McGann District Care Teaching Hospital, Shivamogga Institute of Medical Science, prevalence of akathisia is 10% to 20% in patients with atypical antipsychotic therapy; whereas patients with typical antipsychotic therapy with akathisia are about 20% to 52% [33%].

Pseudoparkinsonism is another side effect which is occurred in patients that were treated with antipsychotic therapy FGA mostly seen in women and geriatric patient 15% and 36%, respectively. The mechanism of this antipsychotic side effect on parkinsonism is neurotransmitter imbalance such as hypoactivity of the D₂ dopaminergic neuron, GABA, and acetylcholine, and hyperactivity of muscarinic cholinergic neuron M₄.³⁴

Hypoactivity of dopamine occurs after administration of FGA and SGA therapy. Dopaminergic

neurons on substantianigra activate D1 dan D2 activatedynorphin neurons; while D2 dopaminergic neurons in the caudate nucleus. In the neurons activate GABAergic neurons nucleus, D1 dopaminergic neuron weakly

Table 3. Side Effects of Antipsychotic FGA and SGA⁸

Side Effects	Low-potency FGAs ^a	High-potency FGAs ^b	SGAs Aripiprazole (Ability)	Clozapine (Clozaril)	Olanzapine (Zyprexa)	Quetiapine (Seroquel)	Risperidone (Risperdal)	Ziprasidone (Geodon)
Anticholinergic effects	+++	+	0	+++	+	+	0	0
Dyslipidemia	++	+	0	+++	+++	++	+	0
Extrapyramidal symptoms	+	+++	+	0	+	0	++	+
Hyperprolactinemia	++	+++	0	0	+	0	+++	+
Neuroleptic malignant syndrome	+	++	+	+	+	+	+	+
Postural hypotension	+++	+	+	+++	+	++	++	+
Prolonged QT interval	++ ^c	+	+	+	+	+	+	++
Sedation	+++	+	+	+++	++	++	+	+
Seizures	+	+	+	+++	+	+	+	+
Sexual dysfunction	+++	++	+	+	+	+	++	+
Type 2 Diabetes Mellitus	+	+	+	++	++	+	+	+
Weight Gain	++	+	0	+++	+++	++	++	0

Note : 0 = rare; + = lower risk; ++ = medium risk; +++ = higher risk.

FGAs = first-generation antipsychotics; SGAs = second-generation antipsychotics.

* --- Effect are approximate, and relative to other antipsychotic medications rather than absolute risk of an adverse effects occuminng.

a --- FGAs with lower potency dopamine D₂ neuroreceptor blockade, including chlorpromazine and thioridazine.

b --- FGAs with higher potency dopamine D₂ neuroreceptor blockade. These include fluphenazine, haloperidol (formely haldol), thiothixene (Navane), and inflluoperazine. Please note that the FGA perphenazine is considered to have intermediate dopamine D₂ neuroreceptor blockade, with an adverse effect profile between the low and high-potency FGAs.

c --- individually, thioridazine has a higher risk of prolonged QT interval and should be used only when no other appropriate options are available.

12

Adapted with permission from Gardner DM, Baldessarini RJ, Waraich P. Modern antipsychotic agents : a critical overview. CMAJ. 2005; 172(13):1703-1711, with additional information from reference 6.

in external globuspallidus. In internal globuspallidus, M4 muscarinic cholinergic neurons which are located in the putamen can increase hyperactivity of acetylcholine. In putamen, there is antagonist interaction between D2 dopaminergic neurons and 5-HT2A serotonin neurons by glutamatergic neurons that inhibit presynaptic NMDA. Thus, SGA with the antagonist effect of 5-HT2A in lower level could promote extrapyramidal syndrome.³⁴

Tardive dyskinesia risk ranged between 0.5% and 62%. This side effect occurred in patients that were treated with long-term typical group antipsychotic therapy and the risk continued to increase in older patients. Risk of tardive dyskinesia significantly lower in patients with atypical group antipsychotic therapy and no case to be reported in patients treated with clozapinemonoth²py. This is because antipsychotic agents such as chlorpromazine, thioridazine, mesoridazine, clozapine, olanzapine, and quetiapine have been reported to be the highest sedation potential²

Neuroleptic Malignant Syndrome (NMS) is a rare side effect of antipsychotic therapy, but it is a life-threatening condition. This side effect is found about 0.5% to 1.0% in patients that are treated with FGA therapy. Mental side effects such as psychosis and delirium could also be found in patients with FGA therapy with higher dose or patients with combination therapy with

anticholinergic.⁶Therapy management to treat extrapyramidal side effects, that are often occurred, is showed on Table 6.

2

Side Effects of Antipsychotic Agents on Cardiovascular System

Orthostatic hypotension has been occurred up to 75% of patients with antipsychotic therapy.

Tabel 4. Prevalence of Side effects of Antipsychotic Agents in Indonesia^{13,14,15}

At Bangli Mental Hospital, Bali Province ¹³	At SambangLihum Mental Hospital, South Kalimantan ¹⁴	In the Hospital Inpatient Ward, Grhasia Yogyakarta ¹⁵
Extrapyramidal syndrome (EPS) (33,3%)	Extrapyramidal syndrome (98,3%)	Extrapyramidal syndrome
Tardif dyskenia (42,9%)	Orthostatic hypotension (86,4%)	Hypotension
Insulin resistance and hyperglycemia (19%)	Anticholinergic effect (76,3%)	
Dyslipidemia (4,8%)	Sedation (44,1%)	
	Nausea (27,1%)	
	Insomnia (16,9%)	
	Anorexia (5,1%)	
	Shortness of breath and cough (1,7%)	

Patients with diabetes mellitus, with the previous history of cardiovascular disease, or elder patients had the greatest risk of orthostatic hypotension.^{9,6} Orthostatic hypotension is defined as decreased blood pressure ($>20/10$ mmHg) in 3 minutes on standing position. The antipsychotic agent that frequently causes orthostatic hypotension side effect is aripiprazole. However, some studies implied that orthostatic hypotension was caused by aripiprazole and clozapine. This side effect was mainly because aripiprazole and clozapine were competitive to $\alpha 1$ adrenoreceptor, ⁹ is promote a higher risk of orthostatic hypotension. As a 5-HT_{2A} antagonist and 5-HT_{1A} partial agonist, aripiprazole could increase the risk of orthostatic hypotension.³⁶

Table 5. Metformin Therapy to Overcome Increasing Body Weight^{30,31,29}

Characteristics	Studies			
	Wu et al (2012)		Wang et al (2012)	
Arms	Metformin 1g/day vs placebo		Metformin 1g/day vs placebo	
Trial duration	6 months		12 weeks	
No. of patients (completers/total)	76/84		66/72	
Primary outcome	Restoration of menstruation.		Change in body weight.	
	Change in weight and BMI.			
Result	Metformin	Placebo	Metformin	Placebo
Mean Weight	-2,3 kg	+2,1 kg	-3,3 kg	+2,5 kg
Mean BMI	-0,93 kg/m ²	0,85 kg/m ²	-1,3 kg/m ²	+0,9 kg/m ²

This orthostatic hypotension side effect could be resolved by administrating fludrocortisone or ephedrine therapy if non-pharmacology therapies were failed. However, this administration should be done with caution because of the side effects. If there was still severe and refractory hypotension, it should be considered to cease clozapine therapy by stepping down the dose gradually in 2 weeks to prevent any cholinergic rebound and to switch into other alternative antipsychotics.³⁷

Alteration in electrocardiography, particularly prolonged QTc could be found in a few patients with antipsychotic agents such as thioridazine, ziprasidone, risperidone, clozapine, olanzapine² and haloperidol.³⁸ Widened or prolonged QTc must be monitored during therapy, and antipsychotic agents must be ceased if this interval consistently surpassed 500 msec. Patients with a history of cardiac or cerebrovascular diseases, also patients in diuretic therapy, should take antipsychotic with caution because the risk of prolonged QTc interval.^{9,6}

Side Effects of Antipsychotic Agents on Alteration of Lipid Profile

Patients treated with atypical antipsychotic therapy such as phenothiazine tended to show increasing triglyceride and cholesterol level. Atypical antipsychotic agents that showed a low risk of lipid alteration were risperidone, ziprasidone, and aripiprazole.^{9,6} Antipsychotic effects on lipid alteration had been associated with increased triglyceride and very low-density lipoprotein release from adiposity. Additionally, antipsychotic could promote expression of sterol regulatory element-binding

proteins (SREBP) and very low-density lipoprotein (VLDL) genes.²⁷

Sterol regulatory element-binding proteins (SREBPs) were the centres to control allosteric from any lipid biosynthesis pathway. Notably, SREBP-1 was responsible as metabolism controller of triglyceride and fatty acid whereas SREBP-2 was the main controller of cholesterol metabolism.^{39,40}

Table 6. Therapy to Treat Extrapyramidal Side Effects³⁵

Side effect	Prevention	Treatment
Acute dystonic reactions	<ul style="list-style-type: none"> 1. Select antipsychotic with low rate EPS * Start with low dose * Increase dose slowly and stepwise 	<ul style="list-style-type: none"> * Oral or intravenous application of 1. anticholinergic drug, e.g., 2-5 mg biperiden, if necessary repeat procedure after 30 min, continue with biperiden oral (maximal 12 mg/d).
Parkinsonism	<ul style="list-style-type: none"> * Select antipsychotic with low risk for parkinsonism * Increase dose slowly and stepwise 	<ul style="list-style-type: none"> * Dose reduction or discontinuation of antipsychotic medication * Switch to SGA * Oral application of anticholinergic drug
Akathisia	<ul style="list-style-type: none"> * Select antipsychotic with low risk for akathisia * Increase dose slowly and stepwise 	<ul style="list-style-type: none"> * Dose reduction * Oral application of beta-receptor blocking agent (e.g., propranolol 30-90 mg/dl) * Switch to certain SGAs * Oral application of benzodiazepines * Trial of an anticholinergic or an antihistaminic agent * Application of vitamin B6 * Application of trazodone
Tardive Dyskinesia	<ul style="list-style-type: none"> * Select antipsychotic with low risk for tardive dyskinesia 1. Evaluate risk factors for tardive dyskinesia 	<ul style="list-style-type: none"> * Switch to cozapine (alternatively to certain other SGAs) 1. Administration of vitamin E * (Administration of tiapride) * ECT (only case reports and case series) * Deep brain stimulation (treatment in severe cases) * Pallidotomy (last resort treatment in extremely severe cases)
Neuroleptic malignant syndrome (NMS)	<ul style="list-style-type: none"> 1. Select antipsychotics with low risk for NMS 	<ul style="list-style-type: none"> * Intensive care management * Stop antipsychotic treatment

* (Application of dantrolene I.V (2.5-10 mg/kg body weight daily)

* (Application of lorazepam 4-8 mg I.V/d).

* In single cases ECT

In the liver, excessive SREBP activity could lead to increased circulation of cholesterol, free fatty acid (FFA), and triacylglyceride [8]. Olanzapine, clozapine, and risperidone could promote significant upregulation of SREBP-1 and SREBP-2, and their downstream target genes that directed to increased lipid and cholesterol synthesis. Even single

intraperitoneal injection with clozapine and olanzapine could induce increased FFA serum, followed by hepatic lipid accumulation.⁴⁰

Monitoring Side Effects of Antipsychotic Agents

While on therapy with antipsychotic agents, patients must regularly be monitored by recording therapy effectivity, the side effect of agents, adherence, and physical health of the patients. Besides, they also should be monitored periodically for body weight, lipid and glucose profile, and ECG.³ Guideline to prevent and monitor metabolic alteration side effect on patients with antipsychotic therapy SGA is showed in Table 7.^{9,42,35}

4 Table 7. Monitoring Side Effect of SGA on Metabolic Alteration^{9,42,35}

Assessment parameter	Baseline	4 weeks	8 weeks	12 weeks	Quarterly	Annually	Every 5 years
Medical and family history, including cardiovascular disease	x					x	
Weight (body mass index)	x	x	x	x	x		
Waist circumference	x					x	
Blood pressure	x			x		x	
Fasting plasma glucose level	x			x		x	
Fasting lipid profile	x			x			x

X = more frequent assessments may be warranted based on clinical status

Reprinted with permission from American Diabetes Association; American Psychiatric Association; American Association of Clinical Endocrinologists; North American Association for the Study of Obesity. Consensus development conference on antipsychotic agents and obesity and diabetes. Diabetes Care. 2004;27(2):599.

3. CONCLUSION

Antipsychotic therapy in schizophrenic patients has side effects such as:

1. Extrapyramidal syndrome. This is caused by imbalance of neurotransmitters such as hypoactivity of the D₂ dopaminergic neuron, GABA, and acetylcholine, and hyperactivity of muscarinic cholinergic neuron M4.
2. Weight gain. This is related to the action of antipsychotics on 5-HT_{2A} and 5-HT_{2C} serotonin, dopamine D₂ and D₃, H₁ Histamine and M₃ muscarinic receptor, increased body weight including increased leptin level, reduced adiponectin level, and alteration in ghrelin level.
3. Sexual dysfunction, decreased libido, irregular menstruation, and gynecomastia due to hyperprolactinemia. This side effects are caused by antipsychotic agents that block D₂ dopamine receptor in tuberoinfundibular thus remove

inhibition of prolactin secretion and increase prolactin plasma level.

4. Orthostatic hypotension. This is because antipsychotics were competitive to α₁ adrenoreceptor. Antipsychotics that work as 5-HT_{2A} antagonists and 5-HT_{1A} partial agonists can increase the risk of orthostatic hypotension.
5. Increased level of triglyceride and cholesterol. This relates to increasing triglyceride, sterol regulatory element-binding proteins (SREBP) and very low-density lipoprotein (VLDL) genes.

The impact of these side effects could lead to discomfort and patient disobedience, the progressiveness of other diseases and increased physical morbidity and mortality. Therefore, it is very important to monitor side effects in schizophrenia patients treated with antipsychotics.

4. References

1. Chang WC, Wong CSM, Chen EYH, Lam LCW, Chan WC, Kin RN, Hung SF, et al. Lifetime Prevalence and Correlates of Schizophrenia-Spectrum, Affective, and Other Non-Affective Psychotic Disorders in the Chinese Adult Population. *Schizophrenia Bulletin*, Oxford University Press. 2017.
2. Keating D, McWilliams S, Schneider I, Hynes C, Cousins G, Strawbridge J, et al. *Pharmacological Guidelines for Schizophrenia : A Systematic Review and Comparison of Recommendations for the First Episode*. *BMJ Open*. 2017 Dec; 7:e013881:1-10.
3. Katona C, Cooper C, Robertson M, editors. *Schizophrenia in Psychiatry at a Glance Fifth Edition*. UK: Wiley-Blackwell Publishing, Ltd; 2012.
4. National Institute of Mental Health. *Schizophrenia* [document on the internet]. USA: 2018 [updated May 2018; cited 2018 May 27]. Available from: <https://www.nimh.nih.gov/health/statistics/schizophrenia.shtml>.
5. Patel KR, Cherian J, Gohil K, and Atkinson D. *Schizophrenia : Overview and Treatment Options*. *Journal Pharmaceutical and Theapeutics*. 2014 Sept;39 (9):638-645.
6. Ayano G. *Schizophrenia: A Concise Overview of Etiology, Epidemiology Diagnosis and Management : Review of Literatures*. *Journal of Schizophrenia Research*. 2016 Aug; 3 (2):1-7.
7. Falkai P, Schmitt A, Cannon TD, editors. *Pathophysiology of Schizophrenia. In Schizophrenia : Current Science and Clinical Practice, First Edition*. UK: A John Wiley & Sons, Ltd, Publication; 2011.
8. Muench J and Hamer AM. *Adverse Effects of Antipsychotic Medications*. *American Academy of Family Physicians*. 2010 March; 81 (5):617-622.
9. Fatani BZ, Aldawod R, Alhawaj FA, Alsadah S, Slaiss FR, Alyaseen EN, et al. *Schizophrenia : Etiology, Pathophysiology and Management - A Review*. *The Egyptian Journal of Hospital Medicine*. 2017 Sept; 69(6):2640-2646.
10. Wells BG, DiPiro JT, Schwinghammer TL, DiPiro CV, editors. *Schizophrenia. In Pharmacotherapy Handbook Ninth Edition*. New York: McGraw-Hill Education; 2015.
11. Hamer S and Haddad PM. *Adverse Effects of Antipsychotics as Outcome Measures*. *British Journal of Psychiatry*. 2007 Apr;191(50):64-70.
12. Jafari S, Enright FF, and Huang XF. *Structural contributions of Antipsychotics Drugs to Their Therapeutic Profiles and Metabolic Side Effects*. *Journal of Neurochemistry*. 2012 Nov; 120:371-384.
13. Subramaniam S, Sasmita NP, and Lesmana CB. *PrevalensiEfekSampingFarmakoterapiterhadapPenderitaSkizofrenia di RumahSakitJiwaBangli, Propinsi Bali*. *E-JurnalMedika*. 2018 Jan; 7(1):22-27.
14. Yulianti MD, Cahaya N, and Srikartika VM. *StudiPenggunaanAntipsikotikdanEfekSampingpadaPasienSkizofrenia di RumahSakitJiwaSambangLihum Kalimantan Selatan*. *JurnalSainsFarmasi&Klinis*. 2017 May; 3(2):153-164.
15. Julaeah, Ananda VD, and Pradana DA. *GambaranEfekSampingAntipsikotikPadaPasienSkizofreniaPadaBangsalRawatInap di RS. Grhasia Yogyakarta*. *Farmasains*. 2016 Apr;3(1):35-41.
16. Park Y, Kim Y, and Lee JH. *Antipsychotic-Induced Sexual Dysfunction and Its Management*. *World J. Mens Health*. 2012 Dec; 30(3):153-159.
17. Montejo AL, Montejo L, and Cremades FV. *Sexual Side-Effects of Antidepressant and Antipsychotic Drugs*. *Curr Opin Psychiatry*. 2015 Nov; 28(6):418-423.
18. Aboraya A, Fullen JE, Ponienman BL, Makela EH and Latocha M. *Hyperprolactinemia Associated with Risperidone : A Case Report and Review of Literature*. *Psychiatry*. 2004 Nov; 29-31.
19. Bargiota SI, Bonotis KS, Messinis IE and Angelopoulos NV. *Review Article The Effects of Antipsychotics on Prolactin Levels and Women's Menstruation*. *Schizophrenia Research and Treatment*. 2013 Nov; 1-10.
20. Just M. *The Influence of Atypical Antipsychotic Drugs in Sexual Function*. *Neuropsychiatric Disease and Treatment*. 2015 Jul;11:1655-1661.
21. John J, Thomas RP, Paul S, Jasmina EK and Job K. *Antipsychotics Induced Sexual Dysfunction*. *Journal of Psychiatry*. 2018 Dec; 21(1):1-2.
22. Baggeley M. *Sexual Dysfunction in Schizophrenia : Focus on Recent Evidence*. *Human Psychopharmacol Clin Exp*. 2008 March; 23:201-209.
23. Chiesa A, Leucci V, Serretti A and Ronchi D. *Antipsychotics and Sexual Dysfunction : Epidemiology, Mechanism and Management*. *Clinical Neuropsychiatry*. 2013 March; 10(1):31-36.
24. Voicu V, Medvedovici A, Ranetti AE and Radulescu FS. *Drug-Induced Hypo- and Hyperprolactinemia : Mechanism, Clinical and Therapeutic Consequences*. *Expert Opin Drug Metab Toxicol*. 2013:1-14.
25. Takechi K, Yoshioka Y, Kawazoe H, Tanaka M, Takatori S, Kobayashi M, et al. *Psychiatric Patients with Antipsychotic Drug Induced Hyperprolactinemia and Menstruation Disorders*. *Biol. Pharm. Bull; The Pharmaceutical Society of Japan*. 2017 Jul; 40(10):1775-1778.
26. Seeman MV. *Review Antipsychotic-Induced Amenorrhea*. *Journal of Mental Health*. 2011 Oct; 20(5):484-491.
27. Madhubhashinee, Dayabandara, Hanwella R, Ratnatunga S, Seneviratne S, Suraweera C, et al. *Antipsychotic-Associated Weight Gain : Management Strategies and Impact on Treatment Adherence*. *Neuropsychiatric Disease and Treatment*. 2017 Aug; 13:2231-2241.
28. Volpato AM, Zugno AL, and Quevedo J. *Recent Evidence and Potential Mechanisms Underlying Weight Gain and Insulin Resistance due to Atypical Antipsychotics*. *Revista Brasileira de Psiquiatria*. 2013 Dec; 35(3):295-304.

29. 29. Prajapati AR. *Review Role of Metformin in the Management of Antipsychotic-Induced Weight Gain*. Progress in Neurology and Psychiatry. 2014 Nov;23:38.
30. 30. Wu RR, Jin H, Gao K, Twamley E, Ou JJ, Shao P, et al. *Metformin or Treatment of Antipsychotic-Induced Amenorrhea and Weight Gain in Women With First-Episode Schizophrenia : A Double-Blind, Randomized, Placebo-Controlled Study*. Am J Psychiatry. 2012 Aug; 169(8):813-821.
31. 31. Wang M, Tong JH, Zhu G, Liang GM, Yan HF, and Wang XZ. *Metformin for Treatment of Antipsychotic-Induced Weight Gain : A Randomized, Placebo-Controlled Study*. Schizophrenia Research. 2012 March; 138(54-57):54-57.
32. 32. Divac N, Prostran M, Jakovcevski I and Cerovac N. *Review Article : Second-Generation Antipsychotics and Extrapyramidal Adverse Effects*. BioMed Research International. 2014 Jun:1-6.
33. 33. Kirgaval RS, Revanakar S and Srirangapatna C. *Prevalence of Extrapyramidal Side Effects in Patients on Antipsychotics Drugs at a Tertiary Care Center*. Journal of Psychiatry. 2017 Jul; 20(5):1-5.
34. 34. Werner FM and Covenas R. *Extrapyramidal Symptoms in Patients Treated with Antipsychotic Drugs*. Journal of Bioequivalence & Bioavailability. 2017 May;9(3):412-415.
35. 35. Hasan A, Falkai P, Lieberman J, Glenthøj B, Gattaz WF, Thibaut F, et al. *World Federation of Societies of Biological Psychiatry (WFSBP) Guidelines for Biological Treatment of Schizophrenia - A Short Version for Primary Care*. International Journal of Psychiatry in Clinical Practice. 2017 Jan:1-9.
36. 36. Lin YS, Ho PS and Liang CS. *Severe Orthostatic Hypotension After Adding Low-Dose Aripiprazole to Clozapine*. Arch Clin Psychiatry. 2017 Oct; 44(3):84.
37. 37. Iqbal MM, Swati S, Aneja A, Iqbal MT, Aneja E and Haque S. *Most Common Side Effects of Clozapine and Their Management*. International Journal of Emergency Mental Health and Human Resilience. 2016; 18(4):1-6.
38. 38. Khasawneh FT and Shankar GS. *Review Article Minimizing Cardiovascular Adverse Effects of Atypical Antipsychotic Drugs in Patients with Schizophrenia*. Cardiology Research and Practice. 2014 Feb:1-8.
39. 39. Skrede S, Steen VM and Ferno J. *Antipsychotic-induced Increase in Lipid Biosynthesis : Activation Through Inhibition*. Journal of Lipid Research. 2013 Dec; 54:307-309.
40. 40. Cai HL, Tan QY, Jiang P, Dang RL, Xue Y, Tang MM, et al. *Original Article A Potential Mechanism Underlying Atypical Antipsychotics-Induced Lipid Disturbances*. Translational Psychiatry. 2015 Oct; 5:1-8.
41. 41. Zhang Y, Liu Y, Su Y, You Y, Ma Y, Yang G, et al. *The Metabolic Side Effects of 12 Antipsychotic Drugs Used for the Treatment of Schizophrenia on Glucose : A Network Meta-Analysis*. BMC Psychiatry. 2017;17(373):1-9.
42. 42. Pramyothin P and Khaodhiar L. *Metabolic Syndrome with the Atypical Antipsychotics*. Current Opinion in Endocrinology, Diabetes & Obesity. 2010;17:460-466.

Side Effects of Antipsychotics on Schizophrenia Patients : A Literature Review

ORIGINALITY REPORT

% 19	% 18	% 10	% 8
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	www.tandfonline.com Internet Source	% 4
2	www.ptcommunity.com Internet Source	% 4
3	www.academia.edu Internet Source	% 2
4	www.depressionforums.org Internet Source	% 2
5	austinpublishinggroup.com Internet Source	% 2
6	wileymicrositebuilder.com Internet Source	% 1
7	Submitted to Northwest Iowa Community College Student Paper	% 1
8	www.nature.com Internet Source	% 1

9	www.archivespsy.com Internet Source	<% 1
10	Submitted to Kingston University Student Paper	<% 1
11	www.jstage.jst.go.jp Internet Source	<% 1
12	"Pharmacovigilance in Psychiatry", Springer Nature, 2016 Publication	<% 1
13	Submitted to Virtual Arkansas School Student Paper	<% 1
14	Albert Muh Haur Lee, Chong Guan Ng, Ong Hui Koh, Jesjeet Singh Gill, Salina Abdul Aziz. "Metabolic Syndrome in First Episode Schizophrenia, Based on the National Mental Health Registry of Schizophrenia (NMHR) in a General Hospital in Malaysia: A 10-Year Retrospective Cohort Study", International Journal of Environmental Research and Public Health, 2018 Publication	<% 1
15	jechbeta.bmj.com Internet Source	<% 1
16	Ramon Cacabelos. "Pharmacogenomics of drugs used to treat brain disorders", Expert	<% 1

Review of Precision Medicine and Drug Development, 2020

Publication

17

worldwidescience.org

Internet Source

<%1

EXCLUDE QUOTES ON

EXCLUDE
BIBLIOGRAPHY ON

EXCLUDE MATCHES < 10
WORDS