

PROCEEDING



THE INTERNATIONAL CONFERENCE ON **SAFETY MANAGEMENT OF CENTRAL CYTOTOXIC RECONSTITUTION IN PHARMACY PRACTICE**



**MAY 25TH, 2013
HOTEL GRAND TJOKRO YOGYAKARTA**

**MAY 26TH-27TH, 2013
BETHESDA HOSPITAL YOGYAKARTA**

Published by :
FACULTY OF PHARMACY
AHMAD DAHLAN UNIVERSITY YOGYAKARTA
INDONESIA

ISBN : 978-979-18458-6-1

PROCEEDING

**INTERNATIONAL CONFERENCE ON
CENTRAL MANAGEMENT OF CENTRAL CYTOTOXIC
RECONSTITUTION**

*Grand Cokro Hotel Yogyakarta
May 25th, 2013*

**THE INTERNATIONAL CONFERENCE ON
CENTRAL MANAGEMENT OF CENTRAL CYTOTOXIC
RECONSTITUTION IN PHARMACY PRACTICE
YOGYAKARTA, INDONESIA, 2013**

Editors :

Dwi Utami M.Si., Apt

Iis Wahyuningsih M.Si., Apt

Wahyu Widyaningsih, M.Si., Apt

Nanik Sulistyani, M.Si., Apt

Published by :

**Faculty of Pharmacy, Ahmad Dahlan University
Jln. Prof. Dr. Soepomo, Warungboto, Yogyakarta
Indonesia**

CONTENTS

Welcome address	i-iii
Content	v
Committee of International Conference	ix
Plenary Session (Invited Speakers)	xi
Compliance → adherence → concordance → self-management → quality of life	1
<i>Prof. dr. A.A. Kaptein</i>	
Developing Pharmacists' skills in Medication Reconciliation	3
<i>Dr. Dyah A Perwitasari, Apt., Ph.D</i>	
Traumatic Experience of Adolescent Female in Floods of Cold Lava after The Eruption of Mount Merapi in The Perspective of Growth and Development in Magelang Regency Shelter	5-9
<i>Retna Tri Astuti, Achir Yani S.Hamid, Novy Helena C.D</i>	
Cost Analysis Therapy of Breast Cancer Patients in Prof. Dr. Margono Soekarjo Purwokerto (OP-02)	11-19
<i>Rizki Khotimah, Budi Raharjo, Heny Ekowati</i>	
Pharmacist Counseling Intervension by Oral Can Increase The Patients Adherence and Decrease Systolic Blood Pressure of Ambulatory Hypertension Patients at Internal Disease Polyclinic PKU Bantul Hospital, Indonesia	21-26
<i>Riza Alfian, Akrom, Endang Darmawan</i>	
Decreasing Systolic Blood Pressure Via Increase Patients Adherence By Short Text Messages (SMS) and Usual Care of Pharmacist on Ambulatory Hypertension Management at Internal Disease Polyclinic, PKU Muhammadiyah Bantul Hospital, Indonesia	27-35
<i>Ginanjjar Zukhruf S., Akrom, Endang Darmawan</i>	
The Use of OTC (Over-The-Counter) Drugs in Self-Medication (Swamedikasi) Effort To The Society in Santan Sumberejo	37-40
<i>Elmiawati Latifah</i>	
Impact of Pharmacist Counseling and Booklet Intervention on Patient's Adherence and Systolic Blood Pressure of Ambulatory Hypertension Patients in Internal Disease Polyclinic Pku Muhammadiyah Bantul, Hospital, Indonesia	41-49
<i>Fitri Setyaningsih, Dyah A. Perwitasari, Akrom</i>	

Disorder of Purine and Pyrimidine Nucleotide Metabolism and Its Therapy	51-56
<i>Mulyadi</i>	
Water Fraction of Sambiloto (<i>Andrographis paniculata</i> Nees) Ethanol Extract Efficacy In Inducing The Number Of Macrophage, Neutrophil, And The Level of TNF-α on Wistar Rats	57-63
<i>Wahyu Dewi Tamayanti, Lidwina Tri Kristanti, Hendra Kurniawan, Martha Ervina, Lannie Hadisoewignyo, Lisa Soegianto, Ratna Megawati Widharna</i>	
Acute Toxicity Test of Rambutan Leaf (<i>Nephelium lappaceum</i> L) Extract in Mice	65-69
<i>Tiara Mega Kusuma, Heni Lutfiyati, Septi Wardani</i>	
Evaluation of Antihypertensive Drugs Utilization in Hospitalized Hypertension Patiens (ICD I.15-2) At X Hospital Bantul Yogvyakarta in 2010 and 2011 By ATC/DDD Method	71-76
<i>Oetari, R.A., Akrom, Perwitasari, D.A.</i>	
Utilization Analysis of Antibiotics For Typhoid Fever in Hospitalized Patient In 2010 And 2011 AT X Hospital in Bantul With ATC/DDD Method	77-81
<i>Pandoyo, Rr. S., Untari, S. S. M., Perwitasari, D. A., Akrom</i>	
Isolate of Actinomycetes Code T34 As Antibiotic Producer Againts <i>Staphylococcus aureus</i> and Bioautography Analysis	83-88
<i>Rizqi Kurniasari, Nanik Sulistyani</i>	
TLC Screening for Antioxidant Activity of Henna (<i>Lawsonia inermis</i> L.) Leaf Extract	89-95
<i>Zainab</i>	
Liposom Formulation as a Thymoquinon Nano-Carrier to Increased the Anticancer Activity	97-101
<i>Nuri Ari Efiana, Tedjo Yuwono</i>	
Formulation of Propanolol Cream With VCO (Virgin Coconut Oil) Contained Base	103-108
<i>Prita Dwi Wulandari, Annas Binarjo</i>	
Antioxidant Activity Assay of Etanolic Extract of Sirsak (<i>Annona muricata</i> L) Leaves	109-114
<i>Laela Hayu Nurani</i>	
Anti Angionesis Activity of Ethanol Extract of Green Algae (<i>Spyrogyra</i> sp) Purified With Chorio Allantoic Membrane (CAM) Method	115-121
<i>Wahyu Widyaningsih, Nina Salamah, Hari Susanti</i>	

- Prevalence and Supportive Factors of Geriatric Self Medication in Pharmacies Gunungkidul Regency at May -July 2012** 123-129
Desy Utami Adi Putri, Andriana Sari, Dyah A. Perwitasari
- The Analysis of Quality of Life in Diabetic Patients Consuming Oral Diabetic Agents** 131-136
Dian Asmidawati, Imaniar Noor Faridah, Dyah A. Perwitasari
- Evaluation on The Implementation of Drug Information Service at Pharmacy in Yogyakarta** 137-141
Faridah Baroroh
- Effect of Turmeric (*Curcuma domestica* Val.) Rhizome Ethanolic Extract to Plasma Lipid Peroxide Level on Wistar Rat Induced by Trimethyltin** 143-147
Heni Puji Astuti, Sapto Yuliani
- The Study of Effects Ethanol Extract *Mimosa pudica* L. and *Manihot utilissima* Pohl. As An Antihyperuricemic in Male Poultry With Induced Chicken Liver Juice** 149-156
Vicko Suswidianoro , Vivi Sofia
- The Effect of Infusa of *Zingiber officinale* Roxb To The Ibuprofen Tablet Bioavailability in Male Rabbits** 157-163
Iis Wahyuningsih, Joko Priyanto, Erika Diah
- Sub Chronic Effect of Ethanol Extract of Nutmeg (*Myristica fragrans* Houtt) Seed in Rat Kidney (PP-08)** 165-169
Moch. Saiful Bachri
- Anti Convulsant Effect of Ethyl Acetate Fraction and Unsolved Ethyl Acetate Fraction from Sirsak Leaf (*Annona muricata*, L.) on Pentylentetrazol induced in Mice** 171-178
Didi Rohadi, Moch. Saiful Bachri, Laela Hayu Nurani
- Anti Convulsant Effect of *Centella asiatica* Fractions and Histopatology Study of Liver and Kidney** 179-183
Affair Masnun, Moch. Saiful Bachri, Laela Hayu Nurani
- Anti Diabetic Activity of Ethanol Extract and Chloroform Extract *Annona muricata* leaf in Alloxan Induced Rats** 185-190
Deni Firmansyah, Moch. Saiful Bachri, Nurkhasanah
- Comparison of Spectrophotometric and TLC-Densitometric Technique in Determination of Phytomelatonin in Green Algae (*Spirogyra* sp) Ethanolic Extract** 191-196
Hari Susanti, Wahyu Widyaningsih, Nina Salamah, Beta Zudia Fertaveni, Efi Puspitasari

**The Organizing Committee of International Conference on
Safety Management of Central Cytotoxic Reconstitution
in Pharmacy Practice**

Advisory Board :

Rector of Ahmad Dahlan University

Prof. Dr. Achmad Mursyidi, Apt.

Prof. Dr. Mulyadi, Apt.

Steering Committee :

Dean of Faculty of Pharmacy Ahmad Dahlan University

Technical Committee :

Chairperson : Dr. rer. nat. Endang Darmawan, Apt

Secretary : Andriana Sari, M.Sc., Apt

Treasure : Ana Hidayati, S. Far., Apt

Sub-committee :

Secretariat : - Imaniar Noor Faridah, M.Sc., Apt
- Faridah Baroroh, M.Sc., Apt

Funding, Sponsorship and Exhibition : - Arief Budi Setianto, M.Sc., Apt.
- Nina Salamah, M.Sc., Apt
- Vivi Sofia, M.Sc., Apt

Scientific Program : - Hendy Ristiono, S.Far., Apt
- Dr. Laela Hayu Nurani, M.Sc., Apt
- Moch Saiful Bachri, M.Sc., Ph.D., Apt
- Nining Sugihartini, M.Sc., Apt

Proceeding Editors : - Dwi Utami M.Si., Apt
- Iis Wahyuningsih M.Si., Apt
- Wahyu Widyaningsih, M.Si., Apt
- Dra. Nanik Sulistyani, M.Si., Apt

Logistic : - Hari Susanti, M.Si., Apt
- drh. Sapto Yuliani, M.PH

- Aprillia, S.Si., Apt
- Warsi, M.Si., Apt
- Promotion and Public Relation : - Annas Binarjo, M.Sc., Apt
- Muhammad Muhlis, Sp.FRS., Apt
- dr. Akrom, M.Kes
- Acomodation, Documentation and Conference Services : - Azis Ikhsanudin, S.Si., Apt
- Adnan, S.Si., Apt

**The Organizing Committee of Workshop on
Safety Management of Central Cytotoxic Reconstitution in Pharmacy
Practice**

Chairperson	: Anna Purwaning Rahaju, S.Si., Apt.
Secretary	: Nolen Mayrani Manik, S. Farm., Apt.
Treasure	: Irine Dyah Widiastuti, S. Si., Apt
Scientific Program	: Dra. L. Endang Budiarti, M. Pharm., Apt. V. Susi Purwanti, S. Si., Apt
Acomodation and Documentation	: Adhitya Nugraha Arisada, S. Farm., Apt.
Funding, and Sponsorship	: Dra. P.E. Wardani, MAB., Apt.
Publication	: Yollita Angela Natasya, S. Farm., Apt.
Logistic	: Dra. Ning Raswani, Apt. Yuliana Sri Suwarti, S.Si., Apt.

WATER FRACTION OF SAMBILOTO (ANDROGRAPHIS PANICULATA NEES) ETHANOL EXTRACT EFFICACY IN INDUCING THE NUMBER OF MACROPHAGE, NEUTROPHIL, AND THE LEVEL OF TNF- α ON WISTAR RATS

Wahyu Dewi Tamayanti, Lidwina Tri Kristanti, Hendra Kurniawan, Martha Ervina, Lannie Hadisoewignyo, Lisa Soegianto, Ratna Megawati Widharna

Faculty of Pharmacy, Widya Mandala Catholic University Surabaya

Email : dewffua@gmail.com

Abstract

Background. *The study of the effect of water fraction of Sambiloto (Andrographis paniculata Nees) ethanol extract on the number of macrophage, neutrophil, and level of TNF- α in the body fluid of male Wistar rats after Staphylococcus aureus induction.*

Methods. *Animals were divided into 3 groups, namely: negative control (NC), treatment (T), and positive control (PC) group. The NC, T and PC groups were administered by 0.9% NaCl, water fraction of Sambiloto ethanol extract of 40mg/200g BW dose, and ibuprofen of 7.2 mg/200g BW dose, respectively. The treatment was performed for 7 consecutive days. The number of macrophages, neutrophils, was identified using light microscope whereas the TNF- α level was identified using ELISA.*

Outcome Measured. *The number of macrophage, neutrophil, and level of TNF- α in the body fluid of male Wistar rats after Staphylococcus aureus induction.*

Results. *The number of macrophages in the KC, the T, and the PC groups were 2.33 ± 1.58 ; 3.80 ± 4.06 ; and 2.67 ± 1.55 cells, respectively. The number of neutrophils were 0.83 ± 0.33 ; 0.58 ± 0.32 ; and 1.33 ± 0.38 cells respective to the KN, T, and PC groups. The level of TNF- α were 17.78 ± 11.67 ; 23.48 ± 15.95 ; and 27.90 ± 30.54 pg/ml respective to the KN, the T, and PC groups.*

Conclusion. *This study indicated that water fraction of Sambiloto ethanol extract able to increase the number of macrophage and TNF- α level yet decrease the number of neutrophil in the body fluids of male Wistar rats after Staphylococcus aureus induction.*

Keywords : *macrophages, neutrophils, TNF- α , Herba Sambiloto, ibuprofen*

INTRODUCTION

Inflammation is a local response to infection or tissue injury. It may occur as a local, systemic, acute, and chronic disorder that lead to the pathologic condition. Inflammatory response precedence by the activation of cell mediated immune system. The cells of immune system respond to the foreign substances that invade the body by several mechanisms, such as chemotaxis, in which the immune cells move to the site of infection, followed by an increase of vascular permeability, and subsequently changes of blood osmotic pressure proceed. Those mechanisms facilitate leucocytes migration to the site of infection (Abbas *et al* 2007).

Development of the inflammatory response plays an important role in initiating pathological condition. Therapeutic approaches to reduce the inflammatory response caused (one of them) by the *Staphylococcus aureus* need to be conducted (Baratawidjaja and Rengganis 2012). Medication choice of inflammation may utilize synthetic drugs and/or natural ingredients. The long term uses of synthetic drugs may initiate unwanted effects on the human body. Therefore, natural ingredients are now becoming chosen alternatives as a healing therapy of inflammatory condition. Some examples are the leaves of Salam (*Syzygium polyanthum*), herbs of Sambiloto (*Andrographis paniculata* Nees.), leaves of Cassava (*Manihot utilissima* Pohl), leaves of Red Betel (*Piper crocatum* Ruiz & Pav), leaves of breadfruit (*Artocarpus altilis*), and others.

Sambiloto (*Andrographis paniculata* Nees) is a natural ingredient which is widely studied because of its efficacy as medicine. Sambiloto (AP) herbs have several active compounds. One of the compounds is known as Andrographolide which posses anti-inflammatory activity (Tewari *et al* 2010). Previously, it was identified that Andrographolide in ethanol extract of AP herbs inhibited inflammation in Wistar rats (Evacuansiany and Soebiantoro, no date). Other studies mentioned that AP herbs active as

anti-inflammation by inhibiting the production of radical oxygen of the neutrophil. Moreover, AP inhibited macrophage migration, and production of TNF- α and IL-12 (Lin and Chao 2010) in the animal serum that treated with AP herbs water fraction containing 10% Andrographolide (Raharjo *et al* 2009).

Among the previous studies of AP anti-inflammatory activity, study of AP especially on phagocytic cells including: monocyte, macrophage, neutrophil, and eosinophil have not yet been conducted. We, therefore, identified the AP anti-inflammatory activity in male Wistar rats.

The objective of this study is to conduct a local inflammation by infecting rats with *Staphylococcus aureus* and identify the response of immune system cells that react on the infected bacteria after an hour. The number of phagocytic cells, namely macrophages and neutrophils were measured as parameter. Since the cells will release cytokines following their activation, therefore, levels of cytokines (TNF- α) also be counted. TNF- α is the main cytokine that is released during acute inflammation against bacteria and other microbes by phagocytic cells (Baratawidjaja and Rengganis 2012). The level of TNF- α measured by ELISA (Enzyme Linked Immunosorbent Assay) method. This study identified the effect of AP herb water fraction of ethanol extract compared to ibuprofen on alleviating local acute inflammation after *Staphylococcus aureus* infection.

METHODS

The apparatuses: distillation equipment, percolator, rotary evaporator (Buchi Rotavapor R-124); separating funnel, chamber, ose, tube, centrifuge apparatus (Hettich Zentrifugen), Eppendorf, syringe, pipette volume, light microscope (Olympus); microplate reader (Thermo Scientific Multiskan-Go, USA).

The materials: Andrographidis Herbs that were collected and determined in the Materia Medica, Batu, East Java; ibuprofen; distilled water; ethanol 96%; ammonia; CHCl₃;

HCl 10% v/v; klorhidrik alcohol; FeCl₃, gelatin solution; Steasny reagent; Na acetate; Na hydroxide; ether; acetic acid glacial; Mayer reagent; Dragendorff solution, n-hexane, ethyl acetate; methanol; butanol, acetic acid; kuercetin; ; *Staphylococcus aureus* (ATCC 25923); MSA medium; BaCl₂ 1%; concentrated H₂SO₄; 0.9% NaCl; Giemsa dye; TNF- α ELISA kit (Abcam, USA).

Animal handling

Animal used were 18 male Wistar rats 2 - 3 months of age and approximately 150-200 g of weight with healthy as well as normal activities. Prior to use, rats were adapted for a week to the new environment. The inclusive criteria of animals were: no symptoms of illness and uniform body weight. Prior to the experiment, rats were fasted for 18 hours (Winter in Hadisoewignyo 2010). Rats were divided into 3 groups of 6 animals. Group I was the negative control (NC), group II was the AP water fraction treated (AP), and group III was treated by ibuprofen (PC).

RESEARCH METHODS

Preparation and Fractionation of AP Crude Extracts

The study was preceded by maceration of crude AP powder by 96% ethanol for 24 hours at room temperature. The obtained extract was filtered and evaporated with a rotary evaporator. The condensed extract was dissolved in 50 ml of methanol:water (7:3). Subsequently, the solution was separated by adding 100 ml of n-hexane and shook in a separator flask for 60 minutes. The formed second layer was separated. The n-hexane addition was repeated for three times. The methanol layer was further separated by ethyl acetate. Ethyl acetate was added to the water layers and shook in a separator flask for 60 minutes. Then, the formed second layer was

separated. This process was repeated for three times.

Anti-inflammatory Activity

In order to conduct anti-inflammatory activity, the rats were injected with *Staphylococcus aureus* (SA). The SA preparation preceded with the rejuvenation of SA in MSA medium and incubation at 37°C for 24 hours. The rejuvenated SA, was mixed to 0.9% NaCl. Subsequently, the turbidity of SA-NaCl suspension was compared to Mc Farland I, in which contained 3×10^8 cfu/ml of bacteria. Before injected to the rats, the suspension was incubated for 2 hour at 37° C (Wasito et al 2008).

Rats were allowed to stand for an hour after SA injection before conducting dissection. Subsequently, the rats were allowed to stand for 5 minutes after 0.9% NaCl injection. Rats that were ready for dissection were anesthetized with ether. Peritoneal fluid was taken from the abdominal cavity. The peritoneal fluid was subsequently smeared and fixed by absolute methanol for 5 minutes. Finally, the fixed peritoneal fluid was stained with Giemsa for 20 minutes, rinsed and dried. The stained peritoneal fluid were identified under a light microscope (400x magnification) and counted for the number of macrophages (Kusmardi et al 2006).

To study the number of the neutrophiles, the blood was taken from the heart, then disposed in a tubes contained EDTA. Prior to centrifugation, part of the blood was taken and subjected to be smeared, and stained as the peritoneal fluid. Subsequently, the stained blood was identified under light microscope (magnification of 400x) to count the number of neutrophil cells. The remaining of the blood was centrifugated at 2,000 rpm for 20 minutes. After centrifugation, serum was taken to be stored at -20° C for measuring the level of of TNF- α by ELISA method (Abcam, 2012).

Table I Standardisation of AP Herbs

No.	Parameter	Qualifications (IHD 1979)	Test results
1	Ashes	< 12%	11.13 ± 0.35
2	Water soluble extract	≥ 18 %	17.03 ± 0.38
3	Ethanol soluble extract	≥ 9.7%	13.53 ± 0.10

IHD : Indonesian Health Departemen

Table II. Rf of AP Herbs Extract and Water Fraction under UV (254 and 366 nm) and UV Visible

Extract	UV λ 254 nm		UV λ 366 nm			UV visible		
	Rf	Compounds	Rf	Color	Compounds	Rf	Color	Compounds
AP Exstract	0.77	Flavon	0.60	Blue	Flavanon	0.56	Yellow	Flavanon
	0.87	Isoflavon	0.79	Blue	Flavanon	0.69	Yellow	Flavanon
			0.87	Red	Isoflavon	0.87	Brown	Isoflavon
AP-water Fraction	0.62	Flavanonol	0.6	Grey	Flavanonol	0.56	Brown	Flavanonol
	0.75	Flavon	0.81	Blue	5-deoksiisofl avon	0.69	Brown	Flavanonol
	0.87	Isoflavon						
Andrografolide	0.81	Andrographolide	0.25	Blue	Andrographo lide			

Table III. The Number of Macrophage, Neutrophil, and TNF-α Leve

Groups	Macropage	Neutrophil	TNF-α
Negative control	2.33 ± 1.58	0.83 ± 0.33	17.7 ± 11.67
AP Herbs	3.80 ± 4.06	0.58 ± 0.32	23.48 ± 15.96
Positive control	2.67 ± 1.54	1.34 ± 0.38	27.90 ± 30.54

AP Herbs: AP Water fraction

Table IV. One Way Anova Analysis ($\alpha=0,05$) of Macrophage, Neutrophil, and TNF- α level Between Groups

Note	F	F table	Significancy
Macrophage	0.42	3.59	0.67
Neutrophil	4.85	4.07	0.04
TNF- α	0.07	3.34	0.94

Table V. LSD Analysis of the Number of Neutrophil

Groups	Compared to	Sig.	Note
Negative control	AP Herbs	0.34	NS
	Positive control	0.07	NS
AP Herbs	Negative control	0.34	NS
	Positive control	0.01	S
Positive control	Negative control	0.07	NS
	AP Herbs	0.02	S

NS: not significant S: significant AP Herbs: AP Water fraction

Results

Before used, several parameter of AP water fraction (AP WF) was standardized including level of ashes, level of water soluble extract, and level of ethanol soluble extracts (Table I). The Indonesian Health Department (1979) qualified that level of AP WF ashes, water soluble extract, and ethanol soluble extract were less than 12%, more than 18%, and more than 9.7%, respectively. The levels of crude AP herbs water-soluble extract (17.03 ± 0.38) were not reach the qualification, therefore, the study was continuously undergone subsequent extraction steps used ethanol as the solvent.

This study used 1 kg of dried AP herbs that macerated by ethanol. The macerated extract obtained was 245 g (24.5%). The extract was subsequently undergone organoleptic test for the color, taste, and smell parameter. The ethanol extract of dried AP herbs was dark green in color, bitter in taste and specific in odor which were qualified according to the previous mentioned data (Indonesian Health Departement 1979).

Following the organoleptic test, the ethanol extract was fractionated extracts by water. The fractionated extract was then subjected to Thin Layer Chromatography (TLC) with butanol: acetic acid: water (3: 1: 1) as mobile phase. The TLC plate was dotted and eluated until reacted with ammonium and so produced a yellowish brown color which was a specific indication of the presence of flavonoids (Harborne 1987) (Table II).

Table II showed the results of Rf level of AP extract and water fraction compared to Andrographolide. When observed at λ 254 and 366 nm, the Rf level of AP extracts and water fractions were different to Andrographolide. It was predicted due to the flavonoids remained in both of AP extracts and water fraction, thus the Andrographolide was not clearly stained. The Rf following observation under the UV visible showed no Rf of andrographolide. It was due to the ammonium vapor was reacted solely to the flavonoids while Andrographolide is belongs to the diterpenoid classes (Lin and Chao 2010).

Table III showed the number of macrophages which was higher than the negative and positive control group. It was probably caused by the injected *Staphylococcus aureus* (SA) were produced an exfoliative which was able to activated macrophages, therefore macrophages were stimulated to release TNF- α , IL-6 and pyrogenic toxin superantigens that subsequently induce the activation of T cells and other macrophages to move to the site of infection (Hidayani no date). The result indicated that AP WF treatment may reduce nitric oxide production and increase macrophage activity (Guan, *et al.* 2012). Moreover, the increased number of macrophages was not different statistically compared to the negative control group that may be caused by the AP WF bacteriostatic and bactericidal activity (Anonymous 2005) which may encounter SA following injection, thus prevent the movement of macrophages to the site of infection. The number of macrophages in the positive control group, that contained ibuprofen, was lower than the AP WF group. This was an indication of less anti-inflammatory activity conducted by ibuprofen since approximately 90% of ibuprofen was bound to protein serum. Therefore, it would be more difficult for ibuprofen to penetrate to the cell membrane to produce anti-inflammatory effects (Wilmana and Gan, 2007).

The number of neutrophil of the AP WF group was lower compared to the negative and positive control group which can be explained due to the antioxidant activity of AP WF that may suppress the synthesis of nitric oxide (NO) and thus less NO will be produced and neutrophil adhesion will be inhibited, consequently (Ezeamuzie, *et al.* 2009). In the positive control group, the number of neutrophil was higher than the negative control and the AP WF group. This was explained due to the action of ibuprofen that affect the biosynthesis of several inflammatory mediators such as prostaglandins and leukotrienes (Wilmana and Gan 2007) that stimulate the increased number of neutrophils (Baratawidjaja and Rengganis 2012).

The levels of TNF- α of the AP WF and the positive control groups showed higher level than the negative control group. This is may be caused by macrophages that activated and released TNF- α into the blood circulation (Abbas, *et al.* 2007) since TNF- α is known as the major cytokine in the acute inflammatory response to bacteria and microbes. In acute inflammation condition, the TNF- α and endothelial leukocytes are working in coordination. Thus, the increased of macrophages is accompanied by elevated level of TNF- α (Baratawidjaja and Rengganis 2012).

CONCLUSION

Conclusively, this study indicate that AP WF may increase the number of macrophages in the peritoneal fluid that had been infected by *Staphylococcus aureus* and consequently increase the level of TNF- α in the serum of the Wistar rats. However, decrease the number of netrophils in the seum of the Wistar rats.

REFERENCES

- Abbas, KA, Lichtman, AH, Pillai, S 2007, *Cellular and Molecular Immunology*, 6th edn, Saunders, Philadelphia, p. 267-301.
- Abcam 2012, *Rat TNF alpha ELISA Kit* (ab46070), viewed 5 october 2012, <<http://www.abcam.com>>
- Anonim 2005, *Tanaman Obat Sambiloto*, viewed 22 Pebruary 2013, <http://www.iptek.net.id/ind/pd_tanobat/view.php?id=152>
- Baratawidjaja, KG, and I, Rengganis 2012, *Imunologi Dasar*, 10th edn, Badan Penerbit FKUI, Jakarta, p. 29, 219-228, 259-260.
- Evacuansiany, EW, F, Soebiantoro (no date), *Pemanfaatan Ekstrak Andrographis paniculata Nees dan Aloe vera L. Sebagai Antiinflamasi*, Bagian Farmakologi Fakultas Kedokteran Universitas Maranatha, Bandung.

- Ezeamuzie, CI, AA, Abu-Ghefreh, H, Canatan 2009, *In Vitro and In Vivo Anti-Inflammatory Effect of Andrographolide*, International Immunopharmacology, Vol. 9, p. 313-318.
- Guan, JH, SW, Bor, CL, Wei, SH, Shyh, YL, Chao, TY, Tsung, and HH, Ming 2012, *Antioxidant and Anti-Inflammatory Properties of Longan (Dimocarpus Longan Lour.) Pericap*, Evidence-Based Complementary and Alternative Medicine, Vol. 2012, p. 1-10.
- Hadisoewignyo, L 2010, *Optimasi Formula Tablet Lepas Lambat Natrium Ibuprofen, Uji Antiinflamasi, dan Uji Ulserogenik pada Mukosa Lambung Tikus*, Dissertation, Universitas Gadjah Mada, Yogyakarta..
- Harborne, JB 1987, *Metode Fitokimia*, Penerbit ITB, Bandung .
- Hidayani, M, NW, Nurelly, FS, Ilyas, D Kadir, S, Rimayani (no date), *Etiopatogenesis Staphylococcal Scalded Skin Syndrome*, Fakultas Kedokteran Universitas Hasanuddin, Makasar, p. 1-15.
- Indonesian Health Departmen 1979, *MMI Jilid III*, Health Departemen of Indonesian Republic, Jakarta.
- Kusmardi, K, Shirly and W, Dwita 2006, *Pengaruh Pemberian Ekstrak Etanol Daun Johar (Cassia siamea Lamk.) Terhadap Peningkatan Aktivitas dan Kapasitas Fagositosis Sel Makrofag*, Makalah Kesehatan, Vol. 10, No. 2, p. 89-93.
- Lin, BF, and WW, Chao 2010, *Isolation and Identification of Bioactive Compounds in Andrographis paniculata (Chuanxinlian), Chinese Medicine*.
- Raharjo, SS, E, Listyaningsih, HP, Diding 2009 *Pengaruh Ekstrak Sambiloto (Andrographis paniculata Nees) Terhadap Kadar Histamin Serum dan Gambaran Histologis Saluran Pernafasan Mencit Balb/c Model Asma Alergi*, Jurnal Bahan Alam Indonesia, Vol. 7, No. 1, p. 1-5.
- Tewari, SK, A, Niranjana, and A Lebri 2010, *Biological activities of Kalmegh (Andrographis paniculata Nees) and its active principle-A review*, Indian Journal of Natural Products and Resources, Vol. 1(2), p. 125-135.
- Wasito, H, SE, Priani, Y, Lumayanti 2008, *Uji Aktivitas antibakteri Madu Terhadap Staphylococcus aureus*, viewed 4 November 2012, <<http://hendriapt.wordpress.com/2008/11/14/uji-aktivitas-antibakteri-madu-terhadap-bakteri-staphylococcus-aureus/>>.
- Wilmana, PF, and S, Gan 2007, *Farmakologi dan Terapi*, 5th edn, Departemen Farmakologi dan Terapeutik Fakultas Kedokteran Universitas, Jakarta, Indonesia, p. 240.