

Diversity of substrate type,
ethnomycology, mineral
composition, proximate, and
phytochemical compounds of
the *Schizopyllum commune* Fr.
in the area along Palu-Koro
Fault, Central Sulawesi,
Indonesia

Submission date: 13-Mar-2023 06:16AM (UTC+0700)
by Ignasius Radix Ap Jati

Submission ID: 2035487354

File name: Diversity_of_substrate_tambahan_P.Radix.pdf (3.09M)

Word count: 13468

Character count: 67922

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Saudi Journal of Biological Sciences

journal homepage: www.sciencedirect.com



Original article

Diversity of substrate type, ethnomycology, mineral composition, proximate, and phytochemical compounds of the *Schizophyllum commune* Fr. in the area along Palu-Koro Fault, Central Sulawesi, Indonesia



Yusran Yusran^{a,*}, Erniwati Erniwati^a, Akhmad Khumaidi^b, Ramadanil Pitopang^c, Ignasius Radix Astadi Praptono Jati^d

^a Department of Forestry, Faculty of Forestry, Tadulako University, Jl. Soekarno-Hatta Km.9, Palu, Central Sulawesi 94118, Indonesia

^b Department of Pharmacy, Faculty of Mathematics and Natural Science, Tadulako University, Jl. Soekarno-Hatta Km.9, Palu, Central Sulawesi 94118, Indonesia

^c Department of Biology, Faculty of Mathematic and Natural Science, Tadulako University, Jl. Soekarno-Hatta Km.9, Palu, Central Sulawesi 94118, Indonesia

^d Department of Food Technology, Widya Mandala Surabaya Catholic University, Jl. Dinoyo 42, Surabaya 60265, Indonesia

ARTICLE INFO

Article history:

Received 5 November 2022

Revised 11 January 2023

Accepted 6 February 2023

Available online 14 February 2023

Keywords:

Schizophyllum commune

Substrate

Nutrition

Palu-Koro Fault

Ethnomycology

Nutritional content

ABSTRACT

Schizophyllum commune Fr. is a wild macro fungus species, which is often used as a food source by the indigenous Kaili tribe along the Palu-Koro fault, Central Sulawesi, Indonesia. This fungus has a wide variety in terms of the weathered wood substrate as a place to grow and is found in almost all types of ecosystems. Although its diversity has been investigated, there is no identification of the weathered wood type as a substrate for growth. Some communities in Indonesia have not also known its potential and benefits. Therefore, this research aims to determine the wood type that grows *S. commune* fungus, ethnomycology, mineral composition, proximate, and phytochemical compounds. It was carried out using the descriptive explanatory approach and the fungi location as well as wood substrate sampling, was determined through the purposive sampling technique in forest areas, agroforestry, and community gardens along the Palu-Koro fault, Central Sulawesi. The samples of unknown wood types were through the collection of tree parts, namely twigs, leaves, flowers, and fruits, which were brought to Herbarium Celebense, Tadulako University for identification. Analysis of mineral content, proximate, and fungal phytochemical compounds was carried out based on the method according to the existing protocol. The results showed that 92 types of rotted wood found where the fungus *S. commune* grew, belonged to 36 families. The nutritional content is also good, although it varies based on the type of wood growing media. Therefore, it can be used and processed into various health-beneficial food products. This showed that domestication of the fungus needs to be carried out to support its commercialization as food and medicine in the future.

© 2023 Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Macro fungus is currently gaining much attention due to its use as food ingredients (Dasanayaka and Wijeyaratne, 2017;

Adejoye et al., 2007; Girma and Tasisa, 2018), drugs (Chandrawanshi et al., 2017; Kamalebo et al., 2018; Espejel-Sánchez et al., 2021), application in bioremediation (Das et al., 2021; Malik et al., 2021; Woldemariam, 2019), as well as in increasing crop production (Elsakhawy and El-Rahem, 2020; Owaid et al., 2017). There is also a high public interest and demand for macro fungus as a food source because of their important nutritional and therapeutic value for humans (Singh, 2017; Waktola and Temesgen, 2018). This fungus has a unique texture, aroma, and good taste that is different from food sourced from plants (Alemu, 2014). Some investigations also reported that several species of macro fungus are a source of bioactive compounds and antioxidants (Sánchez, 2017; Sande et al., 2019; Muszyńska et al., 2017; Zeb and Lee, 2021).

* Corresponding author.

E-mail addresses: yusran@untad.ac.id (Y. Yusran), erniwatiii@untad.ac.id (E. Erniwati), ahmadkhumaidi@untad.ac.id (A. Khumaidi), Ramadhanil@untad.ac.id (R. Pitopang), radix@ukwms.ac.id (Ignasius Radix Astadi Praptono Jati).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

<https://doi.org/10.1016/j.sjbs.2023.103593>

1319-562X/© 2023 Published by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Schizophyllum was first described by Fries in 1815 as the species *Schizophyllum commune* (Fries, 1815), which is also known as the split-gill fungus because it resembles a split gill. Furthermore, it was most easily identified and recognized in eastern states hundreds of years ago as a medicine (Hobs, 2005). The name *Schizophyllum* comes from the Latin syllables *Schizo*, which means split, and *phyllum*, meaning lamella (Carreño-Ruiz et al., 2019). The naming of the species "commune" was according to its meaning, general or widespread, and the occurrence of large intra- and inter-population genetic variations through the spread of spores over great distances and genetic drift (James et al., 1999; Chang and Miles, 2004). Furthermore, it is one of the non-timber forest products that has not been used optimally. This fungus is cosmopolitan, lives both saprophytic and pathogenic, and is easily found on rotting wood or trees in forests, agroforestry lands, plantations, and home gardens. It is very popular worldwide due to the ability to grow in sub-tropical and tropical countries (Yim et al., 2013) such as Mexico and Peninsular Malaysia (Takemoto et al., 2010), North East India, Sumatra Island, Indonesia, and Southern Thailand (Preecha and Thonglumnak, 2015). Most people in these countries use the fungus as food or medicine because it contains fiber, carbohydrates, protein, vitamins, lipids, and mineral elements that are good for the body (Adejoye et al., 2007; Krupodorova and Barshteyn, 2015), anti-diabetic (Chandrawanshi et al., 2019), schizophyllan of anti-cancer compounds, tumors, and immune triggers (Hao et al., 2010; Liu et al., 2015; Ooi and Liu, 2000; Tabata et al., 1981; Yamamoto et al., 1981), schizocommunin of bioactive compounds that can inhibit the growth of cervical cancer cells (Filip et al., 2019), cure infectious diseases caused by bacteria, fungi, and viruses (Mirfat et al., 2014; Kakumu et al., 1991; Komatsu et al., 1973). It is used for schizolysin production, where a hemolysin from this fungus can inhibit HIV disease (Han et al., 2010), the anti-pigmentation agent that is used as a natural ingredient in cosmetology products (Abdul Razak et al., 2018), as well as an ethanol producer (Horisawa et al., 2015).

In Indonesia, *Schizophyllum commune* Fr. has different local names, for example, in Kaili, Central Sulawesi, it is often called *Tanggidi* or *Tanggojo* (Yusran et al., 2021, 2022a), *Kulat Kritip* in Dayak Ngaju, Central Kalimantan and *Kulat Pokok Getah* in West Kalimantan (Nion et al., 2012), *Supa* in Sundanese, Baduy tribe, Banten (Khastini et al., 2018), *jamur gigit* in Javanese, *Tirau* on the island of Sumatra (Kusrinah and Kasiamdari, 2015), *kulat indit-jeng* in Sulawesi and Ngawate (Halmahera), *Keho Kaladede* in Ternate (Bisema 1968), *Keho dlole* in Tidore (Anwar et al., 2020) and *jamur gerigit* in West Papua (Nurlita et al., 2021). Currently, different investigations on *S. commune* have been conducted 30 times in 17 provinces in Indonesia (Nurlita et al., 2021), but the weathered wood type has not been identified as a substrate for growth. In several Indonesian regions, there are still communities that do not know its potential and benefits. These include Sulawesi, an island with a high endemism level, especially in Central Sulawesi Province. There is no report on the type of wood growing media, ethnomycology, mineral composition, proximate, and bioactive compounds. Sulawesi Island is one of the hotspots of the Wallacea Zone, where there are about 1,500 endemic vascular plant types and 30% of their original vegetation cover remains (Myers et al., 2000). Due to the huge potential of forests and trees as substrates for the growth of *S. commune* fungi in this province, it is necessary to have a clear record of the wood type diversity that becomes the substrate for the fungus to grow, including the ethnomycological analysis and the nutritional content influenced by the type of wood growing media. Therefore, this research was conducted to determine the type of substrate wood that grows *S. commune* fungus, the ethnomycology of its use by indigenous peoples in the area

along the Palu-Koro fault, Central Sulawesi, mineral composition, proximate, and their phytochemical compounds based on the different type of weathered wood.

2. Materials and methods

2.1. Location

Observation and identification of the type of weathered wood growing media and sample collection of *Schizophyllum commune* were carried out in natural forests, agroforestry lands, or community gardens along the Palu-Koro fault, Central Sulawesi. This main active fault in Central Sulawesi stretches from the Palu Valley to North Luwu Regency for 220 Km (Patria and Putra, 2020; Socquet et al. 2006), with a higher population density than other areas in the province. The Kaili are the indigenous people and the most dominant. Observation and collection of fungus fruiting body samples were carried out starting from the Palu valley towards the south around Lore Lindu National Park, namely Ngata Baru and Bora villages (Sigi Biromaru Sub-district), as well as other villages which include Rarampadende and Mantikole (West Dolo Sub-district), Bangga and Walatana (South Dolo Sub-district), Binangga (Marawola Sub-district), Uwemanje (Kinovaro Sub-district), Pakuli and Simoro (Tanambulava Sub-district), Toro and Bolapapu (Kulawi Sub-district), Bobo and Tongoa (Palolo Sub-district), Kamarora A and Kadidia (Nokilalaki Sub-district), Sedoa and Kaduwaa (North Lore Sub-district) and Talabosa, Lore Peore Subdistrict (Napu Valley) as well as Doda and Lempe, Lore Tengah Subdistrict (Besoa Valley). The map of the research location is presented in Fig. 1.

The mineral and proximate content were analyzed at the Laboratory of Soil Science, Faculty of Agriculture and Laboratory of Animal Nutrition, as well as the Faculty of Animal Husbandry and Fisheries. Moreover, the analysis of phytochemical compounds was carried out at the Pharmacology Laboratory, Faculty of Mathematics and Natural Sciences, Tadulako University, Palu, Central Sulawesi, Indonesia.

2.2. Observation of weathered wood type, where *Schizophyllum commune* grows, collection of fungus sample fruiting bodies and their ethnomycology.

This exploratory research was carried out using the descriptive explanatory approach. The determination of the location for sampling the *Schizophyllum commune* fungus was carried out intentionally (purposive sampling) in forest areas, agroforestry, and community gardens along the Palu-Koro fault, starting from the valleys of Palu, Kulawi, Palolo, Napu, and Besoa, Central Sulawesi. The samples of weathered wood type were identified by interviewing the owner of the agroforestry/garden. Subsequently, for the wood type in the forest and unknown species, it is carried out by collecting the parts of the tree that are still alive such as leaves, flowers, fruit, and seeds in the field, and was brought to the Herbarium Celebense (CEB), Tadulako University for identification. Interviews were conducted with the community in each village, where the fungus was found and collected, to gain knowledge about the type of weathered wood that grows *S. commune* fungus. Its use as food and medicine, as well as the selling price in traditional markets across the villages, was also determined. Timber/tree type naming was based on *Plants of the World Online (POWO)* (2019) and Royal Botanical Garden (<https://powo.science.kew.org/>).




Fig. 1. Research location along the Palu-Koro fault, Central Sulawesi, Indonesia (Green dots).

2.3. Fungus fruiting body collection and processing

The fruiting bodies of the fungi are cleaned of all adhering dirt with a soft brush without washing first. All fungus samples were cut into small pieces with a sharp knife and packaged separately in plastic bags according to the wood-growing media. The fungi were sun-dried when collected from areas, where electricity is not available. However, most of the fungus samples were dried by electric vacuum. Subsequently, the dried fruit bodies were dried again in the oven at 40 °C for 48 h and ground until smooth using a blender. The samples were stored in airtight plastic containing silica gel during the analysis period for further research purposes. Identification of *S. commune* was based on morphological and microscopic observations using the relevant literature, namely Kuo (2019) via https://www.mushroomexpert.com/schizophyllum_commune.html.

2.4. Analysis of proximate, minerals, and phytochemical compounds on fungi

A total of 15 species of the macro fungus *Schizophyllum commune* samples growing on various types of weathered wood as substrates, as well as macro fungi *Lentinus sajor-caju* and *Auricularia auricula-judae* as comparisons were collected from the forests, agroforestry lands, and community gardens in villages along the Palu-Koro fault, Central Sulawesi. The samples of the *S. commune* macro fungus are:

- A = *S. commune* growing on the wood of *Ficus benyamina* L.
- B = *S. commune* growing on the wood of *Mangifera minor* Blume.
- C = *S. commune* growing on the wood of *Leucaena leucocephala* (Lamk) de Wit.

D = *S. commune* growing on the wood of *Aleurites moluccana* L. Wild.

E = *S. commune* growing on the wood of *Tectona grandis* L.f.

F = *S. commune* growing on the wood of *Pinus merkusii* Jung et de Vriese.

G = *S. commune* growing on the wood of *Lannea coromandelica* (Houtt.) Merr.

H = *S. commune* growing on the wood of *Bambusa* sp.

I = *S. commune* growing on the wood of *Cassia siamea* Lamk.

J = *S. commune* growing on the wood of *Corypha* sp.

K = *S. commune* growing on the wood of *Tamarindus indica* L.

L = *S. commune* growing on the wood of *Anacardium occidentale* L.

M = *S. commune* growing on the wood of *Terminalia catappa* L.

N = *S. commune* growing on the wood of *Gmelina arborea* Roxb.

O = *S. commune* growing on the wood of *Manihot utilisima* L.

P = *Lentinus sajor-caju* growing on the wood of *Cashea siamea* Lamk.

Q = *Auricularia auricula-judae* growing on the wood of *Theobroma cacao* L.

2.4.1. Mineral analysis

The mineral content in the sample was analyzed using Atomic Absorption Spectroscopy (AOAC, 2010). The minerals identified are Nitrogen (N), Phosphorus (P), Potassium (K), Calcium (Ca), Magnesium (Mg), Sulfur (S), Sodium (Na), Iron (Fe), Zinc (Zn), and copper (Cu). A total of 2 g of the sample was put into the digestion flask, followed by the addition of 20 mL of 65% HNO₃ and 10 mL of 70% HClO₄ solution, shaken gently, and let stand for 24 h in the acid cupboard as well as digestion. Subsequently, the digested solution was filtered using Whatman paper no. 42 in a measuring flask and deionized water was added to the boundary line of the flask. The mineral content was determined using a standard curve according to the mineral to be analyzed.

2.4.2. Proximate analysis

Proximate analysis was carried out according to the AOAC (2010) method. The water content was tested using the thermogravimetric principle, where 1 g of the sample was put into a weighing bottle and heated in an oven at 105 °C for 1 h, put in a desiccator, and weighed. This process was repeated until the constant weight data were obtained for the last two measurements. Water content was calculated by the equation (weight of the initial sample-weight of the final sample)/weight of the initial sample * 100%.

Fungus protein content was tested by the Kjeldahl method. A total of 1 g of sample was put in a Kjeldahl flask and 25 mL of sulfuric acid, as well as Kjeldahl tablets, were added. The samples were digested at 400 °C for 2 h, cooled, and attached to the Kjeldahl apparatus circuit. The last step is titration to determine the Nitrogen content of the sample and multiplied by a correction factor to obtain the protein content.

Fat content was tested by the Soxhlet method, where a total of 1 g of sample was put into a Soxhlet flask and 25 mL of hexane solution was added. Soxhlet flasks are installed in the Soxhlet appliance suite. The hexane is expected to extract the fat in the sample and collect it at the bottom of the flask. After 4 h, the hexane solution was evaporated, and the flask was placed in the oven and weighed until it obtained a constant weight. The fat content was obtained from the calculation of the final weight-weight of the empty flask/sample weight * 100%.

Ash content was measured by the thermogravimetric method using a muffle furnace. Approximately 1 g of the sample was put in a crucible of known weight, placed into a muffle furnace for ashing, cooled, and weighed to a constant weight. Fiber analysis of fungus samples was carried out using the thermogravimetric

method. A total of 1 g of sample was hydrolyzed with 1.25% H₂SO₄ and 1.25% NaOH. The residue obtained was washed successively with 20 mL of 10% K₂SO₄, 20 mL of hot distilled water, and 20 mL of 96% alcohol. The filter paper with the remaining residue was roasted at a temperature of 105 °C and weighed until a constant weight was obtained.

2.4.3. Phytochemical analysis of fungus samples

2.4.3.1. Fungus sample extraction. The ethanol extract of each fruiting body of the *S. commune* fungus was carried out by maceration extraction, namely by adding 5 l of pro-analytical 96% ethanol (pa) into a glass container containing 500 g of fungus powder. The powder was soaked for 3x24 hours and stirred every 1 × 24 hours (Kartini *et al.*, 2018). Furthermore, the maceration results are accommodated in a container using a filter and funnel which are concentrated (Raaman, 2006).

2.4.3.2. Phytochemical analysis procedure of fungus samples. The procedure for analyzing the phytochemical compounds of fungus samples uses the following method:

a. Flavonoid Test.

A total of 0.5 g of fungus samples that have been dissolved in ethanol were added to 2 mg of Mg powder. This was followed by the addition of 3 drops of concentrated hydrochloric acid (HCl). When a pink, brown, or reddish color is formed, it indicates the presence of flavonoids (Pooja and Vidyasagar, 2016).

b. Phenolic Test

A total of 0.5 g of fungus samples were dissolved in 5 mL of water. Moreover, a few drops of 5% FeCl₃ solution was added, and when there is a dark green or blue color change, it indicates the presence of phenolic compounds (Sahira and Cathrine, 2015; Pandey and Tripathi, 2014).

c. Saponin Test

Saponins can be detected by the foam test in hot water. The stable foam was discovered for 5 min and did not disappear on the addition of 1 drop of 2 N HCl, indicating the presence of saponins (Nurhasnawati *et al.*, 2019).

d. Steroid/Triterpenoid Test

A total of 2 g of fungus samples were added with 25 mL of ethanol, heated, and filtered. The filtrate was evaporated and ether was added, while the ether layer was pipetted and tested on a spot plate. When 3 drops of Liebermann-Burchard reagent are added and a red/purple color was formed, this shows the presence of positive terpenoids and when green, it indicates steroids (Pooja and Vidyasagar, 2016).

e. Alkaloid Test

A total of 0.5 g of fungus samples were dissolved in ethanol, dripped with HCl, and filtered. The filtrate was tested by adding two drops of Mayer and Dragendorff reagent in different test tubes. A positive reaction is indicated by the presence of a precipitate in the Mayer reagent and a red precipitate in the Dragendorff reagent (Pandey and Tripathi, 2014).

f. Tannin Test

The fungus sample was added with 10 mL of distilled water, filtered, and the filtrate was added. Subsequently, a FeCl₃ reagent was added, and the appearance of a dark blue or black color indicated the presence of tannin. A total of 1% gelatin solution containing sodium chloride was added to the sample, which gave a white precipitate, which indicates the presence of tannin (Pandey and Tripathi, 2014; Zohra *et al.*, 2012).

2.5. Data Analysis

All experiments to determine the mineral composition and proximate were carried out in 3 replications. Data analysis used

Table 1
Type of wood as a substrate that grows *S. commune*.

No.	Local Name	Indonesian Name	Scientific Name	Family	Location
1.	Taipa	Mangga	<i>Mangifera foetida</i> Lour.	Anacardiaceae	3 c,d,e,f
2.	Taipa Dodoro	Mangga Dodor	<i>Mangifera minor</i> Blume	Anacardiaceae	a,b,c,d,e,f
3.	Jambu Sera	Jambu Mete	<i>Anacardium occidentale</i> L.	Anacardiaceae	a,b,c,d,e,f
4.	Kau Jawa	Kayu Jawa	<i>Lannea coromandelica</i> (Houtt.) Merr.	Anacardiaceae	a,b,c,d,e,f
5.	Marantaipa	Rawa-rawa Pipit	<i>Buchania arborescens</i> (Blume) Blume	Anacardiaceae	a,b,c,d,e,f
6.	Siuri	-	<i>Koordersiodendron pinnatum</i> (Blanco) Merr.	Anacardiaceae	c,f, 3
7.	Rao	Senkung	<i>Dracontomelon dao</i> (Blanco) Merr. & Rolfe	Anacardiaceae	g,h,i
8.	Kadondo	Kedondong	<i>Spondias dulcis</i> Soland. Ex Forst.f.	Anacardiaceae	a,b,c,d,e,f
9.	-	Kedondong Hutan	<i>Spondias pinnata</i> (L. F) Kurz	Anacardiaceae	a,b,c,d,e,f
10.	Sarekaya	Sirsak	<i>Annona muricata</i> L.	Annonaceae	a,b,c,d,e,f
11.	Sarekaya Ntovau	Srikaya	<i>Annona squamosa</i> L.	Annonaceae	a,b,c,d,e,f
12.	Andolia (Kaili)	Kenanga	<i>Cananga odorata</i> (Lamk) Hook.f. & Thomson	Annonaceae	b,c,f
13.	Lengaru	Pulai	<i>Alstonia scholaris</i> (L.) R. Br	Apocynaceae	a,b,c,d,e,f
14.	-	-	<i>Wrightia pubescens</i> R. Br.	Apocynaceae	f,h,g
15.	Tirontasi (Kaili)	-	<i>Gastonia serratifolia</i> (Miq.) Philipson	Araliaceae	i,j,k,l
16.	Lui	Silar	<i>Corypha</i> sp.	Arecaceae	a,b,c,d,e,f
17.	Kaluku	Kelapa	<i>Cocos nucifera</i> L.	Arecaceae	a,b,c,d,e,f
18.	Wanga	-	<i>Pigafetta elata</i> Becc.	Arecaceae	i,j
19.	Kalosu	Pinang	<i>Pinanga cease</i> Blume	Arecaceae	a,b,c,d,e,f,g,h,i
20.	-	Kelapa Sawit	<i>Elaeis guineensis</i> Jacq.	Arecaceae	c,f,h 3
21.	Nggonau	Aren	<i>Arenga pinnata</i> (Wurmb) Merr.	Arecaceae	a,b,c,d,e,f,g,h,i,j,k,l
22.	-	Durian	<i>Durio zibethinus</i> Merr.	Bombacaceae	c,g,h,i
23.	Guu (Tau Taa Wana)	Cemara Laut	<i>Casuarina equisetifolia</i> L.	Casuarinaceae	a,b
24.	Donggala	Nyamplung	<i>Calophyllum inophyllum</i> L.	Clusiaceae	b,d
25.	Talise	Ketapang	<i>Terminalia catappa</i> L.	Combretaceae	a,b,c,d,f
26.	-	Simpur	<i>Dillenia indica</i> L.	Dilleniaceae	c,f
27.	Marawola	-	<i>Diospyros macrophylla</i> Bl.	Ebenaceae	b,c,e,f
28.	Sapiri	Kemiri	<i>Aleurites moluccana</i> L. Wild.	Euphorbiaceae	e,f,g,h,i,j
29.	Kasubi	Singkong	<i>Manihot utilisima</i> L.	Euphorbiaceae	a,b,c,d,e,f,g,h,i,j,k,l
30.	Karet	Karet	<i>Hevea brasiliensis</i> Muell.Arg	Euphorbiaceae	c,f
31.	Miapo	Macaranga	<i>Macaranga hispida</i> (Blume) Müll.Arg	Euphorbiaceae	b,c,e,f,g,h,i,j
32.	Balintuma	Kerinjing	<i>Bischofia javanica</i> Blume	Euphorbiaceae	b,c,e,f
33.	-	Kareumbi	<i>Homalanthus populneus</i> (Geiseler) Pax	Euphorbiaceae	h,i,j,k, 4
34.	Beru-beru	Flamboyan	<i>Delonix regia</i> (Bojer ex Hook) Raf.	Fabaceae	a,b,c,d,e,f
35.	Johar	Johar	<i>Cassia siamea</i> Lamk.	Fabaceae	a,b,c,d,e,f
36.	-	Nam-nam	<i>Cynometra ramiflora</i> L.	Fabaceae	a,b
37.	Pete	Petai	<i>Parkia speciosa</i> Hassk.	Fabaceae	a,b,c,e,f
38.	Poi	Asam Jawa	<i>Tamarindus indica</i> L.	Fabaceae	a,b,c,d,e,f
39.	Sengon	Jeungjing	<i>Paraserianthes falcataria</i> (L.) Nielsen	Fabaceae	a,b,c,e,f
40.	Tamalanja/ Kauapse	Lamtoro	<i>Leucaena leucocephala</i> (Lamk) de Wit.	Fabaceae	a,b,c,d,e,f,h
41.	Kau Colo	Trembesi	<i>Samanea samman</i> (Jacq.) Merr.	Fabaceae	a,b,c,d,e,f
42.	Gamal	Gamal	<i>Gliricidia sepium</i> (Jacq.) Kunth ex Walp.	Fabaceae	a,b,c,d,e,f,g,h,i
43.	Doda (Kulawi)	Dadap	<i>Erythrina variegata</i> L.	Fabaceae	3 c,d,e,f,g,h
44.	Akasia	Mangium	<i>Acacia mangium</i> Willd.	Fabaceae	a,b,c,d
45.	-	Akasia	<i>Acacia auriculiformis</i> A. Cunn. Ex Benth	Fabaceae	a,b,c,d,e,f
46.	-	Angsana	<i>Pterocarpus indicus</i> Willd.	Fabaceae	a,e,d
47.	Palili	-	<i>Lithocarpus celebicus</i> (Miq.) Rehder.	Fagaceae	g,h,i
48.	Haleka (Besoa)	-	<i>Castanopsis acuminatissima</i> (Blume) Rheder	Fagaceae	j,k,l
49.	Tavanjuka (Kaili)	Melinjo	<i>Gnetum gnemon</i> L.	Gnetaceae	a,b,c,d,e
50.	-	-	<i>Galbulimima belgraveana</i> (F. Muell.) Sprach.	Himantandraceae	a,c,f,g,h
51.	Jati Puti	Gmelina	<i>Gmelina arborea</i> Roxb	Lamiaceae	A,d,c,d,e,f
52.	Avokad	Alpukat	<i>Persea Americana</i> Mill.	Lauraceae	a,b,c,e,f,g,h,j,j
53.	Kau momi	Medang	Chinnamomum javanicum Blume	Lauraceae	g,h
54.	Uru	Cempaka	<i>Magnolia vrieseana</i> (Miq.) Baill ex Pierre	Magnoliaceae	g,h,l,j
55.	Malapoga	-	<i>Toona ciliata</i> M. Roem	Meliaceae	3
56.	Mindi	Mindi	<i>Melia azedarach</i> L.	Meliaceae	a,b,c,d,e,f
57.	Cokolati	Kakao	<i>Theobroma cacao</i> L.	Malvaceae	a,b,c,e,f,g,h,i,j,k,l
58.	Kalibau	Waru	<i>Hibiscus tiliaceus</i> L.	Malvaceae	a,b,c,d,e
59.	Nunu	Beringin	<i>Ficus benyamina</i> L.	Moraceae	a,b,c,e,f,g,h,i,j
60.	-	-	<i>Ficus minahase</i>	Moraceae	3 j
61.	Ganaga	Nangka	<i>Artocarpus heterophyllus</i> Lmk	Moraceae	a,b,c,d,e,f
62.	Kuu	Sukun	<i>Artocarpus altilis</i> (Parkinson) Fosberg	Moraceae	a,b,c,d,e,f
63.	Kamonji	-	<i>Artocarpus elasticus</i> Reinw. Ex Blume	Moraceae	a,b,c,d,e,f
64.	Sule	Serut	<i>Streblus asper</i> Lour.	Moraceae	a,b,c,d,e,f
65.	-	Leda	<i>Eucalyptus</i> sp.	Myrtaceae	h,i,j,k
66.	Maku	Jambu Bol	<i>Syzygium malaccensis</i> (L.) Merr. & Perry	Myrtaceae	4 c,d,e,f
67.	Alicope	Jembolan	<i>Syzygium cumini</i> (L.) Skeels	Myrtaceae	a,b,c,d,e,f
68.	Jambu jembo	Jambu air	<i>Syzygium samarangense</i> (Blume) Merr. & L. M. Perry	Myrtaceae	a,b,c,d,e,f
69.	Pala	Pala	<i>Myristica fragrans</i> Houtt.	Myristicaceae	e,f,g,h,i
70.	Pinus	Tusam	<i>Pinus merkusii</i> Jung et de Vriese	Pinaceae	3
71.	Avo	Bambu duri	<i>Bambusa blumeana</i> Schult.f	Poaceae	a,b,c,d,e,f
72.	Avo	Bambu	<i>Bambusa maculata</i> Widjaja	Poaceae	a,b,c,d,e,f,g,h,i,j,k,l
73.	Tovu	Tebu	<i>Saccharum officinarum</i> L.	Poaceae	a,b,c,d,e,f,g,h,i

(continued on next page)

Table 1 (continued)

No.	Local Name	Indonesian Name	Scientific Name	Family	Location
74.	Perande	-	<i>Macadamia hildebrandii</i> Van Steen.	Proteaceae	a
75.	Balintuma (Kaili)	-	<i>Bischofia javanica</i> Blume	Phyllanthaceae	e,f,g
76.	Aropi (Kaili)	Buni	<i>Antidesma bunius</i> (L.) Spreng	Phyllanthaceae	c,f
77.	-	Mengkudu	<i>Morinda citrifolia</i> L.	Rubiaceae	a,b,c,d,e,f
78.	Jabon	Jabon	<i>Anthocephalus chinensis</i> (Lam.) A. Rich ex Walp.	Rubiaceae	c,f,h
79.	Jabon	Jabon	<i>Anthocephalus macrophyllus</i> (Roxb) Havil	Rubiaceae	c,f,h
80.	-	Kopi Robusta	<i>Coffea robusta</i>	Rubiaceae	b,c,e,f
81.	Bila	Maja	<i>Aegle marmelos</i> (L.) Correa	Rutaceae	a,b,c,d,e,f
82.	Lotu	Matoa	<i>Pometia pinnata</i> J.R. Forster & J.G. Forster	Sapindaceae	a,b,c,d,e,f
83.	Rambutan	Rambutan	<i>Nephelium lappaceum</i> L.	Sapindaceae	a,b,c,d,e,f
84.	Lengkeng	Klengkeng	<i>Dimocarpus longan</i> Lour.	Sapindaceae	a,b,c,d,e,f
85.	Torode (Besoa, Napu)	Bayur	<i>Pterospermum javanicum</i> Jungh.	Sterculiaceae	g, h, i, j, k, l
86.	-	Bayur Sulawesi	<i>Pterospermum celebicum</i> Miq.	Sterculiaceae	g, h, i, j, k, l
87.	Lekatu (Kaili)	Binuang Laki	<i>Duabanga moluccana</i> Blume	Sonnerataceae	f, g, h, i, j, k, l
88.	Tanjung	Tanjung	<i>Mimusops elengi</i> L.	Sapotaceae	a,b,d,f
89.	Nantu	Nyatoh	<i>Palaquium</i> sp.	Sapotaceae	c,f,h, g,h
90.	Nantu	Nyatoh	<i>Palaquium obovatum</i> (Griff.) Engl.	Sapotaceae	c,f,h
91.	-	-	<i>Gironniera subaequalis</i> Planch.	Ulmaceae	ij
92.	Jati	Jati	<i>Tectona grandis</i> Lf	Verbenaceae	a,b,c,d,e,f,h

Description: a = Ngata Baru and Bora villages (Sigi Biromaru Sub-district); b = Karampadende and Mantikole (West Dolo Sub-district); c = Bangga and Walatana (South Dolo Sub-district); d = Binangga (Marawola Sub-district); e = Uwemanje (Kinovaro Sub-district); f = Pakuli and Simoro (Tanambulava Sub-district); g = Toro and Bolapapu (Kulawi Sub-district); h = Bobo and Tongoa (Palolo Sub-district); i = Kamarora A and Kadidia (Nokilalaki Sub-district); j = Sedoa and Kaduwaa (North Lore Sub-district); k = Talabosa, Lore Peore Subdistrict (Napu Valley); l = Doda and Lempe, Lore Tengah Subdistrict (Besoa Valley).

the statistical package for social sciences 20 (SPSS 20) using one-way analysis of variance (Anova) and the smallest significant difference test was carried out at a probability of 5% to examine whether the treatments were significantly different. The results obtained are represented as the mean ± standard deviation.

3. Results

3.1. Diversity of substrate wood type that grows *Schizophyllum commune* and its ethnomycology

Based on observation data in the field, it was discovered that there were 92 types of weathered wood divided into 36 families, which became the substrate for the growth of *Schizophyllum commune* Fr. along the Palu-Koro fault area as shown in Table 1 below:

Out of the 36 wood families mentioned above, the Fabaceae family has the highest number of wood types, namely 13 species which are the substrate for the growth of *S. commune* fungi, followed by the Anacardiaceae with 9 species, 6 species of Arecaceae, Euphorbiaceae and Moraceae, 4 species of Myrtaceae and Rubiaceae, 3 species of Annonaceae, Poaceae, Sapindaceae and Sapotaceae, Apocynaceae, Fagaceae, Lauraceae, Meliaceae, Malvaceae, Phyllanthaceae and Sterculiaceae each as many as 2 species, and the rest are in the family Araliaceae, Bombaceae, Casuarinaceae, Clusiaceae, Combretaceae, Dilleniaceae, Ebenaceae, Gnetaceae, Himantandraceae, Lamiaceae, Magnoliaceae, Myristicaceae, Pinaceae, Proteaceae, Rutaceae, Sonnerataceae, Ulmaceae, and Verbenaceae with 1 species each as shown in Fig. 2. Furthermore, some examples of macro-morphological fruiting bodies of *S. commune* fungi growing on various types of weathered wood are presented in Fig. 3.




Fig. 2. The number of weathered wood types as substrates for *S. commune* fungi in each different family.



Fig. 3. The morphology of the *Schizophyllum commune* macro fungus growing on several species of weathered wood (A = *Pinus merkusii*, B = *Mangifera foetida*, C = *Gmelina arborea*, D = *Macaranga hispida*, E = *Lannea coromandelica*, F = *Melia azedarach*, G = *Samanea saman*, H = *Artocarpus elasticus*, I = *Anacardium occidentale*, J = *Spondias dulcis*, K = *Arenga pinnata*, L = *Theobroma cacao*, M = *Corypha* sp, N = *Bambusa maculate*, O = *Cocos nucifera*, P = *Tectona grandis*).

The *S. commune* fungus has been used by the Kaili tribe along the Palu-Koro fault area and its surroundings for generations. The Kaili community process this fungus in a simple way, for example, stir-fry with chicken eggs, chilies, tomatoes, and salt, making a soup mixed with vegetables such as cabbage, potatoes, chickpeas, or cooked with coconut milk, chili, and seasonings, which is locally called Utadada Tanggidi. Some people from the Lore tribe in the Napu Valley, Poso Regency, also process by washing with clean water, mixing it with chili and salt, finely ground, and eating the fungus with warm rice. When the harvest is abundant, people usually dry the fruit bodies of this fungus, store them for a certain period in jars and cook the fruit bodies again by soaking them in clean

water for a few minutes to bloom and soften the texture. Generally, this fungus is traded in traditional markets that spread in the Palu valley and its surroundings. The fungus is sold at an uncertain price depending on the season, which is cheap during the rainy season because it is easy to find and grows abundantly on rotting wood. Meanwhile, the price is high during the dry season, because people have to look for the fungus deep in the forest, where it grows a lot due to the microclimate conditions that are very supportive of its growth. In the rainy season, the price per bowl (± 250 g) varies from 2,500–5000 IDR (1 US Dollar = 15,000 IDR) and will increase to 10,000–20,000 IDR per bowl during the dry season. This fungus is sold in fresh condition and sometimes in a processed form that



Fig. 4. The *Schizophyllum commune* Fr. fungus in traditional markets along the Palu Koro fault area, Central Sulawesi, the fruit bodies of the fungi are sold in fresh condition (a,b) and Processed Fungus 'Uta Dada' (c) (Photo taken by Yusran Yusran).

Table 2
Mineral composition some samples of *Schizophyllum commune*, *Lentinus sajor-caju* and *Auricularia auricular-judae* growing on various types of weathered wood.

Treatment	Parameters	N (%)	P (%)	K (%)	Ca (%)	Mg (%)	S (%)	Na (%)	Fe (%)	Zn (%)	Cu (ppm)
A	6.23 ± 0.03e	0.30 ± 0.050c	0.106 ± 0.020f	1.23 ± 0.03c	0.237 ± 0.001b	0.14 ± 0.010b	0.14 ± 0.010a	0.03 ± 0.020d	0.06 ± 0.010b	0.007 ± 0.0005b	0.20 ± 0.05ab
B	6.51 ± 0.02g	0.25 ± 0.005b	0.125 ± 0.005h	1.33 ± 0.01d	0.401 ± 0.005c	0.27 ± 0.001b	0.13 ± 0.002e	0.14 ± 0.002j	0.05 ± 0.002b	0.005 ± 0.0001ab	0.20 ± 0.02ab
C	6.52 ± 0.04g	0.28 ± 0.003e	0.099 ± 0.001e	1.13 ± 0.02b	0.215 ± 0.003h	0.05 ± 0.002bc	0.15 ± 0.005f	0.17 ± 0.010c	0.03 ± 0.0005a	0.012 ± 0.0001d	0.20 ± 0.05ab
D	6.17 ± 0.02d	0.34 ± 0.005d	0.128 ± 0.001h	1.55 ± 0.01f	0.493 ± 0.001d	0.400 ± 0.05c	0.17 ± 0.005c	0.09 ± 0.005 k	0.14 ± 0.002d	0.017 ± 0.0001f	0.20 ± 0.01ab
E	6.52 ± 0.05g	0.20 ± 0.010a	0.100 ± 0.010e	1.28 ± 0.01d	0.298 ± 0.002a	0.07 ± 0.002b	0.12 ± 0.005d	0.11 ± 0.002f	0.03 ± 0.0005a	0.012 ± 0.0001d	0.18 ± 0.005a
F	6.49 ± 0.01f	0.23 ± 0.005ab	0.097 ± 0.001e	1.00 ± 0.05a	0.109 ± 0.001b	0.11 ± 0.005b	0.11 ± 0.002f	0.17 ± 0.005a	0.03 ± 0.0005a	0.013 ± 0.0005de	0.21 ± 0.005b
G	6.50 ± 0.05 fg	0.31 ± 0.005cd	0.118 ± 0.001g	1.48 ± 0.01e	0.274 ± 0.002e	0.45 ± 0.005b	0.14 ± 0.005g	0.27 ± 0.005e	0.09 ± 0.005c	0.009 ± 0.0001c	0.20 ± 0.05ab
H	6.51 ± 0.01g	0.25 ± 0.005b	0.110 ± 0.005f	1.47 ± 0.01e	0.330 ± 0.002e	0.26 ± 0.010bc	0.15 ± 0.005a	0.04 ± 0.001h	0.14 ± 0.002d	0.009 ± 0.0005c	0.20 ± 0.01ab
I	6.49 ± 0.02f	0.23 ± 0.003ab	0.074 ± 0.001d	1.01 ± 0.01a	0.314 ± 0.001c	0.25 ± 0.005c	0.10 ± 0.010a	0.04 ± 0.001g	0.05 ± 0.0001b	0.009 ± 0.0005c	0.19 ± 0.002a
J	6.50 ± 0.05 fg	0.31 ± 0.010cd	0.129 ± 0.001h	1.84 ± 0.01e	0.329 ± 0.001e	0.47 ± 0.005c	0.19 ± 0.010c	0.08 ± 0.005h	0.23 ± 0.005e	0.015 ± 0.0001e	0.22 ± 0.01b
K	6.43 ± 0.02f	0.24 ± 0.030b	0.119 ± 0.002g	1.29 ± 0.01d	0.270 ± 0.010b	0.12 ± 0.020bc	0.15 ± 0.020b	0.07 ± 0.002e	0.17 ± 0.025de	0.013 ± 0.0001de	0.19 ± 0.01a
L	5.88 ± 0.02a	0.24 ± 0.010b	0.057 ± 0.004c	1.14 ± 0.06b	0.248 ± 0.016c	0.23 ± 0.010a	0.12 ± 0.003c	0.08 ± 0.008d	0.10 ± 0.015cd	0.014 ± 0.0001e	0.17 ± 0.02a
M	6.24 ± 0.02e	0.21 ± 0.020a	0.025 ± 0.010a	1.09 ± 0.07b	0.201 ± 0.010b	0.12 ± 0.010b	0.12 ± 0.002a	0.03 ± 0.010b	0.08 ± 0.010c	0.013 ± 0.0001de	0.18 ± 0.03a
N	6.64 ± 0.05h	0.24 ± 0.026b	0.037 ± 0.006b	1.17 ± 0.05c	0.357 ± 0.012d	0.37 ± 0.020b	0.12 ± 0.020a	0.03 ± 0.007i	0.14 ± 0.040d	0.015 ± 0.0001e	0.17 ± 0.02a
O	6.09 ± 0.05c	0.25 ± 0.015b	0.115 ± 0.002g	1.12 ± 0.04b	0.241 ± 0.010b	0.39 ± 0.078b	0.11 ± 0.005a	0.04 ± 0.010d	0.07 ± 0.010bc	0.010 ± 0.0008e	0.26 ± 0.02bc
P	6.50 ± 0.05h	0.20 ± 0.040a	0.109 ± 0.005f	1.47 ± 0.11e	0.207 ± 0.003b	0.11 ± 0.035b	0.14 ± 0.030b	0.06 ± 0.030b	0.07 ± 0.010bc	0.004 ± 0.0002a	2.02 ± 0.04c
Q	5.98 ± 0.01b	0.20 ± 0.010a	0.062 ± 0.001c	1.51 ± 0.001e	0.326 ± 0.001d	0.38 ± 0.005b	0.12 ± 0.004a	0.04 ± 0.002h	0.02 ± 0.003a	0.005 ± 0.0003ab	0.20 ± 0.01ab

Description: The mean value is the average of three replications, ± is the Standard Deviation. The numbers followed by the same letter in the same column mean that they are not significantly different in the Least Significant Difference test at the 5% level.

is ready to be consumed, especially processed mixed with coconut milk (Utadada Tanggidi), which is a favorite dish among the Kaili tribe along the Palu-Koro fault as shown in Fig. 4. For the Kaili tribal community, it is the most preferred and is one of the favorite side dishes because, in addition to its delicious taste, the fungus is easy to obtain since the price is cheap.

3.2. Mineral content of the *Schizophyllum commune* fungus growing on various types of weathered wood

The results of the analysis of the mineral content of *Schizophyllum commune*, *Lentinus sajor-caju* and *Auricularia auricular-judae* growing on various types of weathered wood are presented in Table 2.

3.3. Proximate content of *Schizophyllum commune* samples growing on various types of weathered wood

The analysis results of the proximate content of each sample of *Schizophyllum commune*, *Lentinus sajor-caju* and *Auricularia auricular-judae* are presented in Table 3.

These results show that the water content of *Schizophyllum commune* that grows wild on various types of weathered wood is lower than *Lentinus sajor-caju* and *Auricularia auricular-judae*. It depends on the conditions in which it grows, the water needed to live, and the ability to absorb water from the growing substrate. Although the mean water content of *S. commune* is lower, samples E (*S. commune* grown on *Tectona grandis* L.f) and J (*S. commune* grown on *Corypha* sp. wood) had the lowest water content. The highest water content was in sample N, which was not significantly different from the control.

3.4. Content of phytochemical compounds in *Schizophyllum commune* samples growing on various types of weathered wood

The qualitative analysis results of the phytochemical compounds on each sample of *Schizophyllum commune* based on the wood type are presented in Table 4.

The results showed that the content of phytochemical compounds in *S. commune* samples was relatively not different from one another as shown in Table 4. All samples analyzed were positive for flavonoid, alkaloid, tannin, and phenolic compounds, but did not contain saponin or steroid/terpenoid compounds. In particular, flavonoid compounds were not found in samples of fungi growing on *Anacardium occidentale* L wood, which were detected using AlCl₃ reagent and mg powder, or in those on *Bambusa* sp. and *Manihot utilissima* L.

4. Discussion

These study results are greater than the previous survey conducted by Yusran et al., (2021), which discovered only 16 types of weathered wood that became the substrate for this fungus to grow in Lore Lindu National Park, Indonesia. Furthermore, Herliyana et al., (2011) reported that this is a rotting fungus of several types of wood such as Sengon (*Paraserianthes falcataria*), Karet (*Hevea brasiliensis*), Tusam (*Pinus merkusii*), and Mangium (*Acacia mangium*). In this study, the large number of wood species found to be substrates for *S. commune* fungus was due to observations and explorations carried out in a wider area, both in forest areas, agroforestry, community plantations and agricultural land ranging from lowlands to highlands (0- > 1000 m above sea level) along the Palu-Koro fault, Central Sulawesi, Indonesia. Compared to previous studies which were only in the Lore Lindu National Park area, the size of the area in this study made it possible to find a greater num-

Table 3
Proximate content of the *Schizophyllum commune*, *Lentinus sajor-caju* and *Auricularia auricular-judae* growing on various types of weathered wood.

Treatment	Parameter				
	Water content (%)	Fat (%)	Proteins (%)	Fiber (%)	Ash (%)
A	10.14 ± 0.03c	1.00 ± 0.02c	13.29 ± 0.02c	4.07 ± 0.007h	15.99 ± 0.02c
B	11.02 ± 0.02e	1.20 ± 0.02e	14.43 ± 0.03h	1.22 ± 0.03a	5.79 ± 0.01a
C	11.39 ± 0.21f	1.01 ± 0.01c	13.57 ± 0.02d	2.15 ± 0.03d	8.05 ± 0.01a
D	10.52 ± 0.08d	1.36 ± 0.02g	16.07 ± 0.05j	4.48 ± 0.09i	13.88 ± 0.05c
E	8.64 ± 0.13b	1.05 ± 0.02c	13.71 ± 0.03e	1.45 ± 0.08b	16.53 ± 0.06d
F	11.84 ± 0.06g	0.90 ± 0.01b	13.98 ± 0.01f	3.98 ± 0.02h	17.98 ± 0.01a
G	10.01 ± 0.04c	1.29 ± 0.04f	17.00 ± 0.02l	1.45 ± 0.08b	8.69 ± 0.04b
H	11.16 ± 0.05ef	0.82 ± 0.02a	14.81 ± 0.02i	2.00 ± 0.01c	9.22 ± 0.002b
I	10.92 ± 0.03e	1.34 ± 0.04f	17.60 ± 0.05m	2.83 ± 0.04e	10.07 ± 0.02b
J	8.31 ± 0.02a	1.11 ± 0.03d	19.21 ± 0.02o	10.98 ± 0.01k	18.54 ± 0.05e
K	10.15 ± 0.03c	1.30 ± 0.07f	18.58 ± 0.03n	4.90 ± 0.05j	15.02 ± 0.01d
L	12.38 ± 0.09h	0.91 ± 0.02b	13.95 ± 0.07f	3.17 ± 0.01f	16.04 ± 0.11d
M	10.75 ± 0.52de	0.90 ± 0.04b	16.21 ± 0.18k	2.86 ± 0.12e	19.25 ± 0.57e
N	13.24 ± 0.11i	1.25 ± 0.06e	14.18 ± 0.08g	3.57 ± 0.09g	11.75 ± 0.02c
O	11.30 ± 0.04f	1.57 ± 0.09h	13.98 ± 0.08d	2.23 ± 0.07d	14.15 ± 0.24d
P	13.12 ± 0.03i	1.67 ± 0.02i	12.58 ± 0.02b	1.99 ± 0.05l	10.92 ± 0.03b
Q	13.34 ± 0.02i	1.02 ± 0.01c	12.22 ± 0.05a	31.1 ± 0.01m	9.75 ± 0.02b

Description: The mean value is the average of three replications, ± is the Standard Deviation. The numbers followed by the same letter in the same column mean that they are not significantly different on the 5% level of the Least Significant Difference test.

Table 4
The content of phytochemical compounds in samples of *S. commune*, *Lentinus sajor-caju* and *Auricularia auricular-judae* growing on various types of weathered wood.

Treatment	Flavonoid		Ammonia	Alkaloid		Tanin		Phenolic	Saponins	Steroids/Terpenoids
	AlCl ₃	mg powder		Mayer	Dragen dorff	FeCl ₃	Gelatin			
A	+	+	+	+	+	+	+	+	-	-
B	+	+	+	+	+	+	+	+	-	-
C	+	+	+	+	+	+	+	+	-	-
D	+	+	+	+	+	+	+	+	-	-
E	+	+	+	+	+	+	+	+	-	-
F	+	+	+	+	+	+	+	+	-	-
G	+	+	+	+	+	+	+	+	-	-
H	+	+	-	+	+	+	+	+	-	-
I	+	+	+	+	+	+	+	+	-	-
J	+	+	+	+	+	+	+	+	-	-
K	+	+	+	+	+	+	+	+	-	-
L	-	-	+	+	+	+	+	+	-	-
M	+	+	+	+	+	+	+	+	-	-
N	+	+	+	-	+	+	+	+	-	-
O	+	+	-	+	+	+	+	+	-	-
P	+	+	+	+	+	+	+	+	-	-
Q	+	+	+	+	-	+	+	+	-	-

Information: + = Yes, - = None/not detected.

ber of wood species which became substrates for the *S. commune* fungus, because the types of wood produced by trees that grow spread out and are influenced by altitude above sea level. There are certain types of wood that are the substrate for this fungus and only grow at high altitudes, for example *Pigafetta elata* Becc. and conversely, there are wood-producing trees that only grow in the lowlands. These results indicate that this fungus can grow from the lowlands to the highlands of more than 1000 m asl. The macro fungus has the most growing places and is found in almost all ecosystems. Raper and Miles (1958) and Cooke (1961), stated that it is widely found and can grow on rotting dead wood of at least 150 different genera of flowering plants. According to de Jong (2006), the fungus can also colonize softwood and grass silage.

The yield and productivity of macro fungi depend on the species, substrate, and environmental conditions (Teoh et al., 2017). Meanwhile, *S. commune* will grow well on carbon nutrient sources of sucrose, maltose, and others (Niederpruem and Hunt, 1964). These results indicate that the type of substrate wood, where the fungus grows significantly affects the size and color of the growing fruiting bodies. This showed in the fungus growing on the wood of

Lanea coromandelica (Houtt.) Merr. The weathered ones produce fruit bodies that are larger than those grown on other weathered woods and have a grayish color, while the fungus on *Corypha* sp. has a whiter color. Macroscopically, this fungus has the characteristics of small and short fruiting bodies, hairy lamellae, with a white, or grayish color. Dasanayaka and Wijeyaratne (2017) reported that the edible fungus of *S. commune* can grow on a substrate of 7 species of wood, namely *Alstonia macrophylla*, *Artocarpus heterophyllus*, *Harpullia arborea*, *Mangifera indica*, *Dillenia indica*, *Nephelium lappaceum*, and *Terminalia catappa*. Cultivation on *Artocarpus heterophyllus* wood substrate gave the highest production compared to the other 6 wood substrates.

According to Whitten et al., (1987) and Kefler et al. (2002), almost 15% of the 5000 vascular plant species recorded are endemic to the island of Sulawesi, including more than 2100 woody plant species. The forest area along the Palu-Koro fault is also very rich in plant diversity, therefore, it has the potential as a substrate for food and medicinal macro fungi to grow. One of the important conservation areas along the Palu-Koro fault is Lore Lindu National Park. A total of 166 tree species were also found in sub-montane

and lower montane primary forests, Lore Lindu National Park, Central Sulawesi (Culmsee and Pitopang, 2009). The richness of plant species in this area is a huge potential for the growth and production of *S. commune* fungi as well as other species of food and medicinal macro fungi, with their future development efforts.

Generally, the macro fungus can grow on various types of substrates, but the level of use of the substrate and the growth of the fungus depend on the species of the fungus itself (Kumla *et al.*, 2020). Almost all agricultural waste contains lignocellulosic as the main component needed by fungi and based on the type of plant (Suwannarach *et al.*, 2022). This research is the first to report specifically in large quantities about the type of substrate wood, where the *S. commune* fungus grows in Indonesia.

The bioaccumulation of elements in the fungal body and the biological importance of the accumulation process is strongly influenced by many factors, which are still poorly understood. The fundamental factors that have been known include natural geochemical factors from the source rock, environmental pollution by heavy metals, the environment of the fungus, and the bioaccumulation of trace elements in the fruiting bodies (Kozarski *et al.*, 2015). This research shows that the mineral content of the *S. commune* fungus varies based on the type of weathered wood as the substrate growing media. Furthermore, mineral K has the highest content compared to other minerals, followed by Ca, Mg, and Cu. This is due to the high level of potassium in weathered wood, which is the substrate for the fungus. Alananbeh *et al.*, (2014) stated that the highest mineral in *Pleurotus ostreatus* is K, followed by Mg, Ca, and Zn. The presence of K in the highest concentrations was reported in wild fungi that live on weathered wood substrates (Sanmee *et al.*, 2003).

The concentration of Magnesium and Calcium in fungi is also influenced by environmental factors, as reported in Sanmee *et al.* (2003) and Uzun *et al.* (2011). Calcium is needed by fungi for the growth of mycelia, neutralizing acidic substrate conditions, and increasing the calcium content for consumption (Suzuki, 2020). In the human body calcium functions as a form of bone tissue, teeth, and various other organs, which helps muscle tissue contraction, regulates heart rate, and nerve function. Meanwhile, magnesium plays a role in biological oxidation processes and respiration as well as affects blood pressure and blood sugar levels (Rózsa *et al.*, 2021).

Fungi are known as good accumulators for zinc with fairly high content. The zinc levels in this research are equivalent to other reports by Işiloğlu *et al.* (2001) and Altaf *et al.* (2020). This element is required by fungi for their metabolism and used for various physiological functions such as cell growth and regeneration in humans. Zinc deficiency is a big problem and needs attention alongside iron and vitamin A deficiency (Gupta *et al.*, 2020), therefore, fungi can be used to reduce the level of deficiency in humans.

There is a need to observe the high Cu content in *Lentinus sajorajju*, which grows on decayed *Cassia siamea* wood collected from community gardens. It is suspected to be heavy metal contamination from pesticides used by the community in controlling nuisance organisms. Although the concentration of 2 ppm is not harmful to body health (Uzun *et al.*, 2011), in consuming food macro fungi, the public needs to consider the origin and place of collection. This is to determine whether it is from agricultural and plantation areas, where pesticides are very often applied to avoid contaminating the fruit bodies of the collected fungi.

Falandysz and Borovička (2013) reported that wild-grown edible macro-fungi can accumulate minerals essential for humans at greater levels than cultivated fungi as influenced by the growing habitat (Krupodorova and Sevindik, 2020). Furthermore, Bhattacharyya *et al.*, (2014) stated that trace elements such as Zn, Cu, Fe, and Mn, although required in very small amounts, will

have an essential role as a co-factor of enzymes to carry out biological functions in almost all metabolic processes of the human body.

Based on a previous report, the water content can be influenced by several factors, namely harvesting time, maturity level, and environmental climatic conditions (Beluhan & Ranogajec, 2011). The difference in water content in the mushroom samples in this study was more due to climate differences, especially temperature, humidity and rainfall at the locations where these mushrooms were found, which were influenced by the altitude above sea level. The altitude above sea level greatly affects temperature and humidity, which of course greatly affects the growth of fungi. In addition, the research location in the highlands has higher rainfall than in the lowlands (Palu valley) which is known as a dry area. The next factor that affects the moisture content of the mushrooms is the age of the mushrooms when they are harvested. The fruit bodies are harvested without knowing their age because they grow naturally, which of course will have different levels of maturity, which will affect their water content. The water content needs to be considered because at a high value, it is susceptible to damage such as browning, off-flavor, and the growth of microorganisms after harvesting, thereby reducing the shelf life (Niu *et al.*, 2020).

Fungi are foods with high protein content, hence, *S. commune* has a higher protein content compared to the control. Sample J, which is *S. commune* growing on *Corypha* sp. wood had the highest protein content of 19.21%. These results are in line with Ao & Deb (2019) who investigated wild macro fungi in India with high protein content. Ouzouni *et al.* (2009) showed that fungi have a high protein content compared to the type of ingredients commonly consumed as vegetables, such as green leafy and others. Moreover, other results also show that *S. commune* contains 16–27% protein (Aletor 1995, Longvah and Deosthale 1998; Yusran *et al.*, 2022b). High protein content has the potential to be developed into various food products and consumed to increase people's protein intake.

The fat content of the fungus samples showed a range between 0.82% and 1.57%, which is influenced by metabolic factors and food reserves. The low-fat content shows that *S. commune* can be developed as a low-fat healthy food. This is in line with other research, which explains that fungi have low-fat content (Jacinto-Azevedo *et al.*, 2021; Fogarasi *et al.*, 2018). Moreover, fungi are widely processed for various food products and are believed to provide many health benefits (González *et al.*, 2020). Low levels of fat are reported to loss of weight and the risk of increasing blood pressure (Sun & Niu, 2020). The fiber content of *S. commune* with different growth substrates was in the range of 1–10%, which is lower compared to the control. According to Ao & Deb (2019), fungi are good for human health because they are rich in fiber. This present research proves that fungi with various substrates of weathered wood as growth media have good nutritional levels, therefore, they can be used and processed into several food-beneficial products.

Similarly, Herawati *et al.*, (2021) discovered that the macro fungus *S. commune* that grows wild is positive for flavonoid compounds, steroids, tannins, and coumarins. Okwulehie *et al.*, 2007 also detected the presence of alkaloids, flavonoids, phenols, saponins, and tannins in the *S. commune* growing on rotten mango (*Mangifera indica*) wood. This is different from Acanto and Helen Cuaderes, (2021), who stated that *S. commune* growing in the forest of Minapasuk, Calatrava, Negros Occidental, Philippines contains saponin compounds. Furthermore, Wirth *et al.*, (2021) showed that the fungus contained terpenoid compounds in form of sesquiterpenes.

According to Basso *et al.*, (2020), the content of phytochemical compounds of macro fungus differs based on the type of weathered wood as a substrate for growth media. Furthermore, the detection of phytochemical compounds in the sample is also influenced by several factors, including the solvents and reagents (Shaikh and

Patil, 2020). Teoh and Don (2013) analysis using methanol as an extraction solvent discovered the presence of flavonoid compounds, saponins, and phenols in this fungus, but did not contain alkaloids. Moreover, *S. commune* can produce antioxidant compounds through the shikimate pathway from the central metabolism that generates chorismate precursors. This can eventually form secondary anti-oxidant metabolites such as isoflavones, flavones, flavonols, anthocyanins, tannins, coumarin derivatives, and other phenolic compounds (Boonthatui *et al.*, 2021). The composition of the phenolics content in macrofungi is also influenced by genetic and environmental factors (Yildiz *et al.*, 2017), strains or species of macrofungi, the composition of growing media, harvest time, management techniques, handling conditions, and preparation of substrate for their cultivation, as well as soil/substrate composition or host associated with the wild fungi species (Heleno *et al.*, 2010).

Based on the previous research, it was reported that *S. commune* with its phytochemical compounds has pharmacological activity. Stan *et al.*, (2021) stated that the aqueous extract has anti-dengue activity, while the ethanolic extract has strong antioxidant activity related to phenolic compounds (Basso *et al.*, 2020; Mongkontanawat and Thumrongchote, 2021). The potential immunostimulator and antitumor activity of glucan-associated *S. commune* have also been reported (Vannucci *et al.*, 2013). Fungal antioxidants can vary based on the fruiting body, mycelium, and growth medium (Dulay *et al.*, 2016).

5. Conclusion

This research showed that *Schizophyllum commune* Fr. grows on 92 types of woods that had rotted. This fungus has variations in the color, size, and production of the fruiting bodies, which depend on the weathered wood substrate of the growing medium. The mineral content, proximate, and phytochemical compounds of the macro fungus *S. commune* also varied depending on the type of wood growing media. The results are very useful for local people who are malnourished and have low incomes to consume wild food fungi such as Tanggidi (*S. commune*) because of its high nutritional content and antioxidants. The fungus has affordable selling prices and can be found growing naturally on various types of weathered wood in the area along the Palu-Koro fault, Central Sulawesi, Indonesia. It was also discovered that macro fungi with various decayed wood substrates as growth media have good nutritional levels, therefore, they can be used and processed into various food products that are beneficial for health. Based on these results, it is recommended to carry out chemical compounds analysis on each type of wood to realize the prospect of its use as a substrate in the cultivation of *S. commune* fungi. Therefore, there is a need to domesticate this fungus to support its commercialization as food and medicine in the future.

Funding information

This research was funded by the Directorate General of Research and Development Strengthening, Ministry of Research, Technology and Higher Education, Indonesia, through the Higher Education Applied Research Grant (PTUPT), No. SP DIPA-042.06.1.401516/2019, 5 December 2018.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors are grateful to all participants, namely Aprianto Manggu, Wili Yanto Andisi, Rizal Munandar, Gilno Vanderlik Waro, Angelius Akmon, Robi Gembu, and Abdul Gafur who assisted during the experiment, especially in data and sample collection in the field as well as mineral and proximate analysis of fungus samples in the laboratory. The authors are also grateful to Haris Priyana for the assistance in completing the research site map.

References

- Abdul Razak, D.L., Jamaluddin, A., Abd. Rashid, N.Y., Mohd Fadzil, N.A., Sani, N.A., Abdul Manan, M., 2018. Comparative Evaluation of *Schizophyllum commune* Extracts as Potential Cosmeceutical Bio-Ingredient. *Int. J. Agric. Sci. 5* (1), 2348–3997.
- Acanto, R.B., Helen Cuaderes, V.S., 2021. Antimicrobial Activity And Phytochemical Screening of Split Gill Mushroom (*Schizophyllum commune*) Ethanolic Extract. *Int. J. Sci. Technol. Res.* 10, 4–9.
- Adejoye, O.D., Adebayo-Tayo, B.C., Ogunjobi, A.A., Afolabi, O.O., 2007. Physicochemical Studies on *Schizophyllum commune* (Fries) a Nigerian Edible Fungus. *World Appl. Sci. J.* 2, 73–76.
- Alanabeh, K.M., Bouqellah, N.A., Al Kaff, N.S., 2014. Cultivation of oyster mushroom *Pleurotus ostreatus* on date-palm leaves mixed with other agro-wastes in Saudi Arabia. *Saudi J. Biol. Sci.* <https://doi.org/10.1016/j.sjbs.2014.08.001>.
- Alemu, F., 2014. Cultivation of *Lentinus edodes* on teff straw (agricultural residue) at Dilla University, Ethiopia. *Appl. Microbiol.* 1, 49–59.
- Aletor, V.A., 1995. Compositional studies on edible tropical species of mushrooms. *Food Chem.* 54, 265–268.
- Altaf, U., Lalotra, P., Sharma, Y.P., 2020. Nutritional and mineral composition of four wild edible mushrooms from Jammu and Kashmir, India. *Indian Phytopathol.* 73, 313–320. <https://doi.org/10.1007/s42360-020-00230-1>.
- Anwar, R., Nasichah, A.Z., Roini, C., 2020. Pengetahuan masyarakat Kecamatan Tidore Utara tentang pemanfaatan makroskopis sebagai potensi lokal daerah. *Saintifik@: Jurnal Pendidikan MIPA* 6, 86–92.
- Ao, T., Deb, C.R., 2019. Nutritional and antioxidant potential of some wild edible mushrooms of Nagaland, India. *J. Food Sci. Technol.* 56 (2), 1084–1089. <https://doi.org/10.1007/s13197-018-03557-w>.
- AOAC, 2010. Official methods of analysis of the association of the analytical chemists. Maryland, USA.
- Basso, V., Schiavenin, C., Mendonça, S., de Siqueira, F.G., Salvador, M., Camassola, M., 2020. Chemical features and antioxidant profile by *Schizophyllum commune* produced on different agroindustrial wastes and byproducts of biodiesel production. *Food Chem.* 329. <https://doi.org/10.1016/j.foodchem.2020.127089>.
- Beluhan, S., Ranogajec, A., 2011. Chemical composition and non-volatile components of Croatian wild edible mushrooms. *Food Chem.* 124 (3), 1076–1082.
- Bhattacharyya, A., Chattopadhyay, R., Mitra, S., Crowe, S.E., 2014. Oxidative stress: An essential factor in the pathogenesis of gastrointestinal mucosal diseases. *Physiol. Rev.* 94, 329–354. <https://doi.org/10.1152/physrev.00040.2012>.
- Bisema, J.M., 1968. *Djamur: jang dapat dimakan, jang beracun dan pengusahaan djamur merang*. PT Kinta, Jakarta.
- Boonthatui, Y., Chongsuwat, R., Kittisakulnam, S., 2021. Production of antioxidant bioactive compounds during mycelium growth of *Schizophyllum commune* on different cereal media. *CMUJ. Nat. Sci.* 20 (2), e2021032.
- Carreño-Ruiz, S.D., Lázaro, A.A.A., García, S.C., Hernández, R.G., Chen, J., Navarro, G.K.G., Fajardo, L.V.G., Pérez, N.D.C.J., Cruz, M.T.D.L., Blanco, J.C., Cappello, R.E., 2019. New record of *Schizophyllum* (Schizophyllaceae) from Mexico and the confirmation of its edibility in the humid tropics. *Phytotaxa* 413 (2), 137–148. <https://doi.org/10.11646/phytotaxa.413.2.3>.
- Chandrawanshi, N.G., Tandia, D.K., Jadhav, S.K., 2017. Nutraceutical properties evaluation of *Schizophyllum commune*. *Indian J. Sci. Res.* 13 (2), 57–62.
- Chandrawanshi, N.K., Tandia, D.K., Jadhav, S.K., 2019. Determination of Anti-Diabetic Property of organic and nonorganic solvent extracts of *Schizophyllum commune*. *New Bio. World* 1 (1), 5–8.
- Chang, S., Miles, G.P., 2004. *Mushrooms: Cultivation, Nutritional Value, Medicinal Effects and Environmental Impact*, p. 436. Boca Raton, FL, CRC Press. <https://doi.org/10.1201/9780203492086>.
- Cooke, W.B., 1961. The genus *Schizophyllum*. *Mycologia* 53, 575–599. <https://doi.org/10.2307/3756459>.
- Culmsee, H., Pitopang, R., 2009. Tree diversity in sub-montane and lower montane primary rain forests in Central Sulawesi. *Blume* 54, 119–123. <https://doi.org/10.3767/000651909X475473>.
- Das, S., Bisoyi, S.K., Anmoldeep, P.D., Srivastava, S., 2021. Role of Cultivated Mushrooms in Bioremediation: A Review. *Biol. For.* 13 (1), 160–168.
- Dasanayaka, P.N., Wijeyaratne, S.C., 2017. Cultivation of *Schizophyllum commune* mushroom on different wood substrates. *J. Trop. Fores. Env.* 7 (1), 65–73.
- de Jong, J.F., 2006. *Aerial Hyphae of Schizophyllum commune: Their Function and Formation*. PhD thesis, Univ. Utrecht.
- Dulay, R.M.R., Vicente, J.J.A., Dela, C.A.G., Gagarin, J.M., Fernando, W., Kalaw, S.P., Reyes, R.G., 2016. Antioxidant activity and total phenolic content of *Volvariella*

- volveacea and *Schizophyllum commune* mycelia cultured in indigenous liquid media. *Mycosphere* 7 (2), 131–138. <https://doi.org/10.5943/mycosphere/7/2/4>.
- Elsakhawy, T.A., El-Rahem, W.T.A., 2020. Evaluation of Spent Mushroom Substrate Extract as a Biofertilizer for Growth Improvement of Rice (*Oryza sativa* L.). *Egypt. J. Soil. Sci.* 60 (1), 31–42.
- Espejel-Sánchez, K.I., Espinosa-Solares, T., Reyes-Trejo, B., HernándezRodríguez, G., Cunill-Flores, J.M., Guerra-Ramírez, D., 2021. Nutritional value and thermal degradation of bioactive compounds in wild edible mushrooms. *Rev. Chapingo Ser. Cienc. For. Ambiente.* 27 (3), 337–354. <https://doi.org/10.5154/rchscfa.2020.12.078>.
- Falandysz, J., Borovička, B., 2013. Macro and trace mineral constituents and radionuclides in mushrooms: Health benefits and risks. *Appl. Microbiol. Biotechnol.* 97, 477–501. <https://doi.org/10.1007/s00253-012-4552-8>.
- Filip, R., Shaw, T.A., Nishida, A., Pezacki, J.P., 2019. Fungal Natural Alkaloid Schizocommunin Activates the Aryl Hydrocarbon Receptor Pathway. *Med. Chem. Commun.* 10, 985–990. <https://doi.org/10.1039/C9MD00138G>.
- Fogarasi, M., Socaci, S.A., Dulf, F.V., Diaconasa, Z.M., Fărcaș, A.C., Tofană, M., Semeniuc, C.A., 2018. Bioactive compounds and volatile profiles of five Transylvanian wild edible mushrooms. *Molecules* 23 (12), 3272. <https://doi.org/10.3390/molecules23123272>.
- Fries, E.M., 1815. *Observationes mycologicae*. Gerhardt Bonnier, Copenhagen, Denmark.
- Girma, W., Tasisa, T., 2018. Application of Mushroom as Food and Medicine. *Adv. Biotech Micro* 11, (4). <https://doi.org/10.19080/AIBM.2018.11.555817>.
- González, A., Cruz, M., Losoya, C., Nobre, C., Loreda, A., Rodríguez, R., Contreras, J., Ruth Belmares, R., 2020. Edible mushrooms as a novel protein source for functional foods. *Food Func.* 11 (9), 7400–7414. <https://doi.org/10.1039/D0FO01746A>.
- Gupta, S., Brazier, A.K.M., Lowe, N.M., 2020. Zinc deficiency in low-and middle-income countries: prevalence and approaches for mitigation. *J. Hum. Nutr. Diet.* 33 (5), 624–643. <https://doi.org/10.1111/jhn.12791>.
- Han, C., Zhang, G., Wang, H., Ng, T.B., 2010. Schizolysin, a hemolysin from the split gill mushroom *Schizophyllum commune*. *FEMS Microbiol Lett.* 309 (2), 115–121. <https://doi.org/10.1111/j.1574-6968.2010.02022.x>.
- Hao, L., Xing, X., Li, Z., Zhang, J., Sun, J., Jia, S., Qiao, C., Wu, T., 2010. Optimization of effect factors for mycelia growth and exopolysaccharide production by *Schizophyllum commune*. *Appl. Biochem. Biotechnol.* 160 (2), 621–631. <https://doi.org/10.1007/s12010-008-8507-6>.
- Heleno, S.A., Barros, L., Sousa, M.J., Martins, A., Ferreira, I.C.F.R., 2010. Tocopherols composition of Portuguese wild mushrooms with antioxidant capacity. *Food Chem.* 119 (4), 1443–1450. <https://doi.org/10.1016/j.foodchem.2009.09.025>.
- Herawati, E., Ramadhan, R., Ariyani, F., Marjenah, Kusuma, I.W., Suwinarti, W., Mardji, D., Amirta, R., Arung, E.T., 2021. Phytochemical screening and antioxidant activity of wild mushrooms growing in tropical regions. *Biodivers.* 22 (11), 4716–4721. <https://doi.org/10.13057/biodiv/d221102>.
- Heriyana, E.N., Maryam, L.F., Hadi, Y.S., 2011. *Schizophyllum commune* Fr. Sebagai Jamur Uji Ketahanan Kayu Standar Nasional Indonesia Pada Empat Jenis Kayu Rakyat: Sengon (*P. falcataria*), Karet (*H. brasiliensis*), Tusam (*P. merkusii*), Mangium (*A. mangium*). *J. Silvicultur Tropika.* 2(3), 176–180. doi: <https://doi.org/10.29244/j-siltrop.2.3.325p>.
- Hobs, C.R., 2005. The Chemistry, Nutritional Value, Immunopharmacology, and Safety of the Traditional Food of Medicinal Split-Gill Fungus *Schizophyllum commune* Fr.:Fr (Schizophyllaceae). A Literature Review. *Int. J. Med. Mushrooms.* 7 (1–2), 127–140. <https://doi.org/10.1615/IntJMedMushr.v7.i12.130>.
- Horisawa, S., Ando, H., Ariga, O., Sakuma, Y., 2015. Direct ethanol production from cellulosic materials by consolidated biological processing using the wood rot fungus *Schizophyllum commune*. *Bioresour. Technol.* 197, 37–41. <https://doi.org/10.1016/j.biortech.2015.08.031>.
- İsiloğlu, M., Yılmaz, F., Merdivan, M., 2001. Concentrations of trace elements in wild edible mushrooms. *Food Chem.* 73 (2), 169–175. [https://doi.org/10.1016/S0308-8146\(00\)00257-0](https://doi.org/10.1016/S0308-8146(00)00257-0).
- Jacinto-Azevedo, B., Valdeirama, N., Henríquez, K., Aranda, M., Aqueveque, P., 2021. Nutritional value and biological properties of Chilean wild and commercial edible mushrooms. *Food Chem.* 356. <https://doi.org/10.1016/j.foodchem.2021.129651>.
- James, T.Y., Porter, D., Hamrick, J.L., Vilgalys, R., 1999. Evidence for limited intercontinental gene flow in the cosmopolitan mushroom. *Schizophyllum commune*. *Evolution* 53 (6), 1665–1677. <https://doi.org/10.1111/j.1558-5646.1999.tb04552.x>.
- Kakumu, S., Ishikawa, T., Wakita, T., Yoshioka, K., Ito, Y., Shinagawa, T., 1991. Effect of siozifiran, a polysaccharide, on interferon gamma, antibody production and lymphocyte proliferation specific for hepatitis B virus antigen in patients with chronic hepatitis B. *Int. J. Immunopharmacol.* 13 (7), 969–975. [https://doi.org/10.1016/0192-0561\(91\)90050-h](https://doi.org/10.1016/0192-0561(91)90050-h).
- Kamalebo, H.M., Malale, H.N.S.W., Ndabaga, C.M., Deegreef, J., Kesi, A.D., 2018. Uses and importance of wild fungi: traditional knowledge from the Tshoppe province in the Democratic Republic of the Congo. *J. Ethnobiol. Ethnomed.* 14 (13), 1–12. <https://doi.org/10.1186/s13002-017-0203-6>.
- Kartini, K., Khumaidi, A., Khaerati, K., Ihwan, I., 2018. Ethanol extract of eboni leaf decrease blood glucose level in alloxan induced male rats. *J. Vet.* 19 (3), 329–334.
- Keßler, P.J.A., Bos, M., Sierra Daza, S.E.C., Kop, A., Willemsse, L.P.M., Pitopang, R., Gradstein, S.R., 2002. Checklist of woody plants of Sulawesi, Indonesia. *Blumea Supplement* 14, 1–160.
- Khastini, R.O., Wahyuni, I., Saraswati, I., 2018. Ethnomycology of bracket fungi in Baduy tribe Indonesia. *Biosaintifika* 10 (2), 423–431. <https://doi.org/10.15294/biosaintifika.v10i2.14082>.
- Komatsu, N., Nagumo, N., Okubo, S., Koike, K., 1973. Protective effect of schizophyllan, a polysaccharide of *Schizophyllum commune*, on bacterial infections of mice. *Jpn. J. Antibiot.* 26, 277–283.
- Kozarski, M., Klaus, A., Jakovljevic, D., Todorovic, N., Vunduk, J., Petrović, P., Niksic, M., Vrvic, M.M., van Griensven, L., 2015. Antioxidants of Edible Mushrooms. *Molecules* 20, 19489–19525. <https://doi.org/10.3390/molecules201019489>.
- Krupodorova, T.A., Barshteyn, V.Y., 2015. Amaranth Flour as a New Alternative Substrate for *Schizophyllum commune* Fr.: Fr. and *Cordyceps sinensis* (Berk.) Sacc.. *Growth J. Siberian Fed. Univ. Biol.* 8 (1), 32–44. <https://doi.org/10.17516/1997-1389-2015-8-1-32-44>.
- Krupodorova, T., Sevindik, M., 2020. Antioxidant potential and some mineral contents of wild edible mushroom *Ramaria stricta*. *AgroLife Sci. J.* 9 (1), 186–191.
- Kumla, J., Suwannarath, N., Sujarit, K., Penkhru, W., Kakumyan, P., Jatuwong, K., Vadthananat, S., Lumyong, S., 2020. Cultivation of mushrooms and their lignocellulolytic enzyme production through the utilization of agro-industrial waste. *Molecules* 25 (12), 2811. <https://doi.org/10.3390/molecules25122811>.
- Kuo, M., 2019. *Schizophyllum commune*. Retrieved from mushroomexpert.com. https://www.mushroomexpert.com/schizophyllum_communum.html (accessed in April 2019).
- Kusrinah, Kasiandari, R.S., 2015. Morphological Characteristics and Kinship Relationship of Mushroom *Schizophyllum commune* Fr. *J. Nat. Sci. Res.* 1 (2), 65–71. <https://doi.org/10.21580/jnsr.2015.1.2.1620>.
- Liu, X., Frydenvang, K., Liu, H., Zhai, L., Chen, M., Olsen, C.E., Christensen, S.B., 2015. Iminolactones from *Schizophyllum commune*. *J. Nat. Prod.* 78, 1165–1168. <https://doi.org/10.1021/np500836y>.
- Longvah, T., Deosthale, Y.G., 1998. Compositional and nutritional studies on edible wild mushrooms from northeast India. *Food Chem.* 63 (3), 331–334. [https://doi.org/10.1016/S0308-8146\(98\)00026-0](https://doi.org/10.1016/S0308-8146(98)00026-0).
- Malik, N.A., Kumar, J., Wani, M.S., Tantrar, Y.R., Ahmad, T., 2021. Role of Mushrooms in the Bioremediation of Soil. In: Dar, G.H., Bhat, R.A., Mehmood, M.A., Hakeem, K.R. (Eds.), *Microbiota and Biofertilizers*, Vol 2. Springer, Cham. https://doi.org/10.1007/978-3-030-61010-4_4.
- Mirfat, A.H.S., Noorlidah, A., Vikineswary, S., 2014. Antimicrobial Activities of Split Gill Mushroom *Schizophyllum commune* Fr. *m. J. Res. Commun.* 2 (7), 113–124.
- Mongkantonawat, N., Thumrongchote, D., 2021. Effect of strains and extraction methods on β -glucan production, antioxidant properties, and FTIR Spectra from mushroom fruiting bodies of *Schizophyllum commune* Fr. Thailand. *Food Resc.* 5 (4), 410–415. [https://doi.org/10.26656/fr.2017.5\(4\).259](https://doi.org/10.26656/fr.2017.5(4).259).
- Muszynska, B., Kała, K., Rojowski, J., Grzywacz, A., Opoka, W., 2017. Composition and Biological Properties of *Agaricus bisporus* Fruiting Bodies – a Review. *Pol. J. Food Nutr. Sci.* 67 (3), 173–181. <https://doi.org/10.1515/pjfn-2016-0032>.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., Da FONSECA, G.A., Kent, J., 2000. Biodiversity hotspots for conservation priorities. *Nature* 403, 853–858. <https://doi.org/10.1038/35002501>.
- Niederpruem, D.J., Hunt, S., 1964. Polyols in *Schizophyllum commune*. *Am. J. Bot.* 54 (2), 241–245. <https://doi.org/10.2307/2440804>.
- Nion, Y.S., Djaya, A.A., Kadie, E.M., Lune, S., Wijaya, C.H., 2012. Siklus Hidup Jamur Konsumsi Lokal Kulat Kritis (*Schizophyllum commune*) Pada Daerah Bergambut dan Daerah Bertanah Mineral serta Potensi Nutrisinya (Cycles life of local edible mushroom Kulat Kritis (*Schizophyllum commune*) on peat soil and mineral soil also its nutrition potency). *J. Biologi Indones.* 8 (20), 399–406. <https://doi.org/10.14203/jbi.v8i2.3060>.
- Niu, Y., Yun, J., Bi, Y., Wang, T., Zhang, Y., Liu, H., Zhao, F., 2020. Predicting the shelf life of postharvest *Flammulina velutipes* at various temperatures based on mushroom quality and specific spoilage organisms. *Postharvest Biol. Technol.* 167. <https://doi.org/10.1016/j.postharvbio.2020.111235>.
- Nurhasanawati, H., Sundu, R., Supriningrum, R., Kuspradini, H., Arung, E.T., 2019. Antioxidant activity, total phenolic and flavonoid content of several indigenous species of ferns in East Kalimantan, Indonesia. *Biodivers.* 20 (2), 576–580. <https://doi.org/10.13057/biodiv/d200238>.
- Nurliana, A.I., Putra, I.P., Ikhsan, M., 2021. Catatan Pemanfaatan *Schizophyllum commune* di Kampung Udapi Hilir. *Papua Barat. Integrated Lab. J.* 9 (1), 18–28. <https://doi.org/10.14421/ilj.2021.090104>.
- Okwulehie, I.C., Nwosu, C.P., Okoroafor, C.J., 2007. Pharmaceutical and Nutritional Prospects of Two Wild Macro-Fungi Found in Nigeria. *Biotechnol* 6 (4), 567–572. <https://doi.org/10.3923/biotech.2007.567.572>.
- Ooi, V.E.C., Liu, F., 2000. Immunomodulation and Anti-Cancer Activity of Polysaccharide-Protein Complexes. *Curr. Med. Chem.* 7 (7), 715–729. <https://doi.org/10.2174/0929867003374705>.
- Ouzouni, P.K., Petridis, D., Koller, W.D., Riganakos, K.A., 2009. Nutritional value and metal content of wild edible mushrooms collected from West Macedonia and Epirus, Greece. *Food Chem.* 115 (4), 1575–1580. <https://doi.org/10.1016/j.foodchem.2009.02.014>.
- Owaid, M.N., Abed, I.A., Al-Saeedi, S.S.S., 2017. Applicable properties of the bio-fertilizer spent mushroom substrate in organic systems as a byproduct from the cultivation of *Pleurotus* spp. *Inform. Proc. in Agric.* 4 (1), 78–82. <https://doi.org/10.1016/j.inpa.2017.01.001>.
- Pandey, A., Tripathi, S., 2014. Concept of standardization, extraction and pre phytochemical screening strategies for herbal drug. *J. Pharmacogn. Phytochem.* 2 (5), 115–119.
- Patria, A., Putra, P.S., 2020. Development of the Palu-Koro Fault in NW Palu Valley, Indonesia. *Geosci. Lett.* 7. <https://doi.org/10.1186/s40562-020-0150-2>.

- Plants of the World Online (POWO), 2019, October 30. Royal Botanical Garden (<https://powo.science.kew.org/>).
- Pooja, S., Vidyasagar, G., 2016. Phytochemical screening for secondary metabolites of *Opuntia dillenii* Haw. J. Med. Plants. Stud. 4 (5), 39–43.
- Preecha, C., Thongliumnak, S., 2015. Bag opening technique for bag spawn culture of split gill mushroom (*Schizophyllum commune*). Int. J. Agric. Technol. 11 (2), 367–372.
- Raaman, N., 2006. Phytochemical Techniques. New India Publishing Agensi, India, p. 306.
- Raper, J.B., Miles, P.G., 1958. The Genetics of *Schizophyllum Commune*. 43(3), 530–546. doi: 10.1093/genetics/43.3.530.
- Rózsa, M., Măniuiu, D.N., Egyed, E., 2021. Influence of magnesium (Mg) source on the *Cordyceps militaris* (L.) mushroom mycelium growth. Curr. Trends Nat. Sci 10, 333–340.
- Sahira, B.K., Cathrine, L., 2015. General Techniques Involved in Phytochemical Analysis. Int. J. Adv. Res. Chem. Sci. 2 (4), 25–32.
- Sánchez, C., 2017. Reactive oxygen species and antioxidant properties from mushrooms. Synth. Syst. Biotechnol. 2 (1), 13–22. <https://doi.org/10.1016/j.synbio.2016.12.001>.
- Sande, D., de Oliveira, G.P., e Moura, M.A.F., Martins, B.A., Lima, M.T.N.S., Takahashi, J.A., 2019. Edible mushrooms as a ubiquitous source of essential fatty acids. Food Res. Int. 125. <https://doi.org/10.1016/j.foodres.2019.108524>.
- Sanmee, R., Dell, B., Lumyong, P., Izumori, K., Lumyong, S., 2003. Nutritive value of popular wild edible mushrooms from northern Thailand. Food Chem. 82 (4), 527–532. [https://doi.org/10.1016/S0308-8146\(02\)00595-2](https://doi.org/10.1016/S0308-8146(02)00595-2).
- Shaikh, J.R., Patil, M., 2020. Qualitative tests for preliminary phytochemical screening: An overview. Int. J. Chem. Stud. 8 (2), 603–608. <https://doi.org/10.22271/chemi.2020.v8.i2i.8834>.
- Singh, R., 2017. A Review on Different Benefits of Mushroom. IOSR J. Pharm. Biol. Sci. 12 (1), 107–111. <https://doi.org/10.9790/3008-120102107111>.
- Socquet, A., Simons, W., Vigny, C., McCafrey, R., Subarya, C., Sarsito, D., Ambrosius, B., Spakman, W., 2006. Microblock rotations and fault coupling in SE Asia triple junction (Sulawesi, Indonesia) from GPS and earthquake slip vector data. J. Geophys. Res. Solid Earth 111, B08409. <https://doi.org/10.1029/2005JB003963>.
- Stan, D., Enciu, A.M., Mateescu, A.L., Ion, A.C., Brezeanu, A.C., Stan, D., Tanase, C., 2021. Natural compounds with antimicrobial and antiviral effect and nanocarriers used for their transportation. Front. Pharmacol. 12. <https://doi.org/10.3389/fphar.2021.723233>.
- Sun, L., Niu, Z., 2020. A mushroom diet reduced the risk of pregnancy-induced hypertension and macrosomia: a randomized clinical trial. Food Nutr. Res. 64, 4451. <https://doi.org/10.29219/fnr.v64.4451>.
- Suwanarach, N., Kumla, J., Zhao, Y., Kakumyan, P., 2022. Impact of Cultivation Substrate and Microbial Community on Improving Mushroom Productivity: A Review. Biology 11, 569. <https://doi.org/10.3390/biology11040569>.
- Suzuki, D., 2020. Effects of Phosphoryl Oligosaccharides of Calcium (POs-Ca) on Mycelial Growth and Fruiting Body Development of the Edible Mushroom, *Pleurotus ostreatus*. J. Appl. Glycosci. 67 (3), 67–72. 10.5458%2Fjag.jag.JAG-2020_0001.
- Tabata, K., Ito, W., Kojima, T., Kawabata, S., Misaki, A., 1981. Ultrasonic degradation of schizophyllan, an antitumor polysaccharide produced by *Schizophyllum commune*. Fries. Carbohydr Res 89 (1), 121–135. [https://doi.org/10.1016/S0008-6215\(00\)85234-9](https://doi.org/10.1016/S0008-6215(00)85234-9).
- Takemoto, S., Nakamura, H., Erwin, Imamura, Y., Shimane, T., 2010. *Schizophyllum commune* as a Ubiquitous plant parasite. JARQ 44(4), 357–364. <https://www.jircas.affrc.go.jp>.
- Teoh, Y.P., Don, M.M., 2013. In vitro anti fungal activities and phytochemical analysis of filamentous white-rot fungi, *Schizophyllum commune*. Sains Malays. 42 (9), 1267–1272.
- Teoh, Y.P., Don, M.M., Ujang, S., 2017. Production of Biomass by *Schizophyllum commune* and Its Antifungal Activity towards rubberwood-degrading fungi. Sains Malays. 46 (1), 123–128. <https://doi.org/10.17576/jsm-2017-4601-16>.
- Uzun, Y., Gencelep, H., Kaya, A., Akcay, M.E., 2011. The Mineral Contents of Some Wild Edible Mushrooms. Ekoloji Dergisi. 20 (80), 6–12.
- Vannucci, L., Krizan, J., Sima, P., Stakheev, D., Caja, F., Rajsiglova, L., Horak, V., Saiegh, M., 2013. Immunostimulatory properties and antitumor activities of glucans. Int. J. Oncol. 43 (2), 357–364. <https://doi.org/10.3892/ijo.2013.1974>.
- Waktola, G., Temesgen, T., 2018. Application of mushrooms as food and medicine. Adv. Biotech. Microbiol. 11. <https://doi.org/10.19080/AIBM.2018.11.555817>.
- Whitten, A.J., Muslim, M., Henderson, G.S., 1987. The Ecology of Sulawesi. Gadjah Mada University Press, Yogyakarta, Indonesia.
- Wirth, S., Krause, K., Kunert, M., Broska, S., Paetz, C., Boland, W., Kothe, E., 2021. Function of sesquiterpenes from *Schizophyllum commune* in interspecific interactions. PLoS ONE 16 (1), 1–14. <https://doi.org/10.1371/journal.pone.0245623>.
- Woldemariam, W.G., 2019. Mushrooms in the Bio-Remediation of Wastes from Soil. Adv. Life Sci. Tech. 76, 41–46. <https://doi.org/10.7176/ALST>.
- Yamamoto, T., Yamashita, T., Tsubura, E., 1981. Inhibition of pulmonary metastasis of Lewis lung carcinoma by a glucan, schizophyllan. Inv. Metast. 1, 71–84.
- Yildiz, S., Yilmaz, A., Can, Z., Kiliç, C., Yıldiz, Ü.C., 2017. Total phenolic, flavonoid, tannin contents and antioxidant properties of *Pleurotus ostreatus* and *Pleurotus citrinopileatus* cultivated on various sawdust. GIDA 42, 315–323. <https://doi.org/10.15237/gida.GD16099>.
- Yim, H.S., Chye, F.Y., Rao, V., Low, J.Y., Matanjun, P., How, S.E.C.W., 2013. Optimization of Extraction Time and Temperature on Antioxidant Activity of *Schizophyllum commune* Aqueous Extract Using Response Surface Methodology. J. Food Sci. Technol. 50, 275–283. <https://doi.org/10.1007/s13197-011-0349-5>.
- Yusran, Y., Erniwati, E., Wahyuni, D., Ramadhani, R., Khumaidi, A., 2021. Diversity of macro fungus across three altitudinal ranges in Lore Lindu National Park, Central Sulawesi, Indonesia and their utilization by local residents. Biodiversity 22, 199–210. <https://doi.org/10.13057/biodiv/d220126>.
- Yusran, Y., Erniwati, E., Khumaidi, A., Pitopang, R., Rahmawati, R., Korja, I.N., Labiro, E., Rukmi, R., Muthmainnah, M., Maksum, H., Setyawati, R., 2022a. Diversity of Macro Fungus Species in Several Types of Land Use Around Lore Lindu National Park, Central Sulawesi, Indonesia. Int. J. Des. Nat. Ecodyn. 17, 341–351. <https://doi.org/10.18280/ijnd.170303>.
- Yusran, Y., Erniwati, E., Khumaidi, A., Setiarto, R.H.B., 2022b. Effect of Cooking on the Proximate Composition and Minerals Content of Wild Edible Macro Fungi from Lore Lindu National Park, Central Sulawesi, Indonesia. African. J. Food Agric. Nutr. Dev. 22, 20523–20541. <https://doi.org/10.18697/ajfand.11021660>.
- Zeb, M., Lee, C.H., 2021. Medicinal Properties and Bioactive Compounds from Wild Mushrooms Native to North America. Molecules 26, 251. <https://doi.org/10.3390/molecules26020251>.
- Zohra, S.F., Meriem, B., Samira, S., Alsayadi Muneer, M.S., 2012. Phytochemical Screening and identification of some compounds from Mallow. J. Nat. Prod. Plant Resour. 2, 512–516.

Diversity of substrate type, ethnomycology, mineral composition, proximate, and phytochemical compounds of the *Schizopyllum* commune Fr. in the area along Palu-Koro Fault, Central Sulawesi, Indonesia

ORIGINALITY REPORT

7%

SIMILARITY INDEX

5%

INTERNET SOURCES

6%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1	rcastoragev2.blob.core.windows.net Internet Source	2%
2	Submitted to Higher Education Commission Pakistan Student Paper	2%
3	docplayer.net Internet Source	2%
4	Submitted to Jeju National University Student Paper	1%
5	Yuan-yuan Liu, Wei-hong Sun, Bing-zheng Li, Nan Shang, Yong Wang, Wei-qiao Lv, Dong Li, Li-jun Wang. "Value-added application of <i>Platycodon grandiflorus</i> (Jacq.) A.DC. roots (PGR) by ultrasound-assisted extraction (UAE) process to improve physicochemical quality, structural characteristics and functional properties", Food Chemistry, 2021 Publication	1%



Exclude quotes On

Exclude matches < 1%

Exclude bibliography On