

Info Notifikasi Awal



Participants

Journal Editor (journaleditor)

Section Editor B. P. Tomasouw (editor2)

Financial Editor D. Patty (editor7)

Gregoria Ariyanti (gregoria)

Messages

Note	From
<p>Yth.: Bpk/Ibu Penulis</p> <p>Salam,</p> <p>Bersama ini kami sampaikan bahwa artikel bpk/Ibu, sudah masuk di redaksi jurnal barekeng. Selanjutnya artikel bpk/ibu akan kami proses sesuai dengan tahapan dan aturan yang ditetapkan oleh Redaksi BAREKENG: Jurnal Ilmu Matematika dan Terapan. Harapan kami, bpk/ibu penulis dapat bersabar menunggu proses dari tim redaksi/editor sambil mengecek perkembangan via web OJS Jurnal Barekeng, email dan WA (085243358669).</p> <p>Perkembangan artikel dan Informasi selanjutnya akan kami kabari via web ojs maupun via WhatsApp (WA), mohon dapat menyertakan nomor WA yang bisa dihubungi untuk mempermudah komunikasi.</p>	<p>editor2</p> <p>Jul 13</p>

redaksi/editor sambil mengecek perkembangan via web OJS Jurnal Barekeng, email dan WA (085243358669).

Perkembangan artikel dan Informasi selanjutnya akan kami kabari via web ojs maupun via WhatsApp (WA), mohon dapat menyertakan nomor WA yang bisa dihubungi untuk mempermudah komunikasi.

Terima kasih atas kepercayaan bpk/ibu, untuk mempublikasikan artikelnnya pada Jurnal BAREKENG.

Hormat kami,

Yopi Andry Lesnussa,

Editor in Chief BAREKENG: Jurnal Ilmu Matematika dan Terapan

Email.: barekeng.math@yahoo.com

WhatsApp (WA): +62 85243358669

<p>▶ Terima kasih atas informasi yang Bapak berikan.</p> <p>Berikut nomor WhatsApp (WA) : +62818260906</p> <p>Salam,</p> <p>Gregoria Ariyanti</p>	<p>gregoria</p> <p>Jul 13</p>
---	-------------------------------

Add Message

Info Perbaikan Pertama



Participants

Journal Editor (journaleditor)

Section Editor B. P. Tomasouw (editor2)

Financial Editor D. Patty (editor7)

Gregoria Ariyanti (gregoria)

Messages

Note	From
<p>Yth.: Bpk/Ibu Penulis</p> <p>Salam,</p> <p>Bersama ini kami sampaikan bahwa artikel bpk/Ibu, sudah diperiksa dan dikomentari oleh tim editor Jurnal. Harapan kami, kiranya bpk/ibu dapat memperbaiki artikel tersebut sesuai dengan hasil pemeriksaan dan komentar yang diberikan serta mengirimkannya kembali melalui Menu Add Discussion dengan Subject : "Hasil Perbaikan Pertama"</p> <p>Perkembangan artikel dan Informasi selanjutnya akan kami kabari via web ojs maupun via WhatsApp (WA), mohon dapat menyertakan nomor WA yang bisa dihubungi untuk mempermudah komunikasi.</p>	<p>editor2 Jul 31</p>

Jul 31

Salam,

Bersama ini kami sampaikan bahwa artikel bpk/Ibu, sudah diperiksa dan dikomentari oleh tim editor Jurnal. Harapan kami, kiranya bpk/ibu dapat memperbaiki artikel tersebut sesuai dengan hasil pemeriksaan dan komentar yang diberikan serta mengirimkannya kembali melalui Menu Add Discussion dengan Subject : "Hasil Perbaikan Pertama"

Perkembangan artikel dan Informasi selanjutnya akan kami kabari via web ojs maupun via WhatsApp (WA), mohon dapat menyertakan nomor WA yang bisa dihubungi untuk mempermudah komunikasi.

Terima kasih atas kepercayaan bpk/ibu, untuk mempublikasikan artikelnya pada Jurnal BAREKENG.

Hormat kami,

Yopi Andry Lesnussa,
Editor in Chief BAREKENG: Jurnal Ilmu Matematika dan Terapan

Email: barekeng.math@yahoo.com
WhatsApp (WA): +62 85243358669

[editor2, 6344-Article Text-30186-1-2-20220710 revisi 1.docx](#)

[editor2, English Template of Barekeng 2022 OK.docx](#)

Add Message

Notifications



[BAREKENG: J. Math. & App.] Editor Decision

2022-11-10 12:07 PM

Gregoria Ariyanti:

We have reached a decision regarding your submission to BAREKENG: Jurnal Ilmu Matematika dan Terapan, "A NOTE ON THE SOLUTION OF THE CHARACTERISTIC EQUATION OVER THE SYMMETRIZED MAX-PLUS ALGEBRA".

Our decision is to: Accept Submission

Journal Editor
Jurusan Matematika FMIPA Universitas Pattimura
barekeng.math@fmipa.unpatti.ac.id

[BAREKENG: Jurnal Ilmu Matematika dan Terapan](#)

Notifications



[BAREKENG: J. Math. & App.] Editor Decision

2022-11-18 09:29 PM

Gregoria Ariyanti:

The editing of your submission, "A NOTE ON THE SOLUTION OF THE CHARACTERISTIC EQUATION OVER THE SYMMETRIZED MAX-PLUS ALGEBRA," is complete. We are now sending it to production.

Submission URL: <https://ojs3.unpatti.ac.id/index.php/barekeng/authorDashboard/submission/6344>

Journal Editor
Jurusan Matematika FMIPA Universitas Pattimura
barekeng.math@fmipa.unpatti.ac.id

[BAREKENG: Jurnal Ilmu Matematika dan Terapan](#)

Participants

Journal Editor (journaleditor)

Proofreader M. I. Tilukay (editor3)

Gregoria Ariyanti (gregoria)

Messages

Note

From

Dear Author(s),

journaleditor
Dec 04

Here we send the Galley proof file of your article, so you can have a final check of your article. Please check it carefully and give the response for us. If there is anything you want to revise, please send you message to reply this message (in OJS system) or send in email or our WhatsApp number (085243358669).

Thank you,

best regards,
Editor (Proofreader)

 [journaleditor, 1347-1354 6344-Article Text-30973-1-18-20220802 \(pr\).pdf](#)

▶ Dear Editor,

gregoria
Dec 05

There is not anything I want to revise. Thank you.

A NOTE ON THE SOLUTION OF THE CHARACTERISTIC EQUATION OVER THE SYMMETRIZED MAX-PLUS ALGEBRA

Gregoria Ariyanti^{1*}

¹Department of Mathematics Education, Faculty of Teacher Training and Education,
Widya Mandala Surabaya Catholic University, Madiun City Campus
Madiun, 63131, East Java, Indonesia

Corresponding author e-mail: ^{1*} gregoria.ariyanti@ukwms.ac.id

Abstract. The symmetrized max-plus algebra is an extension of max-plus algebra. One of the problems in the symmetrized max-plus algebra is determining the eigenvalues of a matrix. If the determinant can be defined, the characteristic equation can be formulated as a max-plus algebraic multivariate polynomial equation system. A mathematical tool for solving the problem using operations as in conventional algebra, known as the extended linear complementary problem (ELCP), to determine the solution to the characteristic equation. In this paper, we will describe the use of ELCP in determining the solution to the characteristic equations of matrices over the symmetrized max-plus algebra.

Keywords: symmetrized max-plus algebra, determinant, characteristic equation, elcp

Article info:

Submitted: date, month, year

Accepted: date, month, year

How to cite this article:

First author, second author, etc, "TITLE OF ARTICLE", BAREKENG: J. Il. Mat. & Ter., vol. xx, iss. xx, pp. xxx-xxx, Month, Year.



This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).
Copyright © 2021 First author, Second author, etc.

1. INTRODUCTION

It is known that for \mathbb{R} , the set of all real numbers, given $\mathbb{R}_\epsilon = \mathbb{R} \cup \{\epsilon\}$ with $\epsilon := -\infty$ and $e := 0$ and for every $a, b \in \mathbb{R}_\epsilon$, defined operation \oplus and \otimes as $a \oplus b := \max(a, b)$ and $a \otimes b := a + b$, next one, $(\mathbb{R}_\epsilon, \oplus, \otimes)$ is called max-plus algebra ([1],[2],[3],[4]). Max-plus algebra is an example of a structure semiring that has no inverse of \oplus . In other words, if $a \in \mathbb{R}_\epsilon$ then nothing $b \in \mathbb{R}_\epsilon$ so $a \oplus b = b \oplus a = \epsilon$, unless $a = \epsilon$. In this case, an extension of max-plus algebra has been developed, which is called the symmetrized max-plus algebra and is denoted $(\mathbb{S}, \oplus, \otimes)$ with $\mathbb{S} = (\mathbb{R}_\epsilon^2)/\mathcal{B}$ where \mathcal{B} is an equivalence relation ([3],[5],[6]).

Because the matrix over the symmetrized max-plus algebra can be defined as a determinant ([7],[8]). The equation characteristics can be formulated as a system of equations and polynomial inequality multivariate algebra max. To determine solving systems of equations and polynomial inequalities multivariate max algebra can use operations like in conventional algebra through known mathematical tools with the extended complementary linear problem or ELCP ([9],[10],[11]).

This paper aims to determine the condition of a characteristic equation that has a solution viewed from the matrix over the symmetrized max-plus algebra. In this paper, we will mainly concern with a characteristic equation matrix over the symmetrized max-plus algebra. We show that the solution of a characteristic equation on $\mathbb{S}^\oplus \cup \mathbb{S}^\ominus \cup \mathbb{S}^*$ can be determined with ELCP. Section 2 will review some basic facts for the symmetrized max-plus algebra. The solution of the characteristic equation of the matrix over \mathbb{S} that determine with ELCP is given in Section 3.

2. MATERIALS AND METHODS

2.1 The Symmetrized Max-Plus Algebra

In this section, we review some basic facts for max-plus algebra and the symmetrized max-plus algebra. The characteristic of max-plus algebra can be seen in [1], [5], [12]. Let \mathbb{R} denote the set of all real numbers and $\mathbb{R}_\epsilon = \mathbb{R} \cup \{\epsilon\}$ with $\epsilon := -\infty$ as the null element and $e := 0$ as the unit element. For all $a, b \in \mathbb{R}_\epsilon$, we have $a \oplus b = \max(a, b)$ and $a \otimes b = a + b$. The structure $(\mathbb{R}_\epsilon, \oplus, \otimes)$ is called the max-plus algebra.

Definition 1. [1], [5] Let $u = (x, y), v = (w, z) \in \mathbb{R}_\epsilon^2$. We have

1. Operators \ominus and $(\cdot)^*$ are defined as $\ominus u = (y, x)$ and $u^* = u \oplus u$.
2. An element u is called balances with v , denoted by $u \nabla v$, if $x \oplus z = y \oplus w$.
3. A relation \mathcal{B} is defined as follows : $(x, y) \mathcal{B} (w, z)$ if $\begin{cases} (x, y) \nabla (w, z) & \text{if } x \neq y \text{ and } w \neq z \\ (x, y) = (w, z), & \text{otherwise} \end{cases}$.

Because \mathcal{B} is an equivalence relation, we have the set of factor $\mathbb{S} = (\mathbb{R}_\epsilon^2)/\mathcal{B}$ and the system $(\mathbb{S}, \oplus, \otimes)$ is called the symmetrized max-plus algebra, with the operations of addition and multiplication on \mathbb{S} are defined as follows:

$$\overline{(a, b)} \oplus \overline{(c, d)} = \overline{(a \oplus c, b \oplus d)} \quad \text{and} \\ \overline{(a, b)} \otimes \overline{(c, d)} = \overline{(a \otimes c \oplus b \otimes d, a \otimes d \oplus b \otimes c)} \quad \text{for } \overline{(a, b)}, \overline{(c, d)} \in \mathbb{S}.$$

The system $(\mathbb{S}, \oplus, \otimes)$ is a semiring, because (\mathbb{S}, \oplus) is associative, (\mathbb{S}, \otimes) is associative, and $(\mathbb{S}, \oplus, \otimes)$ satisfies both the left and right distributive [7]. Because \mathbb{S}^\oplus isomorphic with \mathbb{R}_ϵ , so it will be shown that for $a \in \mathbb{R}_\epsilon$, can be expressed by $\overline{(a, \epsilon)} \in \mathbb{S}^\oplus$. Furthermore, it is easily seen that for $a \in \mathbb{R}_\epsilon$ we have :

1. $a = \overline{(a, \epsilon)}$ with $\overline{(a, \epsilon)} \in \mathbb{S}^\oplus$.
2. $\ominus a = \ominus \overline{(a, \epsilon)} = \overline{(a, \epsilon)} = \overline{(\epsilon, a)}$ with $\overline{(\epsilon, a)} \in \mathbb{S}^\ominus$.
 $a^* = a \oplus a = \overline{(a, \epsilon)} \oplus \overline{(a, \epsilon)} = \overline{(a, a)} \in \mathbb{S}^*$.

Lemma 2. [1] For $a, b \in \mathbb{R}_\epsilon, a \oplus b = \overline{(a, b)}$.

Proof:

$$a \oplus b = \overline{(a, \epsilon)} \oplus \overline{(b, \epsilon)} = \overline{(a, \epsilon)} \oplus \overline{(\epsilon, b)} = \overline{(a, b)}. \quad \square$$

Commented [bC1]: S tr t w @ z p f s @ j g w j s a @ r i f p @ t g m i p j y p @ f s z f g m f v e x @ u g z f y @ j c f q @ r j s z @ j k j w s a

Commented [bC2]: S tr t w @ z p f s @ j g w j s a @ r i f p @ t g m i p j y p @ f s z f g m f v e x @ u g z f y @ j c f q @ r j s z @ j k j w s a

Commented [bC3]: 1.Diganti menjadi **RESEARCH METHODS**

Commented [bC4]: S tr t w @ z p f s @ j g w j s a @ r i f p @ t g m i p j y p @ f s z f g m f v e x @ u g z f y @ j c f q @ r j s z @ j k j w s a

Commented [bC5]: S tr t w @ z p f s @ j g w j s a @ r i f p @ t g m i p j y p @ f s z f g m f v e x @ u g z f y @ j c f q @ r j s z @ j k j w s a

Commented [bC6]: Hilangkan bukti

Let \mathbb{S} be the symmetrized max-plus algebra, a positive integer n , and $M_n(\mathbb{S})$ be the set of all $n \times n$ matrices over \mathbb{S} . Operations \oplus and \otimes for matrix over the symmetrized max-plus algebra are defined as follows: $C = A \oplus B \Rightarrow c_{ij} = a_{ij} \oplus b_{ij}$ and $C = A \otimes B \Rightarrow c_{ij} = \bigoplus_l a_{il} \otimes b_{lj}$.

The $n \times n$ zero matrix over \mathbb{S} is ε_n with $(\varepsilon_n)_{ij} = \varepsilon$ and a $n \times n$ identity matrix over \mathbb{S} is E_n with

$$(E_n)_{ij} = \begin{cases} e, & \text{if } i = j \\ \varepsilon, & \text{if } i \neq j \end{cases}$$

Definition 3. We say that the matrix $A \in M_n(\mathbb{S})$ is invertible over \mathbb{S} if $A \otimes B \nabla E_n$ and $B \otimes A \nabla E_n$ for any $B \in M_n(\mathbb{S})$.

2.2 The Extended Complementary Linear Problem

Some optimization problems that are mathematically modeled can appear as a multiplication between variables or in the form of a square of a variable. Problems with these complementary conditions cannot be solved using the linear complementary problem (LCP). Therefore, the concept of the extended complementary linear problem (ELCP) emerged. The form of the ELCP problem is as follows.

Let a matrix $A \in M_{p \times n}(\mathbb{R})$, $B \in M_{q \times n}(\mathbb{R})$, $\bar{c} \in M_{p \times 1}(\mathbb{R})$, and $\bar{d} \in M_{q \times 1}(\mathbb{R})$. Also given the set $\phi_j \subset \{1, 2, \dots, p\}$ where $j = 1, 2, \dots, m$. Determine the vector $\bar{x} \in M_{n \times 1}(\mathbb{R})$ such that it satisfies:

$$\sum_j^m \prod_{i \in \phi_j} (A\bar{x} - \bar{c})_i = 0 \quad (1)$$

with

$$A\bar{x} \geq \bar{c} \text{ and } B\bar{x} = \bar{d} \quad (2)$$

The (1) equation is called the complementary condition. The vector set \bar{x} that satisfies (2) will be a solution to that problem. The ELCP form with constrains (2) can be solved by changing the ELCP form (1) and (2) into homogeneous ELCP structures.

The general form of homogeneous ELCP is as follows.

Given matrix $P \in M_{p \times n}(\mathbb{R})$ and $Q \in M_{q \times n}(\mathbb{R})$, and the set $\phi_j \in \{1, 2, \dots, p\}$ with $j = 1, 2, \dots, m$.

Determine the nontrivial vector $\bar{u} \in M_{n \times 1}(\mathbb{R})$ such that it satisfies:

$$\sum_{j=1}^m \prod_{i \in \phi_j} (P\bar{u})_i = 0 \quad (3)$$

with

$$P\bar{u} \geq 0 \text{ and } Q\bar{u} = 0 \quad (4)$$

The ELCP forms in (1) and (2) can be brought to homogeneous ELCP forms (3) and (4) by taking $\bar{u} = \begin{bmatrix} \bar{x} \\ \alpha \end{bmatrix}$.

$P = \begin{bmatrix} A & -\bar{c} \\ 0_{1 \times n} & 1 \end{bmatrix}$, and $Q = [B \quad -\bar{d}]$ with $\alpha \geq 0$. All \bar{u} vectors that satisfy (4) will form the solution set of the above problem.

2.3 Research Methods

The method used in preparing this research is first to conduct a literature study on symmetrized max-plus algebra, the matrix over symmetrized max-plus algebra, and the general form of ELCP problems. This study is necessary because the matrix over symmetrized max-plus algebra can be defined, so from the formulation of the eigenvalues of the matrix over symmetrized max-plus algebra, characteristic equations are formed. And the characteristic equation can be formulated as a system of equations and polynomial inequalities multivariate algebra max-plus.

Determine the solution to a system of equations and inequalities, polynomial multivariate max algebra can use operations as in conventional algebra through a mathematical tool known as the Extended Linear Complementarity Problem (ELCP).




Figure 1. Workflow for the conceptual study of characterization equation of the symmetrized max-plus algebra

3. RESULTS AND DISCUSSION

For A, b over the symmetrized max-plus algebra, $A \otimes x \nabla b$ is called the linear balanced system, and $A \otimes x \nabla \epsilon$ is the homogeneous linear balanced system. Note that the operator " ∇ " and the homogeneous linear balances have the following characterizations.

Theorem 4. [12],[13] *Given \mathbb{S} is a symmetrized max-plus algebra. The following properties apply :*

1. For all $a, b, c \in \mathbb{C}, a \ominus c \nabla b$ if and only if $a \nabla b \oplus c$.
2. For all $a, b \in \mathbb{S}^{\oplus} \cup \mathbb{S}^{\ominus}, a \nabla b \Rightarrow a = b$.
3. Let $A \in M_n(\mathbb{S})$. The homogeneous linear balance systems $A \otimes x \nabla \epsilon_{n \times 1}$ has a nontrivial solution in \mathbb{S}^{\oplus} or \mathbb{S}^{\ominus} if and only if $\det(A) \nabla \epsilon_{n \times 1}$.

The following is the relationship between Systems of Equations and Inequality Multivariate Polynomials in Max-Plus with ELCP. The general form of a multivariate polynomial system of equations and inequalities in max-plus algebra is as follows:

Given $p_1 + p_2$, with $p_1, p_2 \in \mathbb{N}$, integer $m_1, m_2, \dots, m_{p_1+p_2} \in \mathbb{N} \cup \{0\}$ and $a_{ki}, b_k, c_{kij} \in \mathbb{R}$, for $k = 1, 2, \dots, p_1 + p_2, i = 1, 2, \dots, m_k$, and $j = 1, 2, \dots, n$ and will look for the vector $x \in \mathbb{R}^n$, so that it satisfies:

$$\bigoplus_{i=1}^{m_k} a_{ki} \otimes \bigotimes_{j=1}^n x_j^{\otimes c_{kij}} = b_k, \text{ for } k = 1, 2, \dots, p_1 \quad (5)$$

$$\bigoplus_{i=1}^{m_k} a_{ki} \otimes \bigotimes_{j=1}^n x_j^{\otimes c_{kij}} \leq b_k, \text{ for } k = p_1 + 1, p_1 + 2, \dots, p_1 + p_2 \quad (6)$$

Based on De Schutter and De Moor's statement, there is the following theorem about necessary and sufficient conditions for the extended complementary linear problem ([14],[15]). A multivariate polynomial system of equations and inequalities in max-plus algebra if and only if it is an extended complementary linear problem (ELCP)

From (5) and (6) can be formed matrix $A = \begin{bmatrix} -C_1 \\ -C_2 \\ \vdots \\ -C_{p_1+p_2} \end{bmatrix}$ and $c = \begin{bmatrix} -d_1 \\ -d_2 \\ \vdots \\ -d_{p_1+p_2} \end{bmatrix}$.

If m_k in the equation (5) is equal to 1, then we get: $a_{k1} + \sum_{j=1}^n c_{k1j} x_j = b_k$, for $k = 1, 2, \dots, p_1$. If we add $-a_{k1}$ in two side and define $d_{k1} = b_k - a_{k1}$, we get $\sum_{j=1}^n c_{k1j} x_j = d_{k1}$. So that we get the matrix

$$B = \begin{bmatrix} c_{111} & c_{112} & \dots & c_{11n} \\ c_{211} & c_{212} & \dots & c_{21n} \\ \vdots & \vdots & \ddots & \vdots \\ c_{p_111} & c_{p_112} & \dots & c_{p_11n} \end{bmatrix}, \bar{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \text{ and } \bar{d} = \begin{bmatrix} d_{11} \\ d_{21} \\ \vdots \\ d_{p_11} \end{bmatrix}.$$

Commented [bC7]: S tr t w e z o r p f e w j k j w s a c h y l f p g t g w
l i p j y p k f s z f g m f v e x a n g z f y k j d f q z n r j e z h i k j w s a c

In other words, the equation is obtained in the form of a matrix $B\bar{x} = \bar{d}$.

Defined as much as p_1 set $\phi_1, \phi_2, \dots, \phi_{p_1}$ such that $\phi_j = \{s_j + 1, s_j + 2, \dots, s_j + m_j\}$ with $s_1 = 0$ for $j = 1, 2, \dots, p_1$ and $s_{j+1} = s_j + m_j$ for $j = 1, 2, \dots, p_1 - 1$. Thus obtained:

$$\begin{aligned} \phi_1 &= \{s_1 + 1, s_1 + 2, \dots, s_1 + m_1\} = \{1, 2, \dots, m_1\} \\ \phi_2 &= \{s_2 + 1, s_2 + 2, \dots, s_2 + m_2\} = \{s_1 + m_1 + 1, \\ &\quad s_1 + m_1 + 2, \dots, s_1 + m_1 + m_2\} \\ &= \{m_1 + 1, m_1 + 2, \dots, m_1 + m_2\} \\ &\quad \vdots \\ \phi_{p_1} &= \{s_{p_1} + 1, s_{p_1} + 2, \dots, s_{p_1} + m_{p_1}\} \\ &= \{m_1 + m_2 + \dots + m_{\{p_1-1\}} + 1, m_1 + m_2 + \dots + m_{\{p_1-1\}} \\ &\quad + 2, \dots, m_1 + m_2 + \dots + m_{\{p_1-1\}} + m_{\{p_1\}}\} \end{aligned}$$

Problems in the form (5) and (6) are equivalent to the following ELCP.

Given A, \bar{c} , and $\phi_1, \phi_2, \dots, \phi_{p_1}$, define the vector $\bar{x} \in \mathbb{R}^n$ such that $\sum_{j \in \phi_i} (A\bar{x} - \bar{c})_i = 0$, with $A\bar{x} \geq \bar{c}$. Definition 5 is given a representation of the equation characteristics of a matrix over the symmetrized max-plus algebra.

Definition 5. [3] Let a matrix $A \in M_n(\mathbb{S})$. For $\det(\lambda \otimes E_n \ominus A) \nabla \epsilon$, be in effect

$$\lambda^{\otimes n} \oplus \bigoplus_{k=1}^n a^k \otimes \lambda^{\otimes(n-k)} \nabla \epsilon \tag{7}$$

with $a_k = (\ominus 0)^{\otimes k} \otimes \bigoplus_{\varphi \in C_n^k} \det A_{\varphi\varphi}$ or

$$a^k = (\ominus 0)^{\otimes k} \bigoplus_{i_1, \dots, i_k \in C_n^k} \bigoplus_{\varphi \in P_k} \text{sgn}(\sigma) \otimes \bigotimes_{r=1}^k a_{i_r i_{\sigma(r)}}, \text{ for } k = 1, 2, \dots, n.$$

The relationship of the eigenvalues of the matrix over the symmetrized max-plus algebra with ELCP is given in the following example.

Example 6

Let a matrix $A = \begin{bmatrix} \ominus 2 & 1 & \epsilon \\ 1 & \ominus 0 & 1 \\ \epsilon & 0 & 2 \end{bmatrix}$. From matrix A , using Definition 5 with the formula :

$$a_k = (\ominus 0)^{\otimes k} \otimes \bigoplus_{\varphi \in C_n^k} \det A_{\varphi\varphi}$$

As a result, the obtained characteristic equation is

$$\lambda^{\otimes 3} \oplus a_1 \otimes \lambda^{\otimes 2} \oplus a_2 \otimes \lambda^{\otimes 1} \oplus a_3 \nabla \epsilon$$

or

$$\lambda^{\otimes 3} \oplus 2 \otimes \lambda^{\otimes 2} \oplus \ominus 4 \otimes \lambda \oplus 4 \nabla \epsilon$$

Next,

$$\lambda^{\otimes 3} \oplus 2 \otimes \lambda^{\otimes 2} \oplus 4 \nabla 2 \otimes \lambda^{\otimes 2} \oplus 4 \otimes \lambda \oplus 4$$

So

$$\lambda^{\otimes 3} \oplus 2 \otimes \lambda^{\otimes 2} \oplus 4 = 2 \otimes \lambda^{\otimes 2} \oplus 4 \otimes \lambda \oplus 4$$

We have a equation

$$\lambda^{\otimes 3} \oplus 2 \otimes \lambda^{\otimes 2} \oplus 4 = p \tag{8}$$

$$2 \otimes \lambda^{\otimes 2} \oplus 4 \otimes \lambda \oplus 4 = p \tag{9}$$

Then multiply $p^{\otimes -1}$ in the equation (8) and equation (9) is obtained

$$\lambda^{\otimes 3} \otimes p^{\otimes -1} \oplus 2 \otimes \lambda^{\otimes 2} \otimes p^{\otimes -1} \oplus 4 \otimes p^{\otimes -1} = 0 \tag{10}$$

Commented [bC8]: S tr t w k z o p f s w j j w j s x a r i f i g t a m i i p j y p r f e z f g m f w e x a u g z f y r j c f q r j s z w j k j w s x d

$$2 \otimes \lambda^{\otimes 2} \otimes p^{\otimes -1} \oplus 4 \otimes \lambda \otimes p^{\otimes -1} \oplus 4 \otimes p^{\otimes -1} = 0 \tag{11}$$

Equation (10) and equation (11) can be viewed as **System Multivariate Polynomial Equation in Max-Plus**, consequently obtained :

$$-C_1 = \begin{bmatrix} -3 & 1 \\ -2 & 1 \\ 0 & 1 \end{bmatrix}; -C_2 = \begin{bmatrix} -2 & 1 \\ -1 & 1 \\ 0 & 1 \end{bmatrix}; -\alpha_1 = \begin{bmatrix} 0 \\ 2 \\ 4 \end{bmatrix}; -\alpha_2 = \begin{bmatrix} 2 \\ 4 \\ 4 \end{bmatrix}$$

Next, we form the matrix $A = \begin{bmatrix} -C_1 \\ -C_2 \end{bmatrix} = \begin{bmatrix} -3 & 1 \\ -2 & 1 \\ 0 & 1 \\ -2 & 1 \\ -1 & 1 \\ 0 & 1 \end{bmatrix}$ and $C = \begin{bmatrix} -d_1 \\ -d_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \\ 4 \\ 2 \\ 4 \\ 4 \end{bmatrix}$

Vector $\bar{x} \in \mathbb{R}^2$ with $\bar{x} = \begin{bmatrix} \lambda \\ p \end{bmatrix}$.

We define ϕ_1 and ϕ_2 , so that:

$$\begin{aligned} \phi_1 &= \{s_1 + 1, s_1 + 2, s_1 + 3\} = \{1, 2, 3\} \\ \phi_2 &= \{s_2 + 1, s_2 + 2, s_2 + 3\} = \{s_1 + m_1 + 1, s_1 + m_1 + 2, s_1 + m_1 + 3\} = \{4, 5, 6\}. \end{aligned}$$

We form ELCP homogenous with : $P = \begin{bmatrix} A & -C \\ 0_{1 \times n} & 1 \end{bmatrix} = \begin{bmatrix} -3 & 1 & 0 \\ -2 & 1 & -2 \\ 0 & 1 & -4 \\ -2 & 1 & -2 \\ -1 & 1 & -4 \\ 0 & 1 & -4 \\ 0 & 0 & 1 \end{bmatrix}$; $\bar{u} = \begin{bmatrix} \bar{x} \\ \alpha \end{bmatrix} = \begin{bmatrix} \lambda \\ p \\ \alpha \end{bmatrix}$

$(Pu)_1(Pu)_2(Pu)_3 + (Pu)_4(Pu)_5(Pu)_6 = 0$ with constraints $(Pu) \geq 0$.

With the algorithm in ELCP (Extended Linear Complementarity Problem), we have the set of all

eigen values $\bar{u} = \kappa \begin{bmatrix} -8 \\ 0 \\ 0 \end{bmatrix}$. In ELCP, $\bar{u} = \begin{bmatrix} \bar{x} \\ \alpha \end{bmatrix} = \begin{bmatrix} \lambda \\ p \\ \alpha \end{bmatrix}$.

Hence, $\lambda = \kappa (-8)$. Let $\lambda_1 = 0, \lambda_2 = -2, \lambda_3 = -4$.

The associated eigenvectors for A are :

1). Let $\lambda = 0$. We have the set of all eigenvectors associated with $\lambda = 0$, that is

$$\left\{ \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \nabla \begin{bmatrix} t \\ 1 \otimes t \\ 1 \otimes t \end{bmatrix} \middle| t \in \mathbb{S} \right\}$$

2). Let $\lambda = -2$. We have the set of all eigenvectors associated with $\lambda = -2$, that is

$$\left\{ \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \nabla \begin{bmatrix} s \\ 1 \otimes s \\ 0 \otimes s \end{bmatrix} \middle| s \in \mathbb{S} \right\}$$

3). Let $\lambda = -4$. We have the set of all eigenvectors associated with $\lambda = -4$, that is

$$\left\{ \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \nabla \begin{bmatrix} r \\ 1 \otimes r \\ 0 \otimes r \end{bmatrix} \middle| r \in \mathbb{S} \right\}$$

We construct matrix

$$P = [v_1 \quad v_2 \quad \dots \quad v_r] = \begin{bmatrix} \ominus 1 & \ominus 1 & \ominus 1 \\ \ominus 2 & \ominus 2 & \ominus 2 \\ 2 & 1 & 1 \end{bmatrix}$$

$$\text{with } D = \begin{bmatrix} 0 & \epsilon & \epsilon \\ \epsilon & -4 & \epsilon \\ \epsilon & \epsilon & -2 \end{bmatrix}$$

$$\text{We have } A \otimes P = \begin{bmatrix} \ominus 2 & 1 & \epsilon \\ 1 & \ominus 0 & 1 \\ \epsilon & \epsilon & \epsilon \end{bmatrix} \otimes \begin{bmatrix} \ominus 1 & \ominus 1 & \ominus 1 \\ \ominus 2 & \ominus 2 & \ominus 2 \\ 2 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 3 & 3 \\ 3 & 2 & 2 \\ 1 & 2 & 2 \end{bmatrix}$$

$$\text{Hence } P \otimes D = \begin{bmatrix} \ominus 1 & \ominus 1 & \ominus 1 \\ \ominus 2 & \ominus 2 & \ominus 2 \\ 2 & 1 & 1 \end{bmatrix} \otimes \begin{bmatrix} 0 & \epsilon & \epsilon \\ \epsilon & -4 & \epsilon \\ \epsilon & \epsilon & -2 \end{bmatrix} = \begin{bmatrix} \ominus 1 & \ominus (-3) & \ominus (-1) \\ \ominus 2 & \ominus (-2) & \ominus 0 \\ 2 & (-3) & (-1) \end{bmatrix}$$

Because $A \otimes P \nabla P \otimes D$, so $P^{-1} \otimes A \otimes P \nabla D$.

4. CONCLUSIONS

A mathematical tool for solving the problem using operations as in conventional algebra, known as the extended linear complementary problem (ELCP), to determine the solution to the characteristic equation. We can form the characteristic equations of matrices over the symmetrized max-plus algebra. The ELCP can be used to system multivariate polynomial equations in max-plus. Because the characteristic equations of matrices over the symmetrized max-plus algebra make to system multivariate polynomial equation in max-plus, we used the ELCP to determine the solution of the characteristic equation of matrices over the symmetrized max-plus algebra.

ACKNOWLEDGEMENT

This work is part of a research project supported by the Research and Community Service Institute (Madiun Campus), Widya Mandala Surabaya Catholic University, Indonesia.

REFERENCES

- [1] Ariyanti, G., Suparwanto, A., & Surodjo, B. (2015). Necessary and Sufficient Conditions for The Solution of The Linear Balanced Systems in The Symmetrized Max Plus Algebra. *Far East J. Math. Sci. (FJMS)*, 97(2), 253-266, 2015.
- [2] De Schutter, B., Boom, T.V.D, Xu, J., Farahani, S.S., Analysis and control of max-plus linear discrete-event systems : An introduction, *Discrete Event Dynamic Systems*, 30, 25-54, 2020
- [3] De Schutter, B. *Max-Algebraic System Theory for Discret Event Systems*, PhD thesis Department of Electrical Engineering Katholieke Universiteit Leuven, Leuven, 1996.
- [4] Farlow, K.G., *Maxplus Algebra*, Master's thesis, Virginia Polytechnic, Institut and State University, Virginia, 2009.
- [5] De Schutter, B., & De Moor, B. The Extended Linear Complementarity Problem and Its Application in The Max-Plus Algebra. *Complementarity and Variational Problems: State of the Art* (M.C. Ferris and J.S. Pang, eds.), Philadelphia, Pennsylvania: SIAM, ISBN 0-89871-391-9, pp. 22-39, 1997.
- [6] Kondo, K. Ultradiscrete sine-Gordon Equation over Symmetrized Max-Plus Algebra. *Symmetry, Integrability and Geometry, Methods and Applications*, SIGMA, 9(068), 1-39, 2013.
- [7] Mora, W. et. al. Invertible Matrices Over Idempotent Semirings. *Chamcuri Journal of Mathematics*, 1(2), 55-61, 2009.

