

Effect of bee pollen on the characteristic of bread incorporated with Monascusfermented durian seeds and rice bran

by Chatarina Yayuk Trisnawati

Submission date: 23-Jul-2023 09:24PM (UTC+0700)

Submission ID: 2135352827

File name: 18-Effect_of_bee_pollen_on_the.pdf (1.08M)

Word count: 4035

Character count: 19917

Effect of bee pollen on the characteristic of bread incorporated with *Monascus*-fermented durian seeds and rice bran

Goberto, M.A., *Trisnawati, C.Y., Nugerahani, I., Srinta, I. and Marsono, Y.

Department of Food Technology, Faculty of Agricultural Technology, Widya Mandala Surabaya Catholic University, Surabaya, East Java, Indonesia

Article history:

Received: 8 March 2022

Received in revised form:

18 May 2022

Accepted: 3 November 2022

Available Online: 19 July

2023

Keywords:

Bread,

Monascus-fermented durian

seeds,

Rice bran,

Bee pollen

5

DOI:

[https://doi.org/10.26656/fr.2017.7\(S1\).36](https://doi.org/10.26656/fr.2017.7(S1).36)

Abstract

Bread incorporated with *Monascus*-fermented durian seeds (MFDS) and rice bran (RB) is a functional food which contains bioactive compounds. Although it is beneficial for human health, it has bitterness and unpleasant aroma that caused by phenolic compound in MFDS and fatty acids in RB. Incorporation of bee pollen in this bread is one way to improve the sensory properties of bread. The aim of this study was to observe the effect of different bee pollen concentrations on the physicochemical and sensory properties of bread incorporated with MFDS and RB. This study used Randomized Block Design with six levels of treatment starting from 0, 0.075, 0.150, 0.225, 0.300 and 0.375%. Data were analyzed by Analysis of Variance with $\alpha = 5\%$. The results indicated that difference in bee pollen concentration significantly affected on physicochemical and sensory properties of bread incorporated with MFDS and RB ($p > 0.05$). Higher bee pollen concentration significantly increased the preference score for taste, aroma and overall acceptance ($p > 0.05$). The best treatment was obtained by using 0.375% bee pollen.

1. Introduction

The development of functional bakery products had been widely studied among food scientists as an approach to meet consumer's demand for bread products with extra health benefits. Trisnawati *et al.* (2019) studied the effect of adding functional ingredients such as rice bran (RB) and *Monascus*-fermented durian seeds (MFDS) into bread. Rice bran contains γ -orizanol, γ -tocotrienol and fibre which can regulate blood sugar levels (Premakumari *et al.*, 2013; Sivamaruthi *et al.*, 2018). MFDS contains monascin pigments that can help to reduce blood sugar level and monacolin-K which helps to reduce cholesterol production (Nugerahani *et al.*, 2017; Faroukh and Baumgärtel, 2019).

Incorporation of MFDS flour and RB flour however reduce the preference score for aroma and taste of bread incorporated with MFDS and RB as reported by Trisnawati *et al.* (2019). RB contains monounsaturated fatty acid and polyunsaturated fatty acid which are easily oxidized into short chain fatty acids which have rancid properties (Cho and Samuel, 2009). MFDS contain secondary metabolites such as tannin and alkaloid which produce bitter and astringent taste (Reginio *et al.*, 2016; Hasim *et al.*, 2019).

One way to cover the unpleasant taste and aroma

caused by RB and MFDS is by incorporating bee pollen. There are several volatile compounds in bee pollen such as esters, hydrocarbon, aldehyde, terpenoid, and keton (Neto *et al.*, 2017). Incorporation of bee pollen also may increase furan production, such as furfural and pyrazine in the final product. These compounds produce caramel, floral and fruity flavors (Conte *et al.*, 2020). Unsaturated fatty acid, phytosterol and phospholipid in bee pollen can increase hypoglycemic activity (Komosinska-Vashev *et al.*, 2015). The objective of this study was to observe the effect of bee pollen incorporation on the characteristic of bread incorporated with MFDS and RB.

2. Materials and methods

2.1 Materials

Materials that were used for making bread incorporated with MFDS and RB were bread flour, rice bran flour, instant dry yeast, instant full cream milk powder, granulated sugar, bread improver, table salt, mineral water, margarine which were purchased from local distributor, "Mirah Delima" multiflora bee pollen which was purchased from Mirah Delima Bee Farm and also MFDS flour which was produced in Laboratory of Food Industrial Microbiology, Widya Mandala Surabaya Catholic University. Materials that were used for producing MFDS flour were Petruk durian seeds, pure

*Corresponding author.

Email: chatarina@ukwms.ac.id

culture of *Monascus purpureus* M9, Ca(OH)₂, distilled water and potato dextrose agar (Merck 1.10130.0500).

2.2 Preparation of bread incorporated with *Monascus-fermented durian seeds and rice bran*

Sample preparation began by mixing all of the dry ingredients except table salt for 1 min and followed by adding water. The mixing process then continued for 10 mins. Margarine and table salt was then added and the mixing process continued for 5 mins. The dough was fermented at 26°C for 30 mins. After the dough was shaped into a loaf the dough was proofed at 26°C for 90 mins. The bread was baked in the oven at 180°C for 30 mins then cooled at 26°C for 60 mins. Table 1 shows the composition of bread incorporated with MFDS and RB with different levels of bee pollen concentration.

2.3 *Monascus-fermented durian seeds preparation*

Preparation of MFDS flour began with durian seeds sortation. Durian seeds then washed with water. After that, the durian seeds were boiled with 5% Ca(OH)₂ solution at 85-90°C for 10 mins [durian seeds: 5% Ca(OH)₂ solution = 1:1 (w/v)]. The durian seeds were removed from the Ca(OH)₂ solution, then washed with water. The durian seeds were then cut into 1 cm × 1 cm × 1 cm cubes and went through first drying process at 45°C for 40 mins. The durian seeds were then weighed into 50 g. After that, durian seeds were sterilized at 121°C, 15 lbs/inch² for 10 mins and cooled down at 26°C for 30 mins. The sterilized durian seeds then inoculated with *Monascus purpureus* M9 starter (5% v/w) and were put under aerobic fermentation at 30±1°C for 14 days to produce MFDS. The MFDS were dried at 45°C for 24 hrs, grinded and then sifted with 40 mesh sifters to get MFDS flour (Puspitadewi et al., 2016).

2.4 Moisture content analysis

Moisture content determination was carried out using thermogravimetric method according to AOAC

925.10 (1990). The moisture content of bread incorporated with MFDS and RB was determined using following equation:

$$\text{Moisture content (\%)} = \frac{\text{sample weight (g)} - \text{weight loss (g)}}{\text{sample weight (g)}} \times 100\%$$

2.5 Specific volume

Specific volume determination was carried out using seed displacement method according to Nwosu et al. (2014). An empty container was filled with foxtail millet seeds until overflowed and seeds which were above the container rim were then removed using straight ruler. All the foxtail millet seeds in the container were then poured into a measuring cylinder to measure the volume of the container (V₁). These steps were then repeated except that a loaf of bread sample was already inside the container before it was filled with seeds to obtain V₂.

$$\text{Specific volume (cm}^3/\text{g)} = \frac{V_1 \text{ (mL)} - V_2 \text{ (mL)}}{W \text{ (g)}}$$

2.6 Texture analysis

Prior to analysis, bread samples were cut into 12 mm thickness. Texture analysis was carried out using texture analyzer (TA-XT Plus, Stable Micro System) and performed by two sequential compression events (probe: P36/R, pre-test speed: 5 mm s⁻¹, test speed: 1.5 mm s⁻¹, post-test speed 10 mm s⁻¹, distance: 12 mm, time: 3 s, trigger type: auto, trigger force: 5 g).

Hardness was defined as force that needed to achieve deformation during first compression. Cohesiveness was defined as ratio of the positive first area of the second peak to the first peak. Springiness was defined as ability of the bread to recover in height during the time elapsed between end of first compression and start of the second compression cycle (Dvořáková et al., 2012).

2.7 Color analysis

Table 1. Formulation of bread incorporated with *Monascus-fermented durian seeds and rice bran*.

Ingredients/g	B ₀	B ₁	B ₂	B ₃	B ₄	B ₅
Bread flour	179.85	179.85	179.85	179.85	179.85	179.85
Mineral water	124	124	124	124	124	124
Rice bran flour	20	20	20	20	20	20
Granulated sugar	10	10	10	10	10	10
Margarine	8	8	8	8	8	8
Full cream milk powder	4	4	4	4	4	4
Instant dry yeast	3	3	3	3	3	3
Table salt	2	2	2	2	2	2
Bread improver	0.6	0.6	0.6	0.6	0.6	0.6
MFDS flour	0.15	0.15	0.15	0.15	0.15	0.15
Bee pollen	0	0.075	0.150	0.225	0.300	0.375

B₀ : 0% bee pollen, B₁ : 0.075% bee pollen, B₂ : 0.150% bee pollen, B₃ : 0.225% bee pollen, B₄ : 0.300% bee pollen, B₅ : 0.375% bee pollen.

Color analysis was carried out by measuring the lightness (L*), redness (a*), yellowness (b*), chroma (*C) and hue (°H) of the crumb of bread incorporated MFDS and RB with color reader (Minolta CR-10 Chroma Meter).

2.8 Sensory evaluation

Sensory evaluation was carried out with 50 untrained panelists. Each panelist was asked to evaluate each parameter using a 7-point hedonic scale (Sivile and Sidel, 2004). Scores were assigned in a range 1-7 (1 = extremely dislike; 2 = dislike; 3 = slightly dislike; 4 = neither like nor dislike; 5 = slightly like; 6 = like; 7 = extremely like). Each panelist received four slices of bread incorporated with *Monascus*-fermented durian seed and rice bran which were cut into 5 cm × 5 cm × 1 cm. The tested parameters consisted of preference score for color, aroma, taste, and overall acceptance.

2.9 Statistical analysis

Data were generated in triplicate and subjected to Analysis of Variance (ANOVA) with $\alpha = 5\%$ and followed by Duncan's Multiple Range Test (DMRT) with $\alpha = 5\%$ using SPSS software. Data collected from sensory evaluation were also subjected to spider-web test using Microsoft Excel 2013.

3. Results and discussion

3.1 Moisture content

According to Table 2, incorporation of bee pollen to bread with MFDS and RB resulted in moisture content ranging from 40.29% to 59.25% while Badan Standarisasi Nasional (1995) suggested that maximum moisture content of bread is 40% to support shelf-life against microbial deterioration. Rice bran contains 7-11% fiber (Henderson et al., 2012). Ability of fiber to bind with water then resulted in increased moisture content of bread (Sangle et al., 2017). Bee pollen

Table 2. Effect of bee pollen on the moisture content of bread incorporated with *Monascus*-fermented durian seeds and rice bran.

Bee pollen concentration	Moisture Content (%)
B ₀	40.29±0.11 ^a
B ₁	44.20±0.07 ^b
B ₂	47.34±0.15 ^c
B ₃	51.35±0.16 ^d
B ₄	54.23±0.15 ^e
B ₅	59.25±0.16 ^f

Values are presented as mean±SD (n = 4). Values with different superscripts within column are statistically significantly different evaluated by ANOVA followed by DMRT test ($\alpha = 0.05$). B₀: 0% bee pollen, B₁: 0.075% bee pollen, B₂: 0.150% bee pollen, B₃: 0.225% bee pollen, B₄: 0.300% bee pollen, B₅: 0.375% bee pollen.

[https://doi.org/10.26656/fr.2017.7\(S1\).36](https://doi.org/10.26656/fr.2017.7(S1).36)

contains glucose, fructose and sucrose which were able to form hydroxyl bond with water molecules (Komosinska-Vassev et al., 2013). The bee pollen is high in lysine, arginine, cysteine, tryptophan, and tyrosine which were hydrophilic amino acid (Taha et al., 2017). Cante et al. (2020) also reported that moisture content of gluten free bread increased as bee pollen concentration increased.

3.2 Physical properties

Increase of specific volume in bread samples with 0-0.15% bee pollen as seen in Table 3 was due to additional sugar from bee pollen which are useful CO₂ production by yeast. The CO₂ pushed the gluten matrix to become thinner and resulted in reduced hardness as seen in Table 4. More CO₂ gas also increased viscoelasticity of gluten matrix hence springiness increased. Bread samples with 0.225-0.375% bee pollen showed a decrease in specific volume due to competition in binding water between the hydrophilic molecules (protein and sugar) in bee pollen and gluten forming protein. Hence the gluten matrix became weaker and could not retain CO₂ as much as bread with 0-0.15%. Lower retention of CO₂ also resulted in bread with thicker gluten matrix and lower viscoelasticity hence increase bread's hardness and decrease bread's springiness.

Table 3. Effect of bee pollen on the specific volume of bread incorporated with *Monascus*-fermented durian seeds and rice bran.

Bee pollen concentration	Specific volume (cm ³ /g)
B ₀	3.85±0.03 ^c
B ₁	3.97±0.01 ^d
B ₂	4.07±0.06 ^c
B ₃	3.81±0.02 ^c
B ₄	3.34±0.06 ^b
B ₅	3.23±0.05 ^a

Values are presented as mean±SD (n = 4). Values with different superscripts within column are statistically significantly different evaluated by ANOVA followed by DMRT test ($\alpha = 0.05$). B₀: 0% bee pollen, B₁: 0.075% bee pollen, B₂: 0.150% bee pollen, B₃: 0.225% bee pollen, B₄: 0.300% bee pollen, B₅: 0.375% bee pollen.

The results in Table 5 showed that as bee pollen concentration increased, L* value decreased and a*, b*, C and H value of bread samples increased. Lower L* might be attributed to incorporation of rice bran flour since its color is brown and Maillard reaction which occurred during baking process. Starowicz and Zieliński (2019) reported that free amine groups in protein and carbonyl groups in sugars went through Maillard reaction to produce melanoidin which reduce L* value of food. Brown's base colors are grounded with red and yellow which resulted in increased of a* and b* value.

Table 4. Effect of bee pollen on textural properties for bread incorporated with *Monascus*-fermented durian seeds and rice bran.

Textural properties	B ₀	B ₁	B ₂	B ₃	B ₄	B ₅
Hardness (g)	790.452 ±7.920 ^c	769.631 ±10.910 ^b	665.708 ±5.590 ^a	833.009 ±10.450 ^d	1059.045 ±17.130 ^e	1254.505 ±15.140 ^f
Cohesiveness	0.643 ±0.010 ^c	0.662 ±0.010 ^d	0.677 ±0.010 ^e	0.646 ±0.010 ^c	0.620 ±0.010 ^b	0.596 ±0.010 ^a
Springiness	0.790 ±0.011 ^c	0.819 ±0.003 ^d	0.833 ±0.005 ^e	0.802 ±0.001 ^d	0.768 ±0.002 ^b	0.758 ±0.003 ^a

Values are presented as mean±SD (n = 4). Values with different superscripts within column are statistically significantly different evaluated by ANOVA followed by DMRT test ($\alpha = 0.05$). B₀: 0% bee pollen, B₁: 0.075% bee pollen, B₂: 0.150% bee pollen, B₃: 0.225% bee pollen, B₄: 0.300% bee pollen, B₅: 0.375% bee pollen.

Table 5. Effect of bee pollen on color of bread incorporated with *Monascus*-fermented durian seeds and rice bran.

Color evaluation	B ₀	B ₁	B ₂	B ₃	B ₄	B ₅
Lightness (L*)	69.2±0.25 ^d	67.6±0.39 ^c	66.8±0.99 ^b	66.5±0.20 ^b	65.7±0.55 ^a	65.3±0.17 ^a
Redness (a*)	3.2±0.13 ^a	3.2±0.10 ^a	3.3±0.18 ^a	3.4±0.13 ^b	3.6±0.10 ^c	3.8±0.03 ^d
Yellowness (b*)	15.9±0.13 ^a	17.4±0.28 ^b	18.2±0.22 ^c	18.6±0.45 ^c	19.5±0.45 ^d	20.1±0.32 ^e
Chroma (C)	16.2±0.15 ^a	17.7±0.28 ^b	18.5±0.22 ^c	18.9±0.45 ^c	19.8±0.46 ^d	20.5±0.33 ^e
Hue (°H)	78.7±0.43 ^a	79.6±0.48 ^b	79.8±0.53 ^b	79.7±0.38 ^b	79.5±0.15 ^b	79.2±0.16 ^b

Values are presented as mean±SD (n = 4). Values with different superscripts within column are statistically significantly different evaluated by ANOVA followed by DMRT test ($\alpha = 0.05$). B₀: 0% bee pollen, B₁: 0.075% bee pollen, B₂: 0.150% bee pollen, B₃: 0.225% bee pollen, B₄: 0.300% bee pollen, B₅: 0.375% bee pollen.

Value of b* was also influenced by the yellow pigment β -carotene in bee pollen.

3.3 Sensory evaluation

Table 6 shows that the preference score for color of the bread samples was statistically decreased while preference score of taste, aroma and overall acceptability were statistically increased as BP concentration increased. Spider web in Figure 1. test showed that 0.375% BP was the best concentration for bread incorporated with MFDS and RB even though the color was the least favorable.

Bee pollen consisted of several volatile compounds which comprised of hydrocarbon, esters, terpenoid and alcohol which produced floral and fruity aroma (Neto *et al.*, 2017). Increased in bee pollen concentration also resulted in more Maillard reaction product resulted such as pyrazine that produced nutty and roasted aroma; furan that produced sweet and caramel aroma; acetylpyridine that produced cracker-malty aroma; and pyrrol that produced nutty aroma (van Boekel, 2006). Those volatile compounds together were able to tone down the rancid aroma, which came from the rice bran.

Table 6. Effect of bee pollen on preference score for bread incorporated with *Monascus*-fermented durian seeds and rice bran.

Sensory evaluation	B ₀	B ₁	B ₂	B ₃	B ₄	B ₅
Color	5.45±1.20 ^d	5.40±0.93 ^d	5.05±1.20 ^{cd}	4.85±0.98 ^{bc}	4.55±1.15 ^{ab}	4.33±1.54 ^a
Aroma	3.35±0.92 ^a	4.43±1.36 ^b	4.83±1.01 ^b	4.53±1.22 ^b	4.73±1.26 ^b	5.35±0.95 ^c
Taste	3.33±0.92 ^a	3.38±0.90 ^a	4.10±1.01 ^b	4.48±1.26 ^b	4.93±1.23 ^c	5.33±0.76 ^d
Overall acceptance	3.35±0.77 ^a	4.33±1.05 ^b	4.80±1.18 ^c	4.93±1.21 ^{cd}	5.05±1.20 ^{cd}	5.33±0.97 ^d

Values are presented as mean±SD (n = 50). Values with different superscripts within column are statistically significantly different evaluated by ANOVA followed by DMRT test ($\alpha = 0.05$). B₀: 0% bee pollen, B₁: 0.075% bee pollen, B₂: 0.150% bee pollen, B₃: 0.225% bee pollen, B₄: 0.300% bee pollen, B₅: 0.375% bee pollen.

4. Conclusion

The most suitable bee pollen concentration for bread incorporated with MFDS and RB is 0.375% according to the sensory evaluation. The overall acceptance score of this bread represented that panelist slightly like the bread. This study showed that bee pollen has the potential to improve taste and aroma of bread incorporated with MFDS and RB. Further study to improve specific volume and textural properties of this bread is suggested.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

This research was funded by Ministry of Research, Technology and Higher Education, Republic of Indonesia through competitive research "Penelitian Terapan Unggulan Perguruan Tinggi (PTUPT)" with contract number 130X/WM01.5/N/2020.

References

AOAC (Association of Official Analytical Chemists).

- (1990). Official Methods of Analysis of the Association of Official Analytical Chemists. 15th ed. Arlington, Virginia, USA: AOAC Inc.
- Badan Standarisasi Nasional. (2009). Tepung terigu sebagai bahan makanan (SNI 3751:2009). Retrieved from website: <https://extranet.who.int/nutrition/gina/sites/default/files/IDN%202009%20Tepung%20terigu%20sebagai%20bahan%20makanan%20-%20wheat%20flour.pdf> [In Bahasa Indonesia].
- Cho, S.S. and Samuel, P. (2009). Fiber Ingredients: Food Application and Health Benefits. 1st ed. Boca Raton, USA: CRC Press. <https://doi.org/10.1201/9781420043853>
- Conte, P., Del Caro, A., Urgeghe, P.P., Pertretto, G.L., Montanari, L., Piga, A. and Fadda, C. (2020). Nutritional and aroma improvement of gluten-free bread: is bee pollen effective? *LWT-Food Science and Technology*, 118, 108711. <https://doi.org/10.1016/j.lwt.2019.108711>
- Dvořáková, P., Burešová, I. and Kráčmar, S. (2012). Textural properties of bread formulations based on buckwheat and rye flour. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 6(5), 61-68. <https://doi.org/10.11118/actaun201260050061>
- Faroukh, A. and Baumgärtel, C. (2019). Mini – review: Medication safety of red yeast rice products. *International Journal of General Medicine*, 12, 167-171. <https://doi.org/10.2147/IJGM.S202446>
- Hasim, Faridah, D.N. and Mithania, D. (2019). Penambahan angkak dan bekatul pada agar-agar sebagai alternatif cemilan pengikat kolesterol. *Jurnal Mutu Pangan*, 6(2), 85-90. <https://doi.org/10.29244/jmpi.2019.6.85> [In Bahasa Indonesia].
- Henderson, A.J., Ollila, C.A., Kumar, A., Borresen, E.C., Raina, K., Agarwal, R. and Ryan, E.P. (2012). Chemopreventive properties of dietary rice bran: current status and future prospects. *Advanced Nutrition*, 3(5), 643-653. <https://doi.org/10.3945/an.112.002303>
- Komosinska-Vassev, K., Olczyk, P., Kazmierczak, J., Mencner, L. and Olczyk, K. (2015). Bee pollen: chemical composition and therapeutic application. *Evidence-Based Complementary and Alternative Medicine*, 2015, 297425. <https://doi.org/10.1155/2015/297425>
- Neto, J.S.L., Lopes, J.A.D., Neto, J.M.M., de Lima, S.G., da Luz, C.F.P. and Citó, A.M.G.L. (2017). Volatile compounds and palynological analysis from pollen pots of stingless bees from the mid-north region of Brazil. *Brazilian Journal of Pharmaceutical Sciences*, 53(2), e14093. <https://doi.org/10.1590/s2175-97902017000214093>
- Nugerahani, I., Sutedja, A.M., Srianta, I., Widharna, R.M. and Marsono, Y. (2017). *In vivo* evaluation of *Monascus*-fermented durian seed for antidiabetic and antihypercholesterol agent. *Food Research*, 1(3), 83-88. <https://doi.org/10.26656/fr.2017.3.023>
- Nwosu, U.L., Elochukwu, U.C. and Onwurah, C.O. (2014). Physical characteristics and sensory quality of bread produced from wheat/African oil bean flour blends. *African Journal of Food Science*, 8(6), 351-355. <https://doi.org/10.5897/AJFS2013.1079>
- Premakumari, S., Balasasirekha, R., Gomathi, K., Supriya, S., Alagusundaram, K. and Mohan, R.J. (2013). Evaluation of organoleptic properties and glycemic index of recipes with rice bran. *International Journal of Current Research and Review*, 5(7), 1-8.
- Puspitadewi, S.R.D., Srianta, I. and Kusumawati, N. (2016). Pola produksi pigmen *Monascus* oleh *Monascus sp.* KJR 2 pada media biji durian varietas Petruk melalui fermentasi padat. *Jurnal Teknologi Pangan dan Gizi*, 15(1), 36-42. [In Bahasa Indonesia].
- Reginio, F.C.Jr., Hurtada, W.A. and Dizon, E.I. (2016). Quality and acceptability of *Monascus* biopigment beverage. *International Food Research Journal*, 23 (4), 1492-1500.
- Sangle, J.K., Sawate, A.R., Patil, B.M. and Kshirshagar, R. B. (2017). Nutritional and sensory quality of wheat bread supplemented with stabilized rice bran. *Food Science Research Journal*, 8(1), 112-118. <https://doi.org/10.15740/HAS/FSRJ/8.1/112-118>
- Sivamaruthi, B.S., Kesika, P. and Chaiyasut, C. (2018). A comprehensive review on anti-diabetic property of rice bran. *Asian Pacific Journal of Tropical Biomedicine*, 8(1), 79-84. <https://doi.org/10.4103/2221-1691.221142>
- Starowicz, M. and Zieliński, H. (2019). How Maillard reaction influences sensorial properties (color, flavor, and texture) of food products? *Food Reviews International*, 35(8), 707-725. <https://doi.org/10.1080/87559129.2019.1600538>
- Stone, H. and Sidel, J. (2004). Sensory Evaluation Practices. 3rd ed. Redwood City, California, USA: Academic Press.
- Taha, E.K.A., Kahtani, S.A. dan Taha, R. (2017). Protein content and amino acids composition of bee pollens from major floral sources in Al-Ahsa, Eastern Saudi Arabia. *Saudi Journal of Biological Sciences*, 26(2), 232-237. <https://doi.org/10.1016/j.sjbs.2017.06.003>

- Trisnawati, C.Y., Srianta, I., Nugerahani, I. and Marsono, Y. (2019). Incorporating *Monascus*-fermented durian seeds and rice bran into bread: study on the bread physicochemical and sensory properties. *Food Research*, 3(3), 280-284. [https://doi.org/10.26656/fr.2017.3\(3\).015](https://doi.org/10.26656/fr.2017.3(3).015)
- van Boekel, M.A.J.S. (2006). Formation of flavour compounds in the Maillard reaction. *Biotechnology Advances*, 24(2), 230-233. <https://doi.org/10.1016/j.biotechadv.2005.11.004>

Effect of bee pollen on the characteristic of bread incorporated with Monascusfermented durian seeds and rice bran

ORIGINALITY REPORT

11 %
SIMILARITY INDEX

7 %
INTERNET SOURCES

9 %
PUBLICATIONS

5 %
STUDENT PAPERS

PRIMARY SOURCES

1 www.idosi.org Internet Source **2** %

2 "Gluten-free Bread Technology", Springer Science and Business Media LLC, 2021 Publication **1** %

3 itjfs.com Internet Source **1** %

4 Submitted to Universitas Jember Student Paper **1** %

5 ir.unimas.my Internet Source **1** %

6 link.springer.com Internet Source **1** %

7 Chuan-Mei Yeh, Bi-Yu Kao, Hsuan-Jung Peng. " Production of a Recombinant Type 1 Antifreeze Protein Analogue by . and Its Applications on Frozen Meat and Frozen Dough ", Journal of Agricultural and Food Chemistry, 2009 **1** %

8

Aleksandar Ž. Kostić, Danijel D. Milinčić, Miroljub B. Barać, Mohammad Ali Shariati et al. "The Application of Pollen as a Functional Food and Feed Ingredient—The Present and Perspectives", *Biomolecules*, 2020

Publication

1 %

9

chimie-biologie.ubm.ro

Internet Source

1 %

10

Pimchada Itthivadhanapong, Srinual Jantathai, Gerhard Schleining. "Improvement of physical properties of gluten-free steamed cake based on black waxy rice flour using different hydrocolloids", *Journal of Food Science and Technology*, 2016

Publication

1 %

11

www.hindawi.com

Internet Source

1 %

12

Metin Guldaz. "Effects of Royal Jelly and Bee Pollen on the Growth of Selected Probiotic Bacteria (*Bf. animalis* Spp. *Lactis*, *L. acidophilus* and *L. casei*)", *Journal of Apicultural Science*, 2016

Publication

1 %

13

Submitted to University of Hong Kong

Student Paper

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On