


[VIEW ARTICLES](#)[ISSUES](#)[ABOUT](#)[FOR AUTHORS](#)[FOR REFEREES](#)

FOOD RESEARCH

Volume 6, Issue 3

June 2022

Mini Review

A review on production, application, and toxicological analyses of nanocrystalline cellulose as a novel fat replacer food additive

Aida Safina, A., Chin N.L., Nur Akmal, I., Nor Nadiah, M.Y. and Yus Aniza, Y.

Available Online: 5 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).231](https://doi.org/10.26656/fr.2017.6(3).231)

Aida Safina *et al.* reviewed on the production, application and toxicological analyses of nanocrystalline cellulose as a novel fat replacer food additive.

Full Papers

Moisture content and application rates of inert dust: effects on dust and wheat physical properties

Yao, K.D., Subramanyam, B. and Maghirang, R.G.

Available Online: 5 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).280](https://doi.org/10.26656/fr.2017.6(3).280)

Yao *et al.* studied on the effects on dust and wheat physical properties on the moisture content and application rates of inert dust.

Preparation of a protein drink from fish protein hydrolysate obtained from tilapia skin waste

Osiriphun, S., Wangtueai, S., Rachatanaphun, P. and Jirarattanarangsri, W.

Available Online: 5 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).342](https://doi.org/10.26656/fr.2017.6(3).342)

Osiriphun *et al.* developed a protein drink from fish protein hydrolysate obtained from tilapia skin waste.

The addition of lactic acid bacteria in the soybean soaking process of tempeh

Magdalena, S., Hogaputri, J.E, Yulandi, A. and Yogiara, Y.

Available Online: 5 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).304](https://doi.org/10.26656/fr.2017.6(3).304)

The addition of lactic acid bacteria in the soybean soaking process of tempeh was studied by Magdalena *et al.*

Breadfruit (*Artocarpus altilis*) starch-based nanoparticle formation through dropwise mixing nanoprecipitation

Harsanto, B.W., Pranoto, Y., Supriyanto and Kartini, I.

Available Online: 8 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).308](https://doi.org/10.26656/fr.2017.6(3).308)

Harsanto *et al.* formed breadfruit (*Artocarpus altilis*) starch-based nanoparticle through dropwise mixing nanoprecipitation.

Physical properties and sensory acceptability of gum arabic-coated cherry tomato fruit during storage

Sumonsiri, N., Charoensantisuk, K., Paanoi, N. and Kittayangkul, P.

Available Online: 8 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).395](https://doi.org/10.26656/fr.2017.6(3).395)

The physical properties and sensory acceptability of gum arabic-coated cherry tomato fruit during storage was evaluated by Sumonsiri *et al.*

Physicochemical properties of yam starches from fifty-five lines of *Dioscorea* species

Olayide, S.L., Kehinde, O.S., Adeolu, A.A., Olushola, S.A., Nishinari, K. and Simphiwe, M.N.

Available Online: 8 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).224](https://doi.org/10.26656/fr.2017.6(3).224)

Olayide *et al.* evaluated the physicochemical properties of yam starches from fifty-five lines of *Dioscorea* species

Whey protein concentrate mixed beverages and plasma amino acid response in young males

Klaewkla, J., Hudthagosol, C., Chaijenkij, K., Panya, A., Sang-ngoen, D., Phonsatta, N. and Kaewkul K.

Available Online: 8 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).630](https://doi.org/10.26656/fr.2017.6(3).630)

Klaewkla *et al.* evaluated the plasma amino acid response in young males consuming whey protein concentrate mixed beverages.

Antioxidant activities and polyphenol compounds of kenaf leaf tea infusion after in-vitro gastrointestinal digestion and consumer perception survey

Goh, K.M., Lee, S.W. and Nyam, K.L.

Available Online: 11 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).590](https://doi.org/10.26656/fr.2017.6(3).590)

The antioxidant activities and polyphenol compounds of kenaf leaf tea infusion after in-vitro gastrointestinal digestion and consumer perception survey was conducted by Goh *et al.*

Assessment of *Pseudomonas aeruginosa* biofilm-forming capacities from drinking water in water vending machine

Elexson, N., Sabrina, H., Dalene, L., Eddy, B., Nurul, F.R., Nasra, P., Grace, B., Nick, L., Amirah, Z.J., Nur, D.Z., Dayang, N.A.B., Manju, S. and Tunung, R.

Available Online: 11 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).324](https://doi.org/10.26656/fr.2017.6(3).324)

Elexson *et al.* assessed the biofilm-forming capacities of *Pseudomonas aeruginosa* from drinking water in water vending machine.

Isolation of active compound from *Nephelium lappaceum* L. rind as an antioxidant

Nurani, L.H., Edityaningrum, C.A., Suhaera, Windarsih, A., Riyanto, S. and Rohman, A.

Available Online: 11 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).331](https://doi.org/10.26656/fr.2017.6(3).331)

Nurani *et al.* isolated and studied the active compound from *Nephelium lappaceum* L. rind as an antioxidant.

Halal food: a social responsibility on cartel meat issue in Malaysia

Mohd Riza, N.S., Md Ariffin M.F., Hamdan, M.N. and Ramli, N.

Available Online: 11 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).277](https://doi.org/10.26656/fr.2017.6(3).277)

Mohd Riza *et al.* evaluated the social responsibility on cartel meat issue on halal in Malaysia.

A comparative study of the physico-chemical properties of prominent cocoa bean in Southern Vietnam

Lam, T.V.H., Phan, T.B.T., Truong, T.N. and Ha, T.T.

Available Online: 14 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).359](https://doi.org/10.26656/fr.2017.6(3).359)

Lam *et al.* performed a comparative study on the physico-chemical properties of prominent cocoa bean in Southern Vietnam

Comparison of the mass tissue strength of strawberry fruit between vertical and horizontal compaction

Ansar, A., Murad, M., Sukmawaty, S. and Ilmaknun, L.

Available Online: 14 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).373](https://doi.org/10.26656/fr.2017.6(3).373)

The meat tissue strength of strawberry fruit between vertical and horizontal compaction was compared by Ansar *et al.*

Optimum condition of roasting process of Liberica coffee towards the local and international preference

Halim-Lim, S. A., Wan-Mohtar, W.A.A.Q.I., Surapinchai, S. and Azizan, N.A.Z.

Available Online: 14 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).340](https://doi.org/10.26656/fr.2017.6(3).340)

The optimum conditions of roasting process of Liberica coffee towards the local and international preference was evaluated by Halim-Lim *et al.*

Optimization of heat treatment and pH of red and white pear cactus [*Opuntia ficus-indica* (L.) mill.] fruit juice using response surface methodology

Abdulkadir, N., Solomon, W.K. and Woldetsadik, K.

Available Online: 14 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).306](https://doi.org/10.26656/fr.2017.6(3).306)

Abdulkadir *et al.* optimized the heat treatment and pH of red and white pear cactus [*Opuntia ficus-indica* (L.) mill.] fruit juice using response surface methodology

Meatball model of porcine DNA detection by TaqMan probe real-time PCR

Sajali, N., Ting, S.M.L., Koh, C.C., Desa, M.N.M., Wong, S.C. and Bakar, S.

Available Online: 19 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).384](https://doi.org/10.26656/fr.2017.6(3).384)

Sajali *et al.* evaluated the meatball model of porcine DNA detection by TaqMan probe real-time PCR.

Simultaneous determination of nitrite and nitrate in meat and meat products using ion-exchange chromatography

Mazumdar, R.M., Sharif, M., Khan, T.A., Rahman, M.M. and Abdullah A.T.M.

Available Online: 19 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).339](https://doi.org/10.26656/fr.2017.6(3).339)

Mazumdat *et al.* performed a simultaneous determination of nitrite in meat and meat products using ion-exchange chromatography.

Prediction of diffusion coefficient for losses of minerals from potato during frying

Samir, Z.T., Saeed, S.K., Mohammed, N.K., and Abdul-Rezzak, R.K.

Available Online: 19 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).382](https://doi.org/10.26656/fr.2017.6(3).382)

Samir *et al.* predicted the diffusion coefficient for losses of minerals from potato during frying.

Effects of lyotropic series salts on the functional properties of bambara groundnut (*Voandzeia subterranean*) protein isolate

Lawal, O.S., Sodeinde, K.O., Adediran, A.A., Nishinari, K., Olatunji, O.S., and Ayanda, O.S.

Available Online: 19 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).771](https://doi.org/10.26656/fr.2017.6(3).771)

The effects of lyotropic series salts on the functional properties of bambara groundnut (*Voandzeia subterranean*) protein isolate were studied by Lawal *et al.*

Physicochemical properties of post laying hen breast meat thawed using various methods

Dwiloka, B., Setiani, B.E., Pramono Y.B., Prihatiningsih, R., Nurussyifa, S.Y. and Puspitoasih, A.D.

Available Online: 22 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).296](https://doi.org/10.26656/fr.2017.6(3).296)

The physicochemical properties of post laying hen breast meat thawed using various methods were studied by Dwiloka *et al.*

Anthropometry indicators that are most related to metabolic profiles in female college students

Dieny, F.F., Rose S., Tsani, A.F.A., Jauharany, F.F. and Fitranti, D.Y.

Available Online: 22 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).250](https://doi.org/10.26656/fr.2017.6(3).250)

Dieny *et al.* evaluated the anthropometry indicators that are most related to metabolic profiles in female college students.

Screening of *Lactobacillus rhamnosus*-producing gamma aminobutyric acid (GABA) isolated from Sumbawa mare milk and its potential application to increase GABA content in fermented milk

Nursini, N.W., Antara, N.S., Sugitha, I.M. and Sujaya, I.N.

Available Online: 22 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).380](https://doi.org/10.26656/fr.2017.6(3).380)

Nursini *et al.* screened *Lactobacillus rhamnosus*-producing gamma aminobutyric acid (GABA) isolated from Sumbawa mare milk and studied on its potential to increase GABA content in fermented milk.

The effect of rotary drying temperature on drying characteristic and antioxidant activity of *Etlingera elatior* Jack

Simanjuntak, M.E., Ristiarini, S. and Widyawati, P.S.

Available Online: 22 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).333](https://doi.org/10.26656/fr.2017.6(3).333)

The effect of rotary drying temperature on the drying characteristics and antioxidant activity of *Etlingera elatior* Jack was studied by Simanjuntak *et al.*

The effect of differences in ozonation time and storage temperature on physical, chemical, and sensory characteristics of Japanese spinach (*Spinacia oleracea* L.)

Prabawa, S., Safitri, D.I., Rofandi, H., Amanto, B. and Yudhistira, B.

Available Online: 26 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).350](https://doi.org/10.26656/fr.2017.6(3).350)

Prabawa *et al.* studied the effect of differences in ozonation time and storage temperature on physical, chemical and sensory characteristics of Japanese spinach (*Spinacia oleracea* L.)

Development and acceptability of value-added products from green mussel (*Perna viridis*) in Samar, Philippines

Sorio, J.C. and Arcales, J.A.A.

Available Online: 26 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).320](https://doi.org/10.26656/fr.2017.6(3).320)

Sorio and Arcales developed and evaluated the acceptability of value-added products from green mussel (*Perna viridis*) in Samar, Philippines.

Identification and quantification of sodium benzoate in soft drinks available in Tangail region by high-performance liquid chromatography

Esrafil, M., Akter, S., Alam, M.J., Haque, M.A., Zubair, M.A. and Khan, M.S.H

Available Online: 26 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).348](https://doi.org/10.26656/fr.2017.6(3).348)

Esrafil *et al.* identified and quantified sodium benzoate in soft drinks available in Tangail region using high-performance liquid chromatography.

Comparison between Polymerase Chain Reaction and Loop Mediated Isothermal Amplification for the detection of *Staphylococcus aureus* in food

Bashar, Q.K., Aziz, A.Z. and Kadhim, N.I.

Available Online: 29 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).335](https://doi.org/10.26656/fr.2017.6(3).335)

Bashar *et al.* compared Polymerase Chain Reaction and Loop Mediated Isothermal Amplification for the detection of *Staphylococcus aureus* in food.

Studies on the effect of methionine level on cheese colour as a solid substrate of *Monascus purpureus* JK2 fermentation

Sulandari, L., Utami, T., Hidayat, C. and Rahayu, E.S.

Available Online: 29 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).422](https://doi.org/10.26656/fr.2017.6(3).422)

The effect of methionine level on cheese colour as a solid substrate of *Monascus purpureus* JK2 fermentation was study by Sulandari *et al.*

Effectiveness of banana juice (*Musa acuminata* Linn.) on blood pressure, blood sugar levels, and low-density lipoprotein in elderly

Fitri, Y., Suryana, S., Ahmad, A., Hendra, A., Fitriyaningsih, E., Arnisam and Yuniato, A.E.

Available Online: 29 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).213](https://doi.org/10.26656/fr.2017.6(3).213)

The effectiveness of banana juice (*Musa acuminata* Linn.) on blood pressure, blood sugar levels, and low-density lipoprotein in elderly was studied by Fitri *et al.*

Development and quality evaluation of jelly coated tapai as a cocktail product

Banin, M.M., Azizah, A., Jusni, Farahdina, R., Candra, K.P., Saragih, B. and Yuliani

Available Online: 29 MAY 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).290](https://doi.org/10.26656/fr.2017.6(3).290)

Banin *et al.* developed and evaluated the quality of jelly coated tapai as a cocktail product.

Conversion of left-over ice cream into bakery product for food sustainability

Jamaludin, N.S., Baharuddin, A.S., Karim, S., Wakisaka, M. and Rahman, N.A.A.

Available Online: 5 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).410](https://doi.org/10.26656/fr.2017.6(3).410)

Jamaludin *et al.* studied on the conversion of left-over ice cream into bakery product for food sustainability.

Simultaneous analysis of patin fish oil (*Pangasius micronemus*) and bandeng (*Chanos chanos*) fish oil using FTIR spectroscopy and chemometrics

Ikhsan, A.N., Irnawati, I. and Rohman, A.

Available Online: 5 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).353](https://doi.org/10.26656/fr.2017.6(3).353)

Ikhsan *et al.* performed a simultaneous analysis of patin fish oil (*Pangasius micronemus*) and bandeng (*Chanos chanos*) fish oil using FTIR spectroscopy and chemometrics.

Organoleptic acceptability and nutritional evaluation of innovative *Moringa oleifera* leaves-based herbal teas incorporated various aromatic herbs

Barakat, H.

Available Online: 5 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).411](https://doi.org/10.26656/fr.2017.6(3).411)

Barakat studied the organoleptic acceptability and nutritional evaluation of innovative *Moringa oleifera* leaves-based herbal teas incorporated various aromatic herbs.

Optimization of roasting temperature and time of the durian seed (*Durio zibethinus* L.) as coffee substitution and its flavour profile

Natania, K. and Wijaya, E.

Available Online: 5 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).413](https://doi.org/10.26656/fr.2017.6(3).413)

Natania and Wijaya optimized of roasting temperature and time of the durian seed (*Durio zibenthinus* L.) as coffee substitution and its flavour profile.

Development of pastilles from flesh and rind of watermelon

Din, S.N., Mubarak, A., Lani, M.N., Yahaya M.Z. and Wan Abdullah, W.Z.

Available Online: 12 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).248](https://doi.org/10.26656/fr.2017.6(3).248)

Din *et al.* developed pastilles from flesh and rind of watermelon

The effect of konjac glucomannan and Aloe vera gel concentration on physical and mechanical properties of edible film

Warkoyo, Purnomo, I., Siskawardani, D.D. and Husna, A.

Available Online: 12 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).415](https://doi.org/10.26656/fr.2017.6(3).415)

Warkoyo *et al.* evaluated the effect of konjac glucomannan and Aloe vera gel concentration on physical and mechanical properties of edible film.

The effects of adding lysine to sap on chemical characteristics and antioxidant activity of granulated coconut sugar

Haryanti, P. and Sulisty, S.B.

Available Online: 12 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).420](https://doi.org/10.26656/fr.2017.6(3).420)

The effects of adding lysine to sap on chemical characteristics and antioxidant activity of granulated coconut sugar were studied by Haryanti and Sulisty.

Siam orange (*Citrus nobilis* L.) nectar characteristics with variations in stabilizer and sucrose level

Aini, N., Dwiyan, H., Setyawati, R., Handayani, I., Septiana, A.T., Sustrawan, B. and Aena, D.A.Q.

Available Online: 16 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).386](https://doi.org/10.26656/fr.2017.6(3).386)

Aini *et al.* studied on the Siam orange (*Citrus nobilis* L.) nectar characteristics with variations in stabilizer and sucrose level.

Chemical and microbiological analysis of fermented probiotic watermelon juice

Lani, M.N, MohdMangsor, N.H., Sharifudin, S.A., Abdullah, W.Z.W., MohdIsa, N.S., Jamzuri, M.N.S. and MohdMaidin, N.

Available Online: 16 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).402](https://doi.org/10.26656/fr.2017.6(3).402)

The chemical and microbiological analysis of fermented probiotic watermelon juice were studied by Lani *et al.*

Phytochemical content and antioxidant activity of Komba-komba (*Eupatorium odoratum* L)

Sabarudin, Nuralifah, Zubaydah, W.O.S., Sahumena, M.H., Sari, F.N., Nelisa and Yamin

Available Online: 16 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).495](https://doi.org/10.26656/fr.2017.6(3).495)

The phytochemical content and antioxidant activity of Komba-komba (*Eupatorium odoratum* L) were evaluated by Sabarudin *et al.*

Effect of operational conditions on physicochemical profiles of spray-dried powder of mango (*Mangifera indica* L.) juice from Tu Quy variety in Vietnam

Pham, T.V., Nguyen, M.T.P., Do, L.V., Truong, M.N., Vo, A.N., Van, K.C. and Le, T.D.

Available Online: 16 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).319](https://doi.org/10.26656/fr.2017.6(3).319)

Pham *et al.* studied on the effect of operational conditions on physicochemical profiles of spray-dried powder of mango (*Mangifera indica* L.) juice from Tu Quy variety in Vietnam.

The characterization of Enterobacteriaceae and Pseudomonadaceae isolated from natural salt licks in Sarawak Borneo

Lihan, S., Jalin, F.J.E., Mohd-Azlan, J., Chiew, S.T. and Chai, L.C.

Available Online: 19 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).208](https://doi.org/10.26656/fr.2017.6(3).208)

Lihan *et al.* characterized Enterobacteriaceae and Pseudomonadaceae isolated from natural salt licks in Sarawak Borneo.

Cryogenic freezing preserves the quality of whole durian fruit for the export market

Razali, N.A., Wan Ibrahim, W.M., Safari, S., Rosly, N.K., Hamzah, F.A. and Wan Husin, W.M.R.

Available Online: 19 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).428](https://doi.org/10.26656/fr.2017.6(3).428)

Razali *et al.* studied on the quality of cryogenic freezing of whole durian fruit for the export market.

Effect of jambu mawar [*Syzygium jambos* (L.) Alston] leaves extract on natural microbial populations in food

Ali, S.K., Son, R., Nor Khaizura, M.A.R. and Rukayadi, Y.

Available Online: 19 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).446](https://doi.org/10.26656/fr.2017.6(3).446)

The effect of jambu mawar [*Syzygium jambos* (L.) Alston] leaves extract on the natural microbial populations in food was evaluated by Ali *et al.*

Effects of extraction methods on antioxidants and methoxyflavones of *Kaempferia parviflora*

Chaisuwan, V., Dajanta, K. and Srikaeo, K.

Available Online: 19 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).408](https://doi.org/10.26656/fr.2017.6(3).408)

The effects of extraction methods on the antioxidants and methoxyflavones of *Kaempferia parviflora* were studied by Chaisuwan *et al.*

The effect of red guava (*Psidium guajava* L.) juice on pregnant women's hemoglobin level

Olii, N., Sukaisi, Asriah, Kusika, S.Y., Situmorang, C.C., Haumahu, C.M., Tompunuh, M.M. and Zuraidah

Available Online: 26 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).435](https://doi.org/10.26656/fr.2017.6(3).435)

The effect of red guava (*Psidium guajava* L.) juice on pregnant women's haemoglobin levels was studied by Olii *et al.*

Effect of conventional and ultrasonic-assisted extracts on betacyanin content of red dragon fruit (*Hylocereus polyrhizus*)

Thuy, N.M., Ngoc, P.T.B. and Tai, N.V.

Available Online: 26 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).754](https://doi.org/10.26656/fr.2017.6(3).754)

The effect of conventional and ultrasonic-assisted extracts on betacyanin content of red dragon fruit (*Hylocereus polyrhizus*) was studied by Thuy *et al.*

Replacing a part of wheat flour with starchy food containing high levels of resistant starch in noodles processing

Vuong, K.M., Tram, N.B., Tuyen, L.N., Vy, L.T.T., Tai, N.V. and Thuy, N.M.

Available Online: 26 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).1020](https://doi.org/10.26656/fr.2017.6(3).1020)

Vuong *et al.* evaluated the effects of replacing a part of wheat flour with starchy food containing high levels of resistant starch in noodles processing.

Antioxidant, anti-tyrosinase, and anti-angiogenic activities of dragon fruit (*Hylocereus* spp.)

Cruz, M.M., Reyes, S.B., Angeles, H.G., Del Rosario, J.M., Lirazan, M.B., Estacio, R.C., Corales, L.M. and Dalmacio, L.M.

Available Online: 26 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).400](https://doi.org/10.26656/fr.2017.6(3).400)

Cruz *et al.* studied on the antioxidant, anti-tyrosinase, and anti-angiogenic activities of dragon fruit (*Hylocereus* spp.)

Comparative study of nutritional and functional characteristics of pearl millet, buckwheat, amaranth and unripe banana flours for gluten-free bakery products

Rustagi, S., Khan, S. and Jain, T.

Available Online: 30 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).624](https://doi.org/10.26656/fr.2017.6(3).624)

Rustagi *et al.* performed a comparative study on the nutritional and functional characteristics of pearl millet, buckwheat, amaranth and unripe banana flours for gluten-free bakery products.

Optimization of enzymatic hydrolysis of boso fish (*Oxyeleotris marmorata*) protein based on the degree of hydrolysis and the physical properties of the resultant hydrolysates

Priatni, S., Ratnaningrum, D., Kosasih, W., Eriska, H., Devi, A.F. and Budiari, S.

Available Online: 30 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).593](https://doi.org/10.26656/fr.2017.6(3).593)

Priatni *et al.* optimized the enzymatic hydrolysis of boso fish (*Oxyeleotris marmorata*) protein based on the degree of hydrolysis and the physical properties of the resultant hydrolysates.

Immunonutrition as a potential strategy to prevent and cope with coronavirus disease (COVID-19)

Acevedo-Espinola, R. and Torres-Obregón, S.E.B.

Available Online: 12 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).346](https://doi.org/10.26656/fr.2017.6(3).346)

Acevedo-Espinoala and Torres-Obregón studied on immunonutrition as a potential strategy for COVID-19 prevention.

Analysis of amino acids in food using High Performance Liquid Chromatography with derivatization techniques: a review

Lestari, L.A., Rohman, A., Riswahyuli, Purwaningsih, S., Kurniawati, F. and Irnawati

Available Online: 30 JUNE 2022 | [https://doi.org/10.26656/fr.2017.6\(3\).442](https://doi.org/10.26656/fr.2017.6(3).442)

Lestari *et al.* reviewed on the analysis of amino acids in food using High Performance Liquid Chromatography (HPLC) with derivatization techniques.

VOL. 6 | ISSUE 2

VOL. 6 | ISSUE 4



[VIEW ARTICLES](#)

[ISSUES](#)

[ABOUT](#)

[FOR AUTHORS](#)

[FOR REFEREES](#)

EDITORIAL BOARD

Chief Editor

Son Radu

Malaysia

Editor

Vivian New Chia Yeung

Malaysia

Assistant Editor

Kimberley Rinai

Malaysia

Associate Editors

Prem Narayan Paudel

Kathmandu University

Nepal

Associate Editors

Borarin Buntong

Royal University of Agriculture

Cambodia

Associate Editors

Yoshitsugu Nakaguchi

Ishikawa Prefectural University

Japan

Nasreldin Elhadi

University of Dammam

Saudi Arabia

Dwaipayan Sikdar

University of Chittagong

Bangladesh

Rungsinee Sothornvit

Kasetsart University,

Thailand

Nikos Tzortzakis

Cyprus University of Technology
Cyprus

Pin Kar Yong

Forest Research Institute Malaysia
(FRIM)
Malaysia

Chim Chay

University of Agriculture
Cambodia

Le Van Viet Man

Ho Chi Minh City University of
Technology
Viet Nam

Oscar Herrera-Calderon

Universidad Nacional Mayor de
San Marcos
Peru

Hasmadi Mamat

University Malaysia Sabah
Malaysia

Regiane Ribeiro dos Santos

Federal Rural University of Rio de
Janeiro
Brazil

Koh Soo Peng

Malaysian Agricultural Research
and Development Institute
(MARDI)

Saisamorn Lumyong

Chiang Mai University
Thailand

Hanifah Nuryani Lioe

Bogor Agricultural University
Indonesia

Michelle Galindo de Oliveira

Federal University of Pernambuco
Brazil

Alper Sener

Onzekiz Mart University
Turkey

Angelo Maria Giuffrè

Mediterranea University of Reggio
Calabria
Italy

Javier Garcia-Lomillo

University of Burgos
Spain

Laura Marcela Zárate Polanco

National University of Colombia-
Bogotá
Colombia

Swapan K. Banerjee

HPFB Health Canada,
Canada

Adel Abu Bakr Abd El-Hamid**Ekachai Chukeatirote**

Mae Fah Luang University
Thailand

Saroat Rawdkuen

Mae Fah Luang University
Thailand

Giovanna Giacalone

Università di Torino
Italy

Lorina Acilo Galvez

Visayas State University
Philippines

Sergio Soto Simental

Universidad Autónoma del Estado
de Hidalgo
Mexico

Fernando Cebola Lidon

Universidade Nova de Lisboa
Portugal

Linda Lim Biaw Leng

Universiti Brunei Darussalam
Brunei Darussalam

J.M. Krishanti Jayarukshi Kum**Premarathne**

Wayamba University of Sri Lanka
Sri Lanka

Katherine Ann T. Castillo-Isra

University of the Philippines L

Malaysia

Shatta

Banos

Suez Canal University

Philippines

Egypt

Dinko Hristov Dinko

Trakia University

Bulgaria

Iddya Karunsagar

Nitte University

India

Steve Flint

Massey University

New Zealand

Giuseppe Sortino

University of Palermo

Italy

COPYRIGHT © 2016 - 2023 RYNNYE LYAN RESOURCES

The effect of rotary drying temperature on drying characteristic and antioxidant activity of *Etilingera elatior* Jack

¹Simanjuntak, M.E., ²Ristiarini, S. and ^{2,*}Widyawati, P.S.

¹Mechanical Engineering Department, Medan State Polytechnic, Medan, Indonesia

²Food Technology Department, Widya Mandala Surabaya Catholic University, Surabaya, Indonesia

Article history:

Received: 10 June 2021

Received in revised form: 10 July 2021

Accepted: 18 October 2021

Available Online: 22 May 2022

Keywords:

Etilingera,
Rotary drying,
Drying characteristics,
Antioxidant

DOI:

[https://doi.org/10.26656/fr.2017.6\(3\).333](https://doi.org/10.26656/fr.2017.6(3).333)

Abstract

The study was conducted to investigate the effect of drying temperature on a rotary type of dryer for drying characteristics and antioxidant activity of *Etilingera elatior* flower. This study used a lab-scale rotary dryer which was a modification of the commercial oven heater from the market. The research design was a Randomized Block Design (RCBD) with one factor, i.e. the drying temperature, which consisted of four temperature levels, 60°C, 70°C, 80°C and 90°C. All experiments were repeated six times. The results showed that the drying of *Etilingera elatior* flowers at three different temperatures (70°C, 80°C and 90°C) required a shorter drying time than that of 60°C. The drying rate of the samples at 70°C, 80°C and 90°C was drastically decreased before the 200th min, while 60°C took a long time to 380 mins. The drying rate pattern with the drying temperature of 60°C showed a significantly lower total phenolic content of *Etilingera* flowers compared to 70°C, 80°C and 90°C, while there was no significant difference in total phenolic content among 70°C, 80°C and 90°C. In addition, different drying temperatures did not give a significant effect on the antioxidant activity based on the DPPH (2,2-diphenyl-1-picrylhydrazyl radical) method. This study proposed the effectiveness of drying using a rotary dryer in maintaining the total phenolic content and antioxidant activity of *Etilingera* flowers.

1. Introduction

Etilingera elatior Jack (ginger red/torch ginger) is a spice plant that is included in the *Zingiberaceae* family that is quite widespread in Indonesia. This herb plant can be found in several regions in Southeast Asia (Wijekoon *et al.*, 2010; Nor *et al.*, 2020) with different names, such as *Kincung* in Medan, *Rias* in North Tapanuli, *Sambuung* in Minangkabau, *Kecicang* in Bali, *Siantan* in Malaya and *Daalaa* in Thailand (Health Research and Development Agency of Indonesia, 2000; Lachumy *et al.*, 2010; Nor *et al.*, 2020). *Etilingera elatior* flower is potential as a food flavouring (seasoning) in cooked, sauteed, or heated (Noweg *et al.*, 2003; Juwita *et al.*, 2018) as in fish processing (*pepes* fish, grilled fish, and fried). Fish cooked with *Etilingera* will taste better and the fishy aroma will be reduced (Sukandar *et al.*, 2011). In addition, this plant is efficacious in deodorizing body odour and bad breath (Health Research and Development Agency of Indonesia, 2000; Aldi *et al.*, 2020), natural cosmetic ingredients (Chan *et al.*, 2007). The profile of *E. elatior* Jack is shown in Figure 1a.




Figure 1. *Etilingera elatior* Jack (a) and Rotary dryer lab-scale (b)

The benefits offered by *Etilingera* flower are correlated to phytochemical compounds, such as alkaloids, saponins, tannins, phenolics, flavonoids, triterpenoids, steroids, and glycosides (Naufalin *et al.*, 2005; Setiawati, 2018). A previous study showed that flavonoids, terpenoids, saponins, tannins, and carbohydrate compounds from *etilingera* flowers can be extracted using 80% methanol as the solvent (Lachumy *et al.*, 2010). Furthermore, *E. elatior* has been described to exhibit many biological activities (Lachumy *et al.*, 2010; Chan *et al.*, 2009; Puttarak *et al.*, 2014; Juwita *et al.*, 2018; Nor *et al.*, 2020; Putri, 2021).

*Corresponding author.

Email: paini@ukwms.ac.id

Several investigations revealed that the phytochemical compounds of the *Etlingera* flowers have the potential of becoming a source of antioxidants. The methanolic and ethanolic extracts of them have been proven to scavenge DPPH (2,2-diphenyl-1-picrylhydrazyl) free radical (Chan *et al.*, 2007). The inhibition of DPPH radical was successfully performed by *Etlingera* flowers' methanolic extract at the concentration of 9.14 mg/mL (Lachumy *et al.*, 2010) as well as its aqueous extracts at the concentration of 76.4% (Gasemzadeh *et al.*, 2015). Naufalin and Rukmini (2011) also found that the *Etlingera* flower has a higher potential antioxidant activity (61.61-83.17%) than its leaf (40.64-60.40%) and stem (57.42-84.65%).

In general, fresh *etlingera* flower usually has moisture contents of more than 80%, so it is a very perishable commodity (Sagar and Kumar, 2010). The shelf life of *Etlingera* flower can be extended by reducing its moisture content using drying methods. Several drying methods, such as air-, freeze-, microwave -, vacuum-, oven- and sun-drying, have different effects on the quality of dried material. Variation of drying temperature using a specific drying method can also influence the losses or preservation of antioxidant compounds (Kamiloglu *et al.*, 2016). Can *et al.* (2009) studied the effect of various drying types, such as oven drying, microwave, oven, sun drying, and freeze-drying on the antioxidant properties of *E. elatior* leaves. The total phenolic content of *E. elatior* leaves subjected to freeze-drying was found to be higher compared to other drying methods. Another study by Rohkyani and Suryani (2015) showed drying using an oven at 65°C resulted in the highest DPPH scavenging activity (66.43%) of *Etlingera* flowers compared to 85°C (56.76%).

Rotary drying is another drying method that has not been widely applied in the drying of food commodities. Delele *et al.* (2015) informed that rotary dryers are capable of processing a variety of agricultural products with a wide range of thermo-physical and flow properties, and improve the efficiency of the drying process. To the best of our knowledge, there is still few information regarding the antioxidant characteristics of *E. elatior* flowers dried using a rotary dryer.

Several studies utilized the rotary drying method to preserve food material. Kaleemullah (2005) used rotary drying to 10.5 kg chilli with an initial moisture content of 330% (dry basis) in the temperature range of 50 – 65°C. The study showed that the moisture content is reduced to 10.0% (dry basis) after being dried for 32 hrs, 27 hrs, 23 hrs, and 20 hrs at 50°C, 55°C, 60°C and 65°C, respectively. Tarhana (2010) examined the drying of 15 kg peppermint using a rotary dryer for 15 - 18 hrs and 12

– 15 hrs. Drying can cause the leaves to darken, but the essential oil content is not much different from 2.08 - 2.7 mL/100 g dry matter. The specific energy consumption values have ranged from 7.88 to 15.08 MJ/kg of moisture removed). Daily fluctuations in ambient air conditions directly affected the specific energy consumption of a rotary dryer. Ademiluyi *et al.* (2010) also uses a rotary dryer to dry fermented cassava with parameter tests, including dry air inlet temperature, dry air inlet velocity, relative humidity, feed rate, drum rotational per min and feeding. The results showed that the dry air inlet temperature, the dry air inlet velocity, and the feed rate give a significant effect on the specific heat transfer coefficient and heat load on the material.

This study aimed to investigate the effect of rotary drying temperature on drying characteristics and antioxidant activity of *E. elatior* flowers that including moisture content, drying rate, total phenol, and DPPH free radical scavenging activity. Antioxidant activity analysis was carried out to determine the potential antioxidant activity of *Etlingera* flowers dried under different drying temperatures (60°C, 70°C, 80°C and 90°C) using a rotary dryer. The antioxidant activity was obtained by comparing the antioxidant activity of water extract of *Etlingera* flower petal powder at various drying temperatures with gallic acid and calculating its ability to reduce DPPH free radicals based on the percentage of inhibition (% inhibition). This study used a lab-scale rotary dryer made from a modified commercial oven heater.

2. Materials and methods

Fresh *E. elatior* flowers were obtained from the traditional market around Medan city, Indonesia. All chemicals used were of analytical grade. The DPPH (2,2-diphenyl-1-picrylhydrazyl free radical) was purchased from Sigma Aldrich Chemicals (St. Louis, Missouri, United States), gallic acid, Folin-ciocalteu phenol, sodium carbonate, methanol was obtained from Merck and Co. Pharmaceutical Company (New Jersey, United States), and aquadest was bought from PT. Surabaya Aqua Industry, Indonesia.

The petals of fresh *E. elatior* flowers were collected and the knobs were discarded. The fresh flower petals were chopped to a size of about 1 cm. For each experiment, around 400±0.1 g (Ohaus PA 224) of fresh petals was inserted into the drying basket and filled about 50% volume of the basket. Commercial electric oven (Oxone type OX-8830, 30 L in volume) with modification was pre-heated to the set temperature until a steady condition was achieved, followed by inserting the drying basket containing samples into the oven. The drying air temperatures used were 60°C, 70°C, 80°C,

and 90°C while the basket rotates at around 3 rpm. The experimental setup was shown in Figure 1b. The experimental setup is shown in Figure 1b. During the drying process, the samples were taken out, immediately weighed, and put back into the oven. The process was done periodically every 20 min until the samples reached a constant weight and the drying process was terminated. The moisture content was calculated based on a wet basis with Equation 1:

$$MC (\%) = \frac{M_{H_2O}}{M_{total}} \times 100\% \quad (1)$$

Where MC = moisture content (%), M_{H_2O} = mass of moisture (g) and M_{total} = mass of solid + mass of moisture (g)

The drying rate was calculated from the Equation 2:

$$DR (\%/min) = \frac{MC_{t_0} - MC_{t_1}}{\Delta t} \quad (2)$$

Where DR = drying rate (%/min), MC_{t_0} = moisture content at time t (g), MC_{t_1} = moisture content at t + 20 min (g) and Δt = time interval (min).

The two transfer processes simultaneously occur in thermal drying: i.e. heat or energy transfer and mass transfer. Energy transfer occurs from the environment to vaporize moisture present on the surface of a material/product. Convection heat transfer takes place due to temperature differences between the surface of the material and the dry air. Vaporization is the evaporation of moisture located on the surface of the material. When heat moves from the outside to the inside of the material, moisture that is located inside the material will diffuse out to the material's surface due to differences in moisture content.

2.1 Extraction of samples

The samples were extracted by the soxhlet extraction method (Widyawati et al., 2014). As much as 3 g of fresh *E. elatior* petals were wrapped in a filter paper and put into a soxhlet tube (timbre) that had been filled with 50 mL of water as the solvent. The extraction was carried out at its boiling point for 4 hrs until the solvent's colour in the timbre became colourless. The obtained extract was evaporated by a rotary evaporator under a vacuum at a temperature of 65-80°C, a pressure of 250 - 300 mbar, and a rotational speed of 40 rpm. The evaporation was run for 10 mins to obtain 2.5 mL of a concentrated extract. The concentrated extract was placed in vials and stored in the freezer until further analysis.

2.2 Total phenolic content analysis

Analysis of total phenolic content was based on the

oxidation reaction of phenol compounds to produce a blue solution from reducing yellow hetero polyphosphoric molybdate tungstate anions (Muntana and Prasong, 2010). In brief, 20 μ L of *Etilingera* flower extract was added to 1 mL of 10% Folin Ciocalteu reagent in a 10 mL flask bottle. The mixture was shaken and left for 5 mins. Then, 2.0 mL of 7.5% Na_2CO_3 and distilled water were subsequently added to the mixture until a volume of 10 mL was achieved. The mixture was allowed to stand for 30 mins and the absorbance of the samples was measured at λ 760 nm (UV-Vis spectrophotometer, Shimadzu 1800). Data were analysed against gallic acid as a standard solution (mg GAE/g samples) (Siddiqui et al., 2017).

2.3. DPPH free radical scavenging activity analysis

The assay of DPPH free radical scavenging activity was based on Sompong et al. (2011) with some modifications. As much as 3 mL of DPPH solution (4 mg/100 mL in methanol) was added to 20 μ L of the extract and methanol until the volume reached 10 mL in a 10 mL flask bottle. The samples were incubated for 30 mins in a dark chamber. The absorbance of each sample was measured at λ 517 nm by a spectrophotometer (UV-Vis spectrophotometer, Shimadzu 1800). Data were analysed against gallic acid as a standard solution (mg GAE/g samples).

3. Results and discussion

3.1 Moisture content and drying rate

The initial moisture content of each sample for temperatures 60°C, 70°C, 80°C, and 90°C was 87.53%, 88.03%, 88.1%, and 87.53% (w/w). During the drying process, the changes in moisture content occurred as shown in Figure 3 and these changes in moisture content are proportional. As for the temperature of 60°C, the change is much slower. For each drying temperature of 90°C, 80°C, 70°C, and 60°C, the moisture content will approach 0% after 240, 260, 320, and 480 mins, respectively. The decrease in moisture content follows a straight line until the moisture content approaches 0%. The decrease indicates that the process of moisture evaporation takes place constantly for each period of measurement. The moisture content for each experiment calculated based on Equation 1 is shown in Figure 2.

The drying rate during the experiment calculated using Equation 2 is shown in Figure 3. The constant drying rate looks long enough for all types of temperatures. The constant drying period of around 220 mins was nearly the same for drying temperatures of 70°C, 80°C, and 90°C. During this period, the drying rate is around 1.25 – 30 g/min of measurement. At 60°C, the drying rate is around 0.75 g/min and lasts for about 380

mins. The duration of this constant rate period is due to the high moisture content contained in fresh *E. elatior* flowers. Moisture content for all samples was around 90% and a large amount of moisture content is present on the surface of the sample. The low thickness of the sample also allows the moisture on the inside or middle to evaporate to the surface with minimum difficulty. The fall rate period was shorter than the constant rate period, which was around 40 to 100 mins for experiments with temperatures of 90°C, 80°C, and 70°C. As for the drying temperature of 60°C, the time was around 240 mins.




Figure 2. The moisture content of *Etilingera elatior* Jack flowers at various drying temperatures.




Figure 3. The drying rate of *Etilingera elatior* Jack flowers at various drying temperatures.

The second falling rate period has a shorter time than the previous period. A shorter time indicates that after experiencing a critical condition, the moisture content of *E. elatior* flowers will then be low. During the second falling rate period, the drying rate would be controlled by the diffusion rate of moisture from the inner surface of the sample. This process was conducted until it reached an equilibrium state or until the moisture content has completely evaporated (Nurafifah *et al.*, 2018). Data in Figure 3 showed a constant drying rate lasting up to 20% of moisture content. After that, the drying rate will decrease rapidly. Moisture content in this range is a critical condition where the drying rate will be significantly influenced by the rate of moisture transportation from the centre to the surface.

3.2 Antioxidant activity

The total phenolic content describes the amount of phenol compounds present in a sample. Phenol compounds can facilitate redox reactions, thus can act as antioxidants (Johari and Khong, 2019). The total phenolic content of *Etilingera* flower powder produced using various temperatures of rotary dryer is shown in Figure 4. The results showed total phenolic content of *Etilingera* flower powder produced under the temperature of 60°C (1.216 ± 0.146 mg gallic acid equivalent/g samples) was statistically lower from that of 70°C, 80°C, and 90°C (2.558 ± 0.385 - 2.165 ± 0.609 mg gallic acid equivalent/g samples). The trend of increasing total phenolic content was observed when the drying temperature increased to 70°C, followed by a decrease after the heating temperature was elevated to 90°C. However, no significant difference in total phenolic content was found from the drying temperatures of 70°C, 80°C, 90°C. The obtained results revealed that the phenolic compounds in *E. elatior* flowers were thermostable and could be maximally extracted under the drying temperature of 70°C. Złotek *et al.* (2019) also discovered the thermostability of phenolic compounds derived from white quinoa after being subjected to drying using a single convection dryer under various drying temperatures (30°C, 40°C, 60°C). Generally, the secondary metabolites contained in plants are in the form of free metabolites or bonded to other metabolites. Vanic acid possesses a thermostable characteristic at various dry temperatures, which is similar to p-hydroxybenzoic acid, p-coumaric acid, salicylic acid, and ferulic acid. Naufalin *et al.* (2005) reported *etlingera* flowers contained phytochemical components, such as alkaloids, saponins, tannins, phenolics, flavonoids, triterpenoids, steroids, and glycosides. In contrast, Liaotrakoon and Liaotrakoon (2018) observed a decrease in the total phenolic content and antioxidant activity of mushrooms after being dried using an oven under the temperatures of 40°C, 50°C, and 60°C with an extended drying time.




Figure 4. Total phenolics content of *Etilingera elatior* Jack flowers at various drying temperatures. Values are presented as means of six replicates. Values with different superscripts are significantly different, $p < 0.05$.

Therefore, it can be concluded that drying using the rotary dryer with a higher temperature and longer time up to 4 hrs could maintain the total phenolic content of a material.

DPPH is a free radical that has a maximum absorbance at 517 nm in methanol. The scavenging of DPPH by the addition of antioxidants is indicated by the change in solvent's colour from purple to yellow (Sayed *et al.*, 2015). The ability of *Etlingera* flowers to scavenge DPPH free radicals tended to increase along with increasing drying temperature (Figure 5). In addition, the drying process of *Etlingera* flowers using the rotary dryer with temperatures ranging from 60-90°C for 4 hrs did not significantly change its DPPH scavenging activity, which indicated the ability of the rotary drying method to effectively maintain both total phenolic content and antioxidant activity of *Etlingera* flowers. On the contrary, different results from Liaotrakoon and Liaotrakoon (2018) revealed a decrease in the antioxidant activity of mushrooms after being subjected to the oven drying process.




Figure 5. DPPH Free Radical scavenging activity of *Etlingera elatior* Jack flowers at various drying temperatures.

Furthermore, no correlation was found between the antioxidant activity and the total phenolic content of *Etlingera* flowers. The coefficient determination of less than 0.95 ($R^2 = 0.3318$) indicated an increase in total phenolic content was not closely related to an increase in the DPPH free radical scavenging activity. This result was not in line with Piluzza and Bullitta (2011) where the DPPH and ABTS free radicals scavenging activity were positively correlated with total phenolic content with $R^2 = 0.9152$ and $R^2 = 0.889$, respectively. Gan *et al.* (2017) also noted total phenolic content was linearly correlated with antioxidant activity (iron ion reducing power, hydroxyl free radical scavenging activity and lipid oxidation inhibition activity). Khiya *et al.* (2021) added a high positive correlation between phenolic compounds and antioxidant activity ($R^2 = 0.932$), showing the phenolic compounds may have contributed to the antioxidant activity of *Salvia officinalis* leaves.

This absence of correlation in this study depicted the possibility of other phytochemical compounds influencing the antioxidant activity of *Etlingera* flowers. Naufalin *et al.* (2005) and Setiawati (2018) showed that the phytochemical compounds that comprise *Etlingera* flowers are alkaloids, glycosides, phenolics, terpenoids, steroids, saponins, and flavonoids, with alkaloids as the dominant compound. Alkaloids and phenolic compounds are very important as antioxidants, with alkaloids possessing a stronger antioxidant activity. Quezada *et al.* (2004) showed the presence of alkaloids and flavonoids exhibited high antioxidant potency of Boldo (*Peumus boldus* Molina) extract. Benabdesselam *et al.* (2007) also found that total quinolizidine alkaloid contents of *Fumaria capreolata* (426 mg/100 g) and *Fumaria bastardii* (521 mg/100 g) extracts exhibited a strong total antioxidant activity.

4. Conclusion

The time required for drying *Etlingera* flowers using a rotary dryer did not much differ among the temperatures of 70°C, 80°C, and 90°C and was shorter compared to 60°C. A rapid decrease in drying rate was experienced by *Etlingera* flowers before the 200th min of drying process under 70°C, 80°C, and 90°C, while longer time (380 mins) occurred when using a drying temperature of 60°C. Drying with a rotary dryer was found to effectively maintain the total phenolic content of *Etlingera* flowers, although the drying process was performed under a quite high temperature for 4 hrs. In addition, drying with rotary drying at 60 – 90°C for 4 hrs did not change the DPPH free radical scavenging activity. Accordingly, the rotary drying method was very effective in maintaining the total phenolic content and antioxidant activity of *Etlingera* flowers.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgement

The authors would like to thank the Medan State Polytechnic for funding this research on Collaboration Research Scheme 2019 year and Widya Mandala Surabaya Catholic University.

References

- Ademiluyi, F.T., Abowei, M.F.N., Puyate, Y.T. and Achinewhu, S.C. (2010). Effects of drying parameters on heat transfer during drying of fermented ground cassava in a rotary dryer. *Drying Technology*, 28(4), 550-561. <https://doi.org/10.1080/07373931003622669>

- Aldi, Y., Husni, E. and Yesika, R. (2020). Activity of kincung flowers (*Etlingera Elatior* (Jack) R.M.Sm.) on total leukocytes and percentage of leukocytes in allergic male white mice. *Pharmacognosy Journal*, 12(1), 44–51. <https://doi.org/10.5530/pj.2020.12.8>.
- Benabdesselam, F.M., Khentache, S., Bougoffa, K., Chibane, M., Adach, S., Chapeleur, Y., Max, H. and Mattar, D.L. (2007). Antioxidant activities of alkaloid extracts of two Algerian species of *Fumaria*: *Fumaria capreolata* and *Fumaria bastardii*. *Records Natural Products*, 1(2-3),28-35.
- Chan, E.W.C, Lim, Y.Y. and Wong, S.K. (2009). Effects of different drying methods on the antioxidant properties of leaves and tea of ginger species. *Food Chemistry*, 113(1), 166-172. <https://doi.org/10.1016/j.foodchem.2008.07.090>
- Chan, E.W.C., Lim. Y.Y. and Omar, M. (2007). Antioxidant and antibacterial activity of leaves of *Etlingera* species (*Zingiberaceae*) in Peninsular Malaysia. *Food Chemistry*, 104(4), 1586-1593. <https://doi.org/10.1016/j.foodchem.2007.03.023>
- Delele, D.A., Weigler, F. and Mellmann. J. (2015). Advances in the application of a rotary dryer for drying of agricultural products: A Review. *Drying Technology*, 33(5), 541-558. <https://doi.org/10.1080/07373937.2014.958498>
- Gan, J., Feng, Y., He, Z., Li, X. and Zhang, H. (2017). Correlations between antioxidant activity and alkaloids and phenols of Maca (*Lepidiummeyerii*). *Hindawi Journal of Food Quality*, 2017, 3185945. <https://doi.org/10.1155/2017/3185945>
- Johari, M.A. and Khong, H.Y. (2019). Total phenolic content and antioxidant and antibacterial activities of *Pereskia Bleo*. *Hindawi Advances in Pharmacological Sciences*, 2019, 7428593. <https://doi.org/10.1155/2019/7428593>
- Juwita, T., Puspitasari, I.M. and Levita, J. (2018). Review article torch ginger (*Etlingera elatior*): A review on its botanical aspects, phytoconstituents and pharmacological activities. *Pakistan Journal of Biological Sciences*, 21(4), 151-165. <https://doi.org/10.3923/pjbs.2018.151.165>
- Kaleemullah, S. and Kailappan, R. (2005). Drying kinetics of red chillies in a rotary dryer. *Biosystems Engineering*, 92(1), 15–23. <https://doi.org/10.1016/j.biosystemseng.2005.05.015>
- Kamiloglu, S., Toydemir, G., Boyacioglu, D., Beekwilder, J., Hall, R.D. and Capanoglu, E. (2016). A Review on the effect of drying on antioxidant potential of fruits and vegetables. *Critical Reviews in Food Science and Nutrition*, 56(Suppl. 1), S110-S129. <https://doi.org/10.1080/10408398.2015.1045969>
- Khiya, Z., Oualcadi, Y., Gamar, A., Berrekhis, F., Zair, T. and Hilali, F.E.L. (2021). Correlation of total polyphenolic content with antioxidant activity of hydromethanolic extract and their fractions of the *salvia officinalis* leaves from different regions of Morocco. *Journal of Chemistry*, 2021, 8585313. <https://doi.org/10.1155/2021/8585313>
- Lachumy, S.J.T., Sasidharan, S., Sumathy, V. and Zuraini, Z. (2010). Pharmacological activity, phytochemical analysis and Toxicity of Methanol Extract of *Etlingera Elatior* (Torch Ginger) Flowers. *Asian Pacific Journal of Tropical Medicine*, 3(10), 769-774. [https://doi.org/10.1016/S1995-7645\(10\)60185-X](https://doi.org/10.1016/S1995-7645(10)60185-X)
- Liaotrakoon, W. and Liotrakoon, V. (2018). Influence of drying process on total phenolics, antioxidative activity and selected physical properties of edible bolete (*Phlebopus colossus* (R. Hein) Singer and changes during storage. *Food Science and Technology*, 38(2), 231 – 237. <https://doi.org/10.1590/1678-457x.34116>
- Muntana, N. and Prasong, S. (2010). The Study on total phenol contents and their antioxidant activities on Thai white, red, black rice bran extracts. *Pakistan Journal of Biological Science*, 13(4), 170 – 174. <https://doi.org/10.3923/pjbs.2010.170.174>
- Naufalin, R. and Rukmini. H.S. (2011). Potential antioxidants from kecombrang (*Nicolaia speciosa* Horan) plant extraction during storage, presented at the National Seminar on Building Competitiveness of Food Products Based on Local Raw Materials. Surakarta, Indonesia. [In Bahasa Indonesia].
- Naufalin, R., Jenie, B.S.L., Kusnandar, F., Sudarwanto, M. and Rukmini, H. (2005). Antibacterial activity of kecombrang extract toward pathogenic and food spoilage bacteria. *Journal of Technology and Food Industry*, 16(2), 119-125. [In Bahasa Indonesia].
- Nor, N.A.M., Noordin, L., Bakar, N.H.A. and Ahmad, W.A.N.W. (2020). Evaluation of antidiabetic activities of *Etlingera elatior* flower aqueous extracts *in vitro* and *in vivo*. *Journal of Applied Pharmaceutical Science*, 10(8), 43-51.
- Noweg, T., Abdullah, A.R. and Nidang, D. (2013). Forest plants as vegetables for communities bordering the Crocker Range National Park. *ASEAN Revolution Biodiversity Environment Conservation*, Jan-Mar 2003, 1–18.
- Piluzza, G. and Bullitta, S. (2011). Correlations between phenolic content and antioxidant properties in twenty-four plant species of traditional ethnoveterinary use in the Mediterranean area. *Pharmaceutical Biology*,

- 49(3), 240-247. <https://doi.org/10.3109/13880209.2010.501083>
- Putri, H.P. (2021). *Etilingera elatior* as an antihyperglycemic in diabetes mellitus. *Jurnal Penelitian Perawat Profesional*, 3(1),189-198. <https://doi.org/10.37287/jppp.v3i1.386> [In Bahasa Indonesia].
- Quezada, N., Asencio, M., Del Valle, J.M., Aguilera, J.M. and Gómez, B. (2004). Antioxidant activity of crude extract, alkaloid fraction, and flavonoid fraction from Boldo (*Peumus boldus* Molina) Leaves. *Journal of Food Science*, 69(5), 371-376. <https://doi.org/10.1111/j.1365-2621.2004.tb10700.x>
- Research and Development Agency Ministry of Health of the Republic of Indonesia (2000). Inventory of Indonesian Medicinal Plants (I). 1st ed. Jakarta, Indonesia: Research and Development Agency Ministry of Health of the Republic of Indonesia. [In Bahasa Indonesia].
- Rohkyani, I. and Suryani, T. (2015). Antioxidant activity and organoleptic test of tea bag stem and flower of kecombrang with variations temperature drying. Surakarta, Indonesia: Muhammadiyah University of Surakarta, BSc. Thesis. [In Bahasa Indonesia].
- Siddiqui, N., Rauf, A., Latif, A. and Mahmood, Z. (2017). Spectrophotometric determination of the total phenolic content, spectral and fluorescence study of the herbal Unani drug Gul-e-Zoofa (*Nepeta bracteata* Benth). *Journal of Taibah University Medical Sciences*, 12(4), 360-363. <https://doi.org/10.1016/j.jtumed.2016.11.006>
- Sompong, R., Kandasamy, I., Khumalo, N. and Baijnath, H. (2011). Physicochemical and antioxidative properties of red and black rice varieties from Thailand, China, and Sri Lanka. *Food Chemistry*, 124(1), 132-140. <https://doi.org/10.1016/j.foodchem.2010.05.115>
- Sukandar, D., Radiastutu, N., Jayanegara, I., Muawanah, A.H.A. and Hudaya, A. (2011). Uji Aktivitas Antioksidan Ekstrak Kasar Air Bunga Kecombrang (*Etilingera elatior*) sebagai bahan pangan fungsional. *Indonesian Journal of Applied Chemistry*, 13(1), 1-4. <https://doi.org/10.14203/jkti.v13i1.124>. [In Bahasa Indonesia].
- Tarhana, S., Telci, I., Taner, M.T. and Polatci, H. (2010). Product quality and energy consumption when drying peppermint by rotary drum dryer. *Industrial Crops and Products*, 32(3), 420-427. <https://doi.org/10.1016/j.indcrop.2010.06.003>
- Widyawati, P.S., Budianta, T.D.W., Kusuma, F.A. and Wijaya, E.L. (2014). The difference of solvent polarity to phytochemical content and antioxidant activity of *Pluchea indica* Less leaves extracts, *International Journal of Pharmacognosy and Phytochemical Research*, 6(4), 850-855.
- Wijekoon, J.O., Karim, M.M. and Bhat, R. (2011). Evaluation of nutritional quality of torch ginger (*Etilingera elatior* Jack) inflorescence. *International Food Research Journal*, 18(4), 1415-1420.
- Złotek, U., Gawlik-Dziki, Dziki, D., Swieca, M., Nowak, R. and Martinez, E. (2019). Influence of drying temperature on the phenolic acid composition and antioxidant activity of sprouts and leaves of white and red quinoa. *Hindawi Journal of Chemistry*, 2019, 7125169. <https://doi.org/10.1155/2019/7125169>



food research kecombrang



- 99+ Compose
- Mail
 - Inbox 1,441
 - Starred
 - Snoozed
 - Sent
 - Drafts 2
 - More
- Chat
- Spaces
- Meet

Labels

FR-2021-333 - Article Production External Inbox x



Food Research <foodresearch.my@outlook.com>
to me

Mon, May 9, 2022

Dear Dr Widyawati,

Manuscript ID: FR-2021-333

Manuscript Title: The effect of rotary drying temperature on drying characteristic and antioxidant activity of *Eti*

Before we can proceed with the article production, I would like to clarify a few points that I have commented in refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once you the copy to me as soon as possible. Please note the faster you respond, the quicker we will process your manu

Thanks & Regards,
Vivian New
Editor
Food Research



food research kecombrang



- 99+ Compose
- Mail
- Inbox 1,441
- Starred
- Snoozed
- Sent
- Drafts 2
- More

Labels

From: Paini Sri Widyawati <paini@ukwms.ac.id>
Sent: Wednesday, 23 February, 2022 11:24 AM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: Give me Information

Dear Dr. Son Radu

Thanks for information

The best Regards

Paini Sri Widyawati

On Wed, Feb 23, 2022 at 1:54 AM Food Research <foodresearch.my@outlook.com> wrote:

Dear Paini Sri Widyawati

It is under technical review. Please expect some delay as we are experiencing a high volume of publication at
Thank you for your understanding.

Best regards,
Son Radu, PhD
Chief Editor



food research kecombrang



- 99+ Compose
- Mail
- Inbox 1,441
- Starred
- Snoozed
- Sent
- Drafts 2
- More

Labels

From: Painsi Sri Widyawati <paini@ukwms.ac.id>
Sent: Tuesday, 22 February, 2022 5:45 PM
To: Food Research <foodresearch.my@outlook.com>
Subject: Give me Information

Dear Editor of Food Research

Related to my manuscript with title " The effect of temperature on drying characteristic and antioxidant activities of *Eth* using a rotary dryer" with Manuscript ID: FR-2021-333. Please give me information about status this manuscript.

Thanks for information

The best regards

Painsi Sri Widyawati

One attachment • Scanned by Gmail





food research kecombrang



99+

Compose

Mail

Inbox 1,441

Chat

Starred

Spaces

Snoozed

Sent

Meet

Drafts 2

More

Labels



Paini Sri Widyawati <paini@ukwms.ac.id>
to Food

May 10, 2022,

Dear Dr. Vivian New

I send again my manuscript that has been revised

Thanks for attention

The Best Regards

Paini Sri Widyawati

One attachment • Scanned by Gmail





food research kecombrang



- 99+ Compose
- Mail
- Inbox 1,441
- Starred
- Snoozed
- Sent
- Drafts 2
- Meet
- More

Labels

One attachment • Scanned by Gmail



Food Research <foodresearch.my@outlook.com>
to me

May 11, 2022

Dear Dr Widyawati,

Received with thanks.

Thanks & Regards

Vivian New

Editor

Food Research

From: Paini Sri Widyawati <paini@ukwms.ac.id>

Sent: Tuesdav. 10 May. 2022 10:02 PM



food research kecombrang



- 99+ Compose
- Mail
- Inbox 1,441
- Starred
- Snoozed
- Sent
- Drafts 2
- More

Labels

F **Food Research** <foodresearch.my@outlook.com> May 15, 2022,
to me

Dear Dr Widyawati,

Please refer to the attachment for the galley proof of your manuscript FR-2021-333 entitled 'The effect of rotar drying characteristic and antioxidant activity of *Etilingera elatior* Jack' . Please check the content of the galley for mistakes, please comment and highlight in the PDF itself and revert to us within two (2) days of receipt. Once version, your manuscript will be published online for early viewing.

Please see the attachment for the invoice INV22129. We hope that you can make the payment as soon as possible for us to complete the publication of your manuscript. The manuscript information e.g. volume, issue, page number provided once we have received the payment.

Thanks & Regards,
Vivian New
Editor
Food Research

From: Food Research <foodresearch.my@outlook.com>
Sent: Friday, 13 May, 2022 4:12 PM



food research kecombrang



- 99+
- Mail
- Compose
- Inbox 1,441
- Starred
- Snoozed
- Sent
- Drafts 2
- More
- Chat
- Spaces
- Meet

Labels

Dear Dr. Vivian New We agree with the manuscript from the galley proof and I have paid the publication fee. I attach proof of t



Food Research <foodresearch.my@outlook.com>
to me

May 18, 2022

Dear Dr Widyawati,

Thank you very much for the payment. I'll notify you of the article's publication soon.

Thanks & Regards
Vivian New
Editor
Food Research

From: Paini Sri Widyawati <paini@ukwms.ac.id>

Sent: Wednesday, 18 May, 2022 4:36 PM

- You are welcome.
- Thanks a lot.
- Noted with thanks.