

# Membranes and deoxyribonucleic acid of hippocampal neurons damage due to lowdensity polyethylene microplastics in blood of Wistar rats

*by* Yudhiakuari Sincihu

---

**Submission date:** 27-Mar-2024 11:10AM (UTC+0700)

**Submission ID:** 2332450882

**File name:** 8-Membranes\_and\_deoxyribonucleic.pdf (894.65K)

**Word count:** 5409

**Character count:** 29424

**How to Cite:**

Sincihu, Y., Sudiana, I. K., Kusumastuti, K., Keman, S., Sulistyorini, L., Lusno, M. F. D., Elias, S. M., & Hasina, S. N. (2022). Membranes and deoxyribonucleic acid of hippocampal neurons damage due to low-density polyethylene microplastics in blood of Wistar rats. *International Journal of Health Sciences*, 6(S7), 3490–3503. <https://doi.org/10.53730/ijhs.v6nS7.12621>

## **Membranes and deoxyribonucleic acid of hippocampal neurons damage due to low-density polyethylene microplastics in blood of Wistar rats**

**Yudhiakuari Sincihu**

Faculty of Medicine, Widya Mandala Surabaya Catholic University, Surabaya, Indonesia and Doctoral Program of Public Health, Universitas Airlangga, Surabaya, Indonesia

**I Ketut Sudiana**

Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

**Kurnia Kusumastuti**

Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

**Soedjajadi Keman**

Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

Corresponding author email : [soedja\\_keman@fkm.unair.ac.id](mailto:soedja_keman@fkm.unair.ac.id)

**Lilis Sulistyorini**

Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

**Muhammad Farid Dimjati Lusno**

Doctoral Program of Public Health, Universitas Airlangga, Surabaya, Indonesia

Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

**Saliza Mohd Elias**

Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, Serdang, Malaysia

**Siti Nur Hasina**

Universitas Nahdlatul Ulama Surabaya, Surabaya, Indonesia

**Abstract**---Microplastic particles in the blood can cause damage to organs such as the brain. This study aimed to analyze the effect of microplastic particles in the blood on membrane damage (expression of malondialdehyde metabolites) and deoxyribonucleic acid damage (expression of 8-hydroxy-2'-deoxyguanosine metabolites) in hippocampus

neurons of Wistar rats. Methods: Forty-two Wistar rats were used and equally divided into six groups. The study groups X1, X2, X3, X4 and X5 was given 0.0375mg, 0.075mg, 0.15mg, 0.3mg, and 0.6mg of low-density polyethylene microplastic powder mixed with 2cc distilled water respectively, while the control group only given distilled water. Microplastic administration was carried out for 90 days. Results: Microplastic particles were found in the blood of Wistar rats. The level of microplastics particle was higher along with the higher exposure dose. The mean expression of malondialdehyde and 8-hydroxy-2'-deoxyguanosine metabolites in the hippocampal neurons in CA1 and CA3 areas were significantly increased with higher exposure doses (Kruskal-Wallis test  $p < 0.01$ ). The Spearman's correlation showed a strong relationship between the levels of microplastic particles in blood and the expression of metabolites malondialdehyde and 8-hydroxy-2'-deoxyguanosine (all  $p < 0.01$ ). Microplastic in the blood of Wistar rats has increased expression of malondialdehyde and 8-hydroxy-2'-deoxyguanosine metabolites in hippocampal neurons.

**Keywords**--8-hydroxy-2'-deoxyguanosine, Environmental pollution, Hippocampal neurons damage, Malondialdehyde, Microplastics.

## Introduction

Microplastics (MPs <5mm) have become pollutants in the human food chain (Cox KD et al, 2019; Jiang S et al, 2021). This is evidenced by the discovery of plastic fibre particles accumulating in the organs of living things (Karbalaei S, Hanachi P, Walker TR, Cole M, 2018). Human faeces contain MPs polymeric particles of polyethylene, polystyrene, polyvinyl chloride and polyethylene terephthalate were reported (Schwabl P et al, 2019; Yan Z et al, 2019). The entry route for MPs in humans is not only through contaminated food, but also through inhalation of polluted air and skin contact with MPs. MPs with a diameter of ~20µm will circulate in the blood 1-2 days post ingestion for distribution to all organs and tissues of the body (Wright SL, Kelly FJ, 2017). MPs with a diameter of ≤10µm can penetrate the blood brain barrier by passive transport because they have a fat-soluble composition of hydrophobic organic contaminants [7]. Whole component of MPs will trigger biological responses, such as inflammation, genotoxicity, oxidative stress, hypersensitivity, and unwanted immune responses that end in cell death (Karbalaei S et al, 2018; Companale C et al, 2020; Hwang J et al, 2019).

In body tissue, MPs are persistent, bio-accumulative, and toxic due to the free radicals formed (Deng Y et al, 2017; Shengchen W et al, 2021). In the brain, it causes neurodegeneration of neurons. The dose, particle size and length of exposure time determine the effects of brain damage and disease (Dehaut A et al, 2016). Normally, free radicals are present in every living thing. However, high levels cause cellular damage via reactive oxygen species (ROS) formation (Pomatto LCD et al, 2018). Damage occurs because cellular antioxidant levels are lower than oxidant levels (Trestrail C et al, 2020). Various studies have mentioned low antioxidant levels in animal exposed to MPs (Karbalaei S, 2018; Companale C, 2020; Trestrail C et al, 2020). The number of neurons damage

causes disruption of cognitive abilities as the main function of the brain. The hippocampus is one of the anatomical structures that play a role in cognitive function. There are 4 areas called the Cornu Ammonis (CA), namely CA1, CA2, CA3 and CA4. The learning function is played by the CA1 area and the learning function in the CA3 area [14,15] (Cao Y et al, 2016; Konen LM et al, 2020).

Cellular death can be through membrane or deoxyribonucleic acid of neurons damage. Free radicals in the form of hydroxyl radical (OH\*) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) formed from the Haber-Weiss and Fenton reaction play a role in damaging membrane and deoxyribonucleic acid neurons (Xu TANZ et al, 2019; Ribeiro IML et al, 2020). Cellular metabolite of phospholipid membrane damage is malondialdehyde (MDA), while deoxyribonucleic acid damage is 8-oxo-7,8-dihydro-2-deoxyguanosine (8-OHdG) (Song P et al, 2021). These cellular metabolites can be observed under a microscope by immunohistochemical staining. The presence of these cellular metabolites indicates neurons damage. Another important point in this study is about an adequate dose of MPs that can cause neurons damage in the hippocampus

Wistar rats were used as a model to assess the presence of cell damage in hippocampal neurons due to the administration of toxic substances (microplastics). Wistar rats are genetically homologous to humans so the results of this study can be used as the basis for further research on higher mammalian orders. The hippocampus of Wistar rats is structurally different from humans but is functionally similar (Golcaves EN et al, 2018).

Therefore, the aim of this study was to analyze low-density polyethylene microplastic particles in the blood that cause membrane and deoxyribonucleic acid damage in hippocampal neurons of Wistar rats based on observed expression of malondialdehyde (MDA) and 8-hydroxy-2'-deoxyguanosine (8-OHdG) metabolites in hippocampal neurons.

## Method

### Experimental research design

This was an experimental laboratory study with a post-test only control group design to examine the effects of blood microplastic particles on the hippocampus neurons. The research was carried out at the Biomedical Laboratory Widya Mandala Surabaya Catholic University and Research Centre Universitas Airlangga, Indonesia. The research was conducted from October 2021 until March 2022. Lemeshow formula at  $\alpha$  0.05 and  $\beta$  0.05, obtained 7 Wistar rats for each group. Forty-two (N =42) Wistar rats were divided into six groups by random allocation. Wistar rats were obtained from Farma Veterinary Center Surabaya and had a certificate of eligibility as experimental animals. Wistar rats were male, 12 weeks old, weighing 160-175 grams, fed and watered at libitum, clean care environment, stable air circulation, room temperature 18-26°C, and humidity 40-70%. Each Wistar rat in the experimental group was given of dry low-density polyethylene microplastics powder per day dissolved in 2cc of distilled water through an oral probe. Study group 1 (X1) was given 0.0375mg, study group 2 (X2) was given 0.075mg, study group 3 (X3) was given 0.15mg, study group 4 (X4) was given

0.3mg, and study group 5 (X5) given 0.6 mg of microplastic per day. While the control group (K) only given distilled water without microplastics. After 90 days, there was no drop out, Wistar rats were anesthetized and 1cc of blood was taken through cardiac puncture to measure the number of microplastic particles in the blood. After the blood collection procedure was completed, Wistar rats were euthanized with cervical dislocation technique until vital signs of life such as heart rate were not detected and pupillary reflexes were negative. Wistar rats were dissected through the head and removed the brain. After making histopathological preparations in the hippocampus area, anti-MDA and 8-OHdG immunohistochemical staining were made. Results of the microscopic examination were statistically analyzed to assess the effect of microplastic administration on the levels of microplastic particles in the blood of Wistar rats. After that, we analyzed the correlation between the level of microplastic particles in blood and the expression of malondialdehyde (MDA) and 8-hydroxy-2'-deoxyguanosine (8-OHdG) metabolites in hippocampal neurons. Adequate doses of microplastics in the blood that can damage hippocampal neurons were also be analyzed using statistical tests.

#### Microplastics processing

Microplastic (MPs) particles were made from Low-Density Polyethylene (plastic wrap) which was fragmented using Miller FCT-Z100 (Fomac, Indonesia). Milling until the size of MPs particle is fine. The plastic powder filtered with an 800mesh sieve (Anping Tianhao Wire Mesh Products Co., Ltd, China) to produce particles measuring <20µm.

#### Assessment of blood microplastics levels

The procedure for examining blood levels of MPs refers to the method previously described by Monteleone et al (2019) namely destruction of the blood, isolation of microplastic particles, and examination of blood microplastic particles level. Wistar rat blood samples were prepared at Biomedical Laboratory of Widya Mandala Surabaya Catholic University. The level of MPs particles in blood was calculated by clinical pathologist at Neuron Clinic Laboratory in Surabaya using binocular light microscope.

#### Immunohistochemical staining procedures

Examination of malondialdehyde (MDA) metabolites expression in hippocampal neurons following the manufacturer's protocol for JAI-MMD-030N anti-Malondialdehyde [MDA] mAb (1F83) (AdipoGen Life Sciences, Inc® San Diego). This protocol was used and described previously by Li et al, 2013. Examination of the expression of 8-hydroxy-2'-deoxyguanosine (8-OHdG) metabolites in hippocampal neurons followed the manufacturer's protocol for 8-OHdG monoclonal antibody (Immundiagnostik AG® Germany). This protocol was used and previously described by Topal et al 2017. This immunohistochemical preparation procedure was carried out at the Anatomical Pathology Laboratory, Universitas Airlangga.

#### Assessment of immunohistochemical slide



Immunohistochemical assessment of hippocampal neurons of Wistar rats was performed by an anatomic pathologist using binocular light microscope at 40x and 400x magnification. Calculations of the mean number of hippocampal neurons expressing malondialdehyde (MDA) and 8-hydroxy-2'-deoxyguanosine (8-OHdG) metabolites was performed in Cornu Ammonis 1 (CA1) and Cornu Ammonis 3 (CA3) areas in each of five fields of view. The expression of metabolites MDA and 8-OHdG showed dark brown cells (red arrows on figure) due to the use of the DAB chromogen. The brown precipitate in these cells will not dissolve in alcohol and xylene during the rehydration process.

#### Research data analysis

The results in this study were presented in the form of a graph of the mean ( $\bar{x}$ )  $\pm$  standard deviation (SD) and immunohistochemical microscopic images of the expression of malondialdehyde (MDA) and 8-hydroxy-2'-deoxyguanosine (8-OHdG) metabolites in hippocampal neurons. Normality test was performed using the Shapiro-Wilk test. The test showed a group with p value  $<0.05$  which indicate that the research data was not normally distributed. Therefore, the statistical analysis was performed by nonparametric test. Kruskal-Wallis test was used to determine the significance of differences in the number of MPs particle in blood, expression of malondialdehyde neurons, and expression of 8-hydroxy-2'-deoxyguanosine neurons between groups at 95% confidence intervals. The correlation between the number of MPs particle in the blood and the expression of malondialdehyde (MDA) and 8-hydroxy-2'-deoxyguanosine (8-OHdG) metabolites in hippocampal neurons used Spearman's Rho test. Multiple comparison tests were used to determine the adequate dose that caused damage to membrane and deoxyribonucleic acid of hippocampal neurons. The Mann-Whitney U test was used as a multiple comparison.

#### Ethics statement

Research ethics approval was obtained from the Health Research Ethics Committee (HREC) Widya Mandala Surabaya Catholic University with a certificate number 209/WM12/KEPK/DOSEN/T/2021.

#### Discussion




Figure 1. Mean ± SD levels of blood microplastic particles




Figure 2. Mean ± SD expression of malondialdehyde metabolites in hippocampal neurons




Figure 3. Histopathological images of hippocampal neuron membrane damage in Wistar rats, stained with immunohistochemical. Description: a) control group with CA1 and CA3 area, 40x magnification; b-c) MDA expression in CA1 and CA3 area, unexposed to MPs, microscope 400x magnification; d) study group with CA1 and CA3 area, 40x magnification; e-f) MDA expression in CA1 and CA3 area, exposed to MPs, microscope 400x magnification; red arrows is hippocampal neurons expressing MDA; green arrows is normal cell neuron.





Figure 4. Mean ± SD expression of 8-hydroxy-2'-deoxyguanosine metabolites in hippocampal neurons





group with CA1 and CA3 area, 40x magnification; b-c) 8-OHdG expression in CA1 and CA3 area, unexposed to MPs, microscope 400x magnification; d) study group with CA1 and CA3 area, 40x magnification; e-f) 8-OHdG expression in CA1 and CA3 area, exposed to MPs, microscope 400x magnification; red arrows is hippocampal neurons expressing 8-OHdG; green arrows is normal cell neuron.

Table 1  
Comparison of MPs particle in blood, expression of MDA and 8-OHdG metabolite in hippocampal neurons between groups

Variable	p value of Kruskal-Wallis statistical test
MPs particle in blood	<0.01
MDA metabolite expression at CA1	<0.01
MDA metabolite expression at CA3	<0.01
8-OHdG metabolite expression at CA1	<0.01
8-OHdG metabolite expression at CA3	<0.01

Table 2  
The correlations between number of MPs particle in blood with the expression of MDA and 8-OHdG metabolites

Parameter	Expression of MDA Metabolites	Expression of 8-OHdG Metabolites
Correlation Coefficient	.755**	.842**

p value of Spearman's rho <0.01 <0.01

Table 3  
Multiple comparison analysis between groups on MDA and 8-OHdG metabolites expression variable in hippocampal neurons

Group	p value of Mann-Whitney U test on MDA metabolites expression						p value of Mann-Whitney U test on 8-OHdG metabolites expression					
	K	X1	X2	X3	X4	X5	K	X1	X2	X3	X4	X5
K	-	1.00	0.15	0.00	0.00	0.00	-	0.05	0.00	0.00	0.00	0.00
		0	5	1	1	1		9	3	4	2	2
X1	1.00	-	0.15	0.00	0.00	0.00	0.05	-	0.04	0.01	0.00	0.00
		0	5	1	1	1		9	0	7	3	3
X2	0.15	0.15	-	0.02	0.00	0.00	0.00	0.04	-	0.22	0.02	0.00
		5	5	0	6	2		3	0	2	1	4
X3	0.00	0.00	0.02	-	0.33	0.01	0.00	0.01	0.22	-	0.14	0.01
		1	1	0		3		4	7	2	0	7
X4	0.00	0.00	0.00	0.33	-	0.05	0.00	0.00	0.02	0.14	-	0.15
		1	1	6	3	4		2	3	1	0	8
X5	0.00	0.00	0.00	0.01	0.05	-	0.00	0.00	0.00	0.01	0.15	-
		1	1	2	0	4		2	3	4	7	8

We all already know that plastic pollutants have polluted sea water, fresh water and soil ecosystems globally (Walker TR, 2019). Currently, small size of plastic particles was reported to contaminate human food and drink (Cox KD et al, 2019; Jiang S et al, 2021). One of them is low-density polyethylene (LDPE). This will be a serious problem in the future, especially in the health sector (Jiang S et al, 2021; Jambeck JR et al, 2015; Miraj SS et al, 202). In this study, LDPE was formed into microplastic (MPs) particles. These microplastics are <20µm in diameter with sharp and irregular edges. The shape of MPs determines the biological response (Wright SL, Kelly FJ, 2017).

The findings of this study indicate that MPs particles have been found in blood of Wistar rats given orally MPs for 90 days. The number of MPs particles found in the blood was higher in the Wistar rats group given higher doses of MPs (Figure 1). Previous research stated that MPs particles with diameter ≤20µm in the digestive tract can enter the blood circulation system through endocytosis mechanism, then can be distributed to various organs in the body (Lei L et al, 2018; Wu B et al, 2019; wang YL et al, 2019). At sizes <10 µm, MPs particles can pass through cell membranes and are found in the placenta, muscle, and brain (Campanale C et al, 2020). Administration of higher MPs doses will cause more contact between MPs particles and the surface of microfold cells at the tips of the intestinal villi, thereby increasing the potential for MPs particle endocytosis. These results are similar to the findings of (Hamed et al, 2019), in Tilapia was given 1mg/L, 10mg/L, and 100mg/L of MPs for 15 days. They stated that the number of MPs particle per gram of fish tissue increased along with higher exposure doses. In their study, it was not known how many doses of MPs particle were consumed by Tilapia, but our study explained between the MPs exposure dose and the number of MPs particle found in the blood.

The current study found that the MPs particles level in blood is significantly and positively correlated with the neuron damage in the Wistar rats. The findings of this study indicate the occurrence of membrane (expression of malondialdehyde metabolite) and deoxyribonucleic acid (expression of 8-hydroxy-2'-deoxyguanosine metabolite) damage in hippocampal neurons of Wistar rats due to low-density polyethylene microplastics. This damage can occur because MPs particles in the blood circulation or accumulated in the organ tissues cannot be destroyed by phagocytic cells. Even so, cellular defence continue to destroy them through oxygen-dependent and oxygen-independent mechanisms. Oxygen-dependent mechanisms such as oxidative stress, while oxygen-independent mechanisms such as the role of inflammatory mediators (Othman AR et al, 2021; Li Z et al 2020). The final response of this mechanism will induce cell damage (Hwang J et al, 2019; Trestrail C et al, 2020).

Malondialdehyde (MDA) as an indirect index of oxidative damage caused by lipid peroxidation in hippocampal neuronal membranes. Deoxyguanosine (dG) is one of the bases that compose the deoxyribonucleic acid of hippocampal neurons, which when oxidized will become 8-hydroxy-2'-deoxyguanosine (8-OHdG) (Song P, 2021). Initially, microplastics consumed will increase intracellular antioxidant enzymes such as superoxide dismutase and catalase. However, if microplastics continue to enter the body, these enzymes will decrease because they are unable to reduce the free radicals that are formed. This has been proven by many studies with exposure to MPs (Trestrail C et al, 2020; Li Z et al, 2020; Sokmen TO et al, 2020). In this condition, cell damage begins to occur, such as an increase in the expression of MDA and 8-OHdG metabolites as an indicator of free radical reactions in experimental animals (Shengchen W et al, 2021).

This finding is in line with the study by Li et al, 2020, in Wistar rats fed 5mg/L of polystyrene MPs for 90 days ( $\approx$  0.15mg of dry MPs particles per day) showed that myocardium damage was triggered by oxidative stress. Study by Sokmen et al (2020), stated that similar to the findings of this study, oxidative stress occurred in zebrafish embryo brain tissue exposed to 20nm diameter polystyrene MPs for 120 hours and increased expression of the 8-OHdG metabolites. Study by Cheng et al (2020), which used LDPE MPs with a diameter of 550-1000 $\mu$ m for 28 days showed an increase in oxidative stress which led to increased expression of MDA and 8-OHdG metabolites genes in *Eisenia Fetida*. Similarly, in the study by Chen et al (2020), in *Eisenia Fetida* exposed to LDPE MPs with a diameter <400 $\mu$ m at a minimum dose of 1 g/kg dry powder showed an increase in MDA metabolites in intestinal tissue.

The toxicity of LDPE MPs comes from plastic particles that cannot be destroyed through cellular defense mechanisms. In addition, it also comes from toxic compounds contained in these particles due to the production process or absorbed when low-density polyethylene is in nature (Wright SL, Kelly FJ, 2017). The toxic compounds are still found in the form of fragmented plastics (Wright SL, Kelly FJ, 2017; CampanaleC et al, 2020; Brennecke D, 2016).]. When these plastic particles or compounds enter the body, it will trigger the breaking of covalent bonds of atom into unpaired electrons (Trestrail C, 2020). Free radicals are molecules or atoms that have one or more unpaired electrons in their outer



shell so they are highly reactive and radical. This reaction will occur continuously because the MPs particles cannot be destroyed by the body's defense mechanisms. If the concentration of cellular antioxidants is insufficient to neutralize these free radicals, it will cause oxidative stress. Oxidative stress will continue with damage to cell structure. Continuous damage will cause a disease (Pomatto LCD et al 2018; Trestrail C, 2020). In the brain, oxidative stress causes neurodegeneration through neurons damage (Xu TANZ et al, 2019; Ribeiro IML, 2020; Song P et al, 2021). Brain neurodegeneration is a pathophysiology of decreased brain function. One of them is cognitive function played by the hippocampus.

Wistar rats hippocampal neurons damage was seen significantly when given MPs at dose of 0.075mg per day for 90 days (equivalent to 45 MPs particles in milliliters of blood). In humans weighing  $\pm$  70 kilograms, this MP exposure dose is equivalent to 3.47mg. This means that if humans consume as much as 3.47mg of MP particles every day, there will be significant damage to their hippocampal neurons.

This study has proven that MPs in the blood cause damage to the hippocampal neurons as seen from the expression of MDA and 8-OHdG metabolites in hippocampal neurons. The results of this study cannot be generalized to other neurons in the brain. In addition, this study has limitations such as not examining the presence of MPs particles in the brain of Wistar rats, the size of the MPs particle used varies up to 20 $\mu$ m in diameter, and does not examine the concentration of other toxic compounds contained in the blood of Wistar rats due to MPs exposure. As our recommendation, in other studies it is necessary to measure the levels of MPs particle in the brain (especially the hippocampus) and identify various toxic substances in the blood due to MPs exposure. This substance also has a toxic effect on brain neurons such as phthalates, bisphenol-A, polybrominated diphenyl ether, dichloro-diphenyl-dichloroethylene, dichloro-diphenyl-trichloroethane, polycyclic aromatic hydrocarbons, and polychlorinated biphenyls.

### **Conclusion**

Administration of low-density polyethylene MPs <20 $\mu$ m in diameter for 90 days caused the presence of microplastic particles in the blood of Wistar rats. The higher number of MPs particles in the blood causes an increase in the expression of MDA and 8-OHdG metabolites in the hippocampal neurons. These expressions indicate the hippocampal neurons damage in membranes and deoxyribonucleic acid.

### **Acknowledgments**

This research was financially supported by Universitas Airlangga as an overseas partner collaborative mandate research grant. Registered with grant number 777/UN3.15/PT/2021. We also express our deepest gratitude to head and all staff of the Laboratory of Pathology Anatomical Universitas Airlangga, Niluh Suwasanti as a head of Clinical Pathology at Neuron Clinic Laboratory, and also



all staff of Biomedical Laboratory of Widya Mandala Surabaya Catholic University as places for conducting research.

## References

- Brennecke, D., Duarte, B., Paiva, F., Caçador, I., & Canning-clode, J. (2016). Microplastics as vector for heavy metal contamination from the marine environment. *Estuarine, Coastal and Shelf Science*, 178, 189–195. <https://doi.org/10.1016/j.ecss.2015.12.003>
- Campanale, C., Massarelli, C., Savino, I., Locaputo, V., & Uricchio, V. F. (2020). A detailed review study on potential effects of microplastics and additives of concern on human health. *International Journal of Environmental Research and Public Health*, 17(4). <https://doi.org/10.3390/ijerph17041212>
- Cao, Y., Gou, Z., Du, Y., Fan, Y., Liang, L., Yan, Y., Lin, P., Jin, M., & Du, Y. (2016). Glutamatergic and central cholinergic dysfunction in the CA1, CA2 and CA3 fields on spatial learning and memory in chronic cerebral ischemia — Induced vascular dementia of Rats. *Neuroscience Letters*, 620, 169–176. <https://doi.org/10.1016/j.neulet.2016.03.039>
- Chen, Y., Liu, X., Leng, Y., & Wang, J. (2020). Defense responses in earthworms (*Eisenia fetida*) exposed to low-density polyethylene microplastics in soils. *Ecotoxicology and Environmental Safety*, 187(109788), 1–6. <https://doi.org/10.1016/j.ecoenv.2019.109788>
- Cheng, Y., Zhu, L., Song, W., Jiang, C., Li, B., Du, Z., Wang, J., Wang, J., Li, D., & Zhang, K. (2020). Combined effects of mulch film-derived microplastics and atrazine on oxidative stress and gene expression in earthworm (*Eisenia fetida*). *Science of the Total Environment*, 746(141280), 1–11. <https://doi.org/10.1016/j.scitotenv.2020.141280>
- Cox, K. D., Covernton, G. A., Davies, H. L., Dower, J. F., Juanes, F., & Dudas, S. E. (2019). Human consumption of microplastics [Research-article]. *Environmental Science & Technology*, 53, 7068–7074. <https://doi.org/10.1021/acs.est.9b01517>
- Dehaut, A., Cassone, A. L., Frère, L., Hermabessiere, L., Himber, C., Rinnert, E., Rivière, G., Lambert, C., Soudant, P., Huvet, A., Duflos, G., & Paul-Pont, I. (2016). Microplastics in Seafood: Benchmark protocol for their extraction and characterization. *Environmental Pollution*, 215, 223–233. <https://doi.org/10.1016/j.envpol.2016.05.018>
- Deng, Y., Zhang, Y., Lemos, B., & Ren, H. (2017). Tissue accumulation of microplastics in mice and biomarker responses suggest widespread health risks of exposure. *Scientific Reports*, 7(March), 1–10. <https://doi.org/10.1038/srep46687>
- Gonçalves, E. N., Faustino-rocha, A. I., Seixas, F., Colaço, B., Ferreira, R., Fardilha, M., Oliveira, P. A., & Ginja, M. (2018). Modelling human prostate cancer: Rat models. *Life Sciences*, 203, 210–224. <https://doi.org/10.1016/j.lfs.2018.04.014>
- Hamed, M., Soliman, H. A. M., Osman, A. G. M., & Sayed, A. E. H. (2019). Assessment the effect of exposure to microplastics in Nile Tilapia (*Oreochromis niloticus*) early juvenile: I. blood biomarkers. *Chemosphere*, 228, 345–350. <https://doi.org/10.1016/j.chemosphere.2019.04.153>
- Hwang, J., Choi, D., Han, S., Choi, J., & Hong, J. (2019). An assessment of the toxicity of polypropylene microplastics in human derived cells. *Science of the*

- Total Environment*, 684, 657–669.  
<https://doi.org/10.1016/j.scitotenv.2019.05.071>
- Jambeck, J. R., Geyer, R., Wilcox, C., Siegler, T. R., Perryman, M., Andrady, A., Narayan, R., & Law, K. L. (2015). Plastic waste inputs from land into the ocean. *Science Journal*, 347(6223), 768–772.
- Jiang, S., Wang, F., Li, Q., Sun, H., Wang, H., & Yao, Z. (2021). Environment and food safety: a novel integrative review. *Environmental Science and Pollution Research*, 28, 54511–54530.
- Kantari, C., Ribeil, M. P., & Sarsat, éronique W. (2008). The role of neutrophils and monocytes in innate immunity. *Trends in Innate Immunity*, 15, 118–146.  
<https://doi.org/10.1159/000136335>
- Karbalaei, S., Hanachi, P., Walker, T. R., & Cole, M. (2018). Occurrence, sources, human health impacts and mitigation of microplastic pollution. *Environmental Science and Pollution Research*, 25(36), 36046–36063.  
<https://doi.org/10.1007/s11356-018-3508-7>
- Konen, L. M., Wright, A. L., Royle, G. A., Morris, G. P., Lau, B. K., Seow, P. W., Zinn, R., Milham, L. T., Vaughan, C. W., & Vissel, B. (2020). A new Mouse line with reduced GluA2 Q / R site RNA editing exhibits loss of dendritic spines , hippocampal CA1-neuron loss , learning and memory impairments and NMDA receptor-independent seizure vulnerability. *Molecular Brain*, 13(27), 1–19.
- Lei, L., Liu, M., Song, Y., Lu, S., Hu, J., Cao, C., Xie, B., Shi, H., & He, D. (2018). Polystyrene (nano) microplastics cause size-dependent neurotoxicity, oxidative damage and other adverse effects in *Caenorhabditis elegans*. *Environmental Science: Nano*, 5(8), 2009–2020.  
<https://doi.org/10.1039/c8en00412a>
- Li, G., Feng, Y., Cheng, T. S., Yin, J., & Zhang, C. (2013). Edaravone, a novel free radical scavenger, prevents steroid-induced osteonecrosis in Rabbits. *Rheumatology*, 52, 438–447. <https://doi.org/10.1093/rheumatology/kes313>
- Li, Z., Zhu, S., Liu, Q., Wei, J., Jin, Y., Wang, X., & Zhang, L. (2020). Polystyrene microplastics cause cardiac fibrosis by activating Wnt/ $\beta$ -catenin signaling pathway and promoting cardiomyocyte apoptosis in rats. *Environmental Pollution*, 265, 115025. <https://doi.org/10.1016/j.envpol.2020.115025>
- Miraj, S. S., Parveen, N., & Zedan, H. S. (2021). Plastic microbeads: small yet mighty concerning. *International Journal of Environmental Health Research*, 31(7), 788–804. <https://doi.org/10.1080/09603123.2019.1689233>
- Monteleone, A., Schary, W., Fath, A., & Wenzel, F. (2019). Validation of an extraction method for microplastics from human materials. *Clinical Hemorheology and Microcirculation*, 73(1), 203–217.  
<https://doi.org/10.3233/CH-199209>
- Othman, A. R., Hasan, H. A., Muhamad, M. H., Ismail, N. 'Izzati, & Abdullah, S. R. S. (2021). Microbial degradation of microplastics by enzymatic processes : a review. *Environmental Chemistry Letters*, 19, 3057–3073.  
<https://doi.org/10.1007/s10311-021-01197-9>
- Pomatto, L. C. D., & Davies, K. J. A. (2018). Adaptive homeostasis and the free radical theory of ageing. *Free Radical Biology and Medicine*, 124(August), 420–430. <https://doi.org/10.1016/j.freeradbiomed.2018.06.016>
- Ribeiro, I. M. L., Pereira, W. L., Nogueira, L. B., Oliveira, L. A. M., Teixeira, R. R., & Nogueira, K. de O. P. C. (2020). Neuroprotective effect of isobenzofuranones on hydrogen peroxide-mediated redox imbalance in

- primary cultures of hippocampal neurons. *Brazilian Archives of Biology and Technology*, 63, 1–8.
- Schwabl, P., Koppel, S., Königshofer, P., Bucsecs, T., Trauner, M., Reiberger, T., & Liebmann, B. (2019). Detection of various microplastics in human stool. *Annals of Internal Medicine*, 171, 453–458. <https://doi.org/10.7326/M19-0618>
- Shengchen, W., Jing, L., Yujie, Y., Yue, W., & Shiwen, X. (2021). Polystyrene microplastics-induced ROS overproduction disrupts the skeletal muscle regeneration by converting myoblasts into adipocytes. *Journal of Hazardous Materials*, 417(April), 1–12. <https://doi.org/10.1016/j.jhazmat.2021.125962>
- Sökmen, T. Ö., Sulukan, E., Türkoğlu, M., Baran, A., Özkaraca, M., & Ceyhun, S. B. (2020). Neurotoxicology polystyrene nanoplastics (20 nm) are able to bioaccumulate and cause oxidative DNA damages in the brain tissue of zebrafish embryo (Daniorerio)\*. *Neurotoxicology*, 77, 51–59. <https://doi.org/10.1016/j.neuro.2019.12.010>
- Song, P., Yan, B., Lei, F., Qiu, Z., Zhang, C., Wu, Y., Chen, S., Yang, X., Shen, D., & Ma, P. (2021). Continuous artificial light at night exacerbates diisononyl phthalate-induced learning and memory impairment in Mice: Toxicological evidence. *Food and Chemical Toxicology*, 151(May), 1–9.
- Suryasa, I. W., Rodríguez-Gámez, M., & Koldoris, T. (2022). Post-pandemic health and its sustainability: Educational situation. *International Journal of Health Sciences*, 6(1), i-v. <https://doi.org/10.53730/ijhs.v6n1.5949>
- Topal, A., Alak, G., Ozkaraca, M., Kokturk, M., Atamanalp, M., Alak, G., Ozkaraca, M., Kokturk, M., & Atamanalp, M. (2017). Neurotoxic responses in brain tissues of rainbow trout exposed to imidacloprid pesticide: Assessment of 8-hydroxy-2-deoxyguanosine activity, oxidative stress and acetylcholinesterase activity. *Chemosphere*, 175, 186–191. <https://doi.org/10.1016/j.chemosphere.2017.02.047>
- Trestrail, C., Nugegoda, D., & Shimeta, J. (2020). Invertebrate responses to microplastic ingestion: Reviewing the role of the antioxidant system. *Science of the Total Environment*, 734(138559), 1–18. <https://doi.org/10.1016/j.scitotenv.2020.138559>
- Walker, T. R. (2019). (Micro)plastics and the UN sustainable development goals. *ScienceDirect*, 30, 1–9. <https://doi.org/10.1016/j.cogsc.2021.100497>
- Wang, Y. L., Lee, Y. H., Chiu, I. J., Lin, Y. F., & Chiu, H. W. (2020). Potent impact of plastic nanomaterials and micromaterials on the food chain and human health. *International Journal of Molecular Sciences*, 21(5). <https://doi.org/10.3390/ijms21051727>
- Wright, S. L., & Kelly, F. J. (2017). Plastic and human health: a micro issue? *Environmental Science and Technology*, 51(12), 6634–6647. <https://doi.org/10.1021/acs.est.7b00423>
- Wu, B., Wu, X., Liu, S., Wang, Z., & Chen, L. (2019). Size-dependent effects of polystyrene microplastics on cytotoxicity and efflux pump inhibition in human Caco-2 cells. *Chemosphere*, 221, 333–341. <https://doi.org/10.1016/j.chemosphere.2019.01.056>
- Xu, T. A. N. Z., Shen, X. Y. A. N., Sun, L. L., & Chen, Y. A. L. I. (2019). Ginsenoside Rg1 protects against H<sub>2</sub>O<sub>2</sub> - induced neuronal damage due to inhibition of the NLRP1 inflammasome signalling pathway in hippocampal neurons in vitro. *International Journal of Molecular Medicine*, 43, 717–726. <https://doi.org/10.3892/ijmm.2018.4005>

Yan, Z., Zhao, H., Zhao, Y., Zhu, Q., Qiao, R., Ren, H., & Zhang, Y. (2019). An efficient method for extracting microplastics from feces of different species. *Journal of Hazardous Materials*, 121489(October), 1–39. <https://doi.org/10.1016/j.jhazmat.2019.121489>



# Membranes and deoxyribonucleic acid of hippocampal neurons damage due to lowdensity polyethylene microplastics in blood of Wistar rats

## ORIGINALITY REPORT

6%

SIMILARITY INDEX

5%

INTERNET SOURCES

6%

PUBLICATIONS

2%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.ijicc.net">www.ijicc.net</a> Internet Source	1%
2	Kun Niu, Jia-Lin Qin, Guo-Fang Lu, Jian Guo, John P. Williams, Jian-Xiong An. "Dexmedetomidine Reverses Postoperative Spatial Memory Deficit by Targeting Surf1 and Cytochrome c", Neuroscience, 2021 Publication	1%
3	<a href="http://avesis.aybu.edu.tr">avesis.aybu.edu.tr</a> Internet Source	1%
4	<a href="http://repository.unair.ac.id">repository.unair.ac.id</a> Internet Source	1%
5	"Handbook of Microplastics in the Environment", Springer Science and Business Media LLC, 2022 Publication	1%
6	Ni Putu Wulan Purnama Sari. "The Unmet Needs of Women Living with Cancer in Community Setting: A Professional	1%

# Concern", Indonesian Journal of Cancer, 2022

Publication

7

[ira.lib.polyu.edu.hk](http://ira.lib.polyu.edu.hk)

Internet Source

1 %

8

[link.springer.com](http://link.springer.com)

Internet Source

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On