

BAB 5

SIMPULAN

5.1. Simpulan

Dari data penelitian yang telah diinterpretasikan, dapat ditarik kesimpulan :

- Tablet likuisolid ibuprofen yang menggunakan gliserin sebagai pelarut *non volatile* dan PVP K-30 sebagai polimer dapat meningkatkan disolusi ibuprofen dibandingkan dengan tablet ibuprofen konvensional (non likuisolid).
- Penambahan PVP K-30 sebagai polimer hidrofilik dapat meningkatkan konstanta laju disolusi tablet likuisolid ibuprofen dengan gliserin sebagai pelarut *non volatile*. Formula B dan C mempunyai konstanta laju disolusi lebih besar dibandingkan formula A, sedangkan formula D yang mengandung konsentrasi PVP K-30 sebesar 15% dari *liquid medication* memiliki konstanta laju disolusi yang tidak berbeda bermakna dengan Formula A.

5.2. Alur Penelitian Selanjutnya

Dapat dilakukan penelitian tablet likuisolid dengan mencari formula optimum berdasarkan berbagai faktor seperti konsentrasi obat dalam pelarut, konsentrasi polimer, konsentrasi *carrier*, dan perbandingan antara *carrier* dan *coating*.

DAFTAR PUSTAKA

- Anonim, 1979, **Farmakope Indonesia**, ed. III, Departemen Kesehatan RI, Jakarta, 6-8.
- Anonim, 1995, **Farmakope Indonesia**, ed. IV, Departemen Kesehatan RI, Jakarta, 4, 166, 449-450, 488-489, 515, 683, 783-784, 999-1000.
- Anonim, 1997, **AHFS**, Drug Information AMERICAN Society of Health System Pharmacist, Inc., Bethesda, 1499-1504.
- Anonim, 2007^a, **US Pharmacopeia XXX**, US Pharmacopeial Convention Inc., Rockville.
- Anonim, 2007^b, **Informasi Spesialite Obat Indonesia**, vol. 42, Ikatan Sarjana Farmasi Indonesia, Jakarta, 233-249.
- Ansel, H. C., 2008, **Pengantar Bentuk Sediaan Farmasi**, ed. 4, terjemahan Farida Ibrahim, Universitas Indonesia, Jakarta, 57.
- Banakar, U.V., 1992, **Pharmaceutical Disolution Testing**, Marcel Dekker Inc., New York, 19-25.
- Banker, G.S. and N.R. Anderson, 1986, Tablet, in: **The Theory and Practice of Industrial Pharmacy: Tablet**, L. Lachman, H.A. Lieberman, and J.L. Kanig (Eds.), 3rd ed., Lea and Febiger, Philadelphia, 259, 299, 316 – 329.
- Banker, G.S. and C.T. Rhodes, 1996, **Modern Pharmaceutics**, 3rd ed., Marcel Dekker Inc., New York, 341 – 348.
- Davies, P., 2001, Oral Solid Dosage Forms, in: **Pharmaceutical Preformulation and Formulation: A Practical Guide from Candidate Drug Selection to Commercial Dosage Form**, M. Gibson (Ed.), vol. 199, 2nd ed., Informa Healthcare USA, Inc., New York, 373.
- Dibbern, H.W., R.M. Muller, and E. Wirbitzki, 2002, **UV and IR Spectra: Pharmaceutical Substances (UV and IR) and Pharmaceutical and Cosmetic Excipients (IR)**, Editio Cantor Verlag, Germany.

- Duchene, D., 1983, Tablet Disintegration, In: **Topic in Pharmaceutical Science**, P.D. Breimer, and P. Spesier (Eds.), Proceeding of The 43rd International Congress of Pharmaceutical, Amsterdam – New York – Oxford, 387 – 399.
- Fierse, E. F. and A. T. Hagen, 1986, Pre formulation, In: **The Theory and Practice of Industrial Pharmacy**, L. Lachman, H. A. Lieberman, and J. L. Kanig (Eds.), 3rd ed., Lea and Febiger, Philadelphia, 183-184.
- Green, J.M., 1996, A Practical Guide to Analytical Method Validition, **Analytical Chemistry**, 68, 305-309.
- Gubbi S. and R. Jarag, 2009, Liquisolid Technique for Enhancement of Dissolution Properties of Bromhexine Hydrochloride, **J. Pharm**, 2(2), 382 – 386.
- Hartono, 2004, **Statistik untuk Penelitian**, Lembaga Studi Filsafat, Kemasyarakatan, Kependidikan, dan Perempuan (LSFK₂P) bekerja sama dengan pustaka pelajar, Yogyakarta, 236.
- Javadzadeh, Y., M.R. Siahi-Shabdar, S. Asnaashari, and A. Nokhodchi, 2007^a, Liquisolid Technique as a Tool for Enhancement of Poorly Water-Soluble Drugs and Evaluation of Their Physicochemical Properties, **Acta Pharm**, 57, 99 – 109.
- Javadzadeh, Y., B. J. Navimipour, and A. Nokhodchi, 2007^b, Liquisolid Technique for Dissolution Rate Enhancement of a High Dose Water Insoluble Drug (Carbamazepine), **International Journal of Pharmaceutics**, 341, 26 – 34.
- Karmarkar, A.B., I.D.Gonjari, A.H. Hosmani, P.N. Dhabale, and S.B. Bhise, 2009^a, Dissolution Rate Enhancement of Fenofibrat using Liquisolid Tablet technique, **Latin American Journal of Pharmacy**, 28(2), 219.
- Karmarkar, A.B., I.D.Gonjari, A.H. Hosmani, P.N. Dhabale, and S.B. Bhise, 2009^b, Liquisolid Tablets: A Novel Approach for Drug Delivery, **International Journal of Health Research**, 2(1), 45-50.
- Khan, K.A., 1975, The Concept of Dissolution Efficiency. **J. Pharm**, 27(1), 48-49.

Kibbe, A.H. , 2000, **Handbook of Pharmaceutical Excipients**, 3rd ed., The Pharmaceutical Press, London, 102, 143, 305, 501, 555.

Langenbucher, F., 1972, Linearization of Dissolution Rate Curve by Weibull Distribution, **Journal of Pharmaceutical Sciences**, 24, 979-981.

Liao, C.C. and C.I. Jarowski, 2006, Dissolution rates of corticoid solutions dispersed on silicas, **Journal of Pharmaceutical Sciences**, 73(3), 401-403.

Martin, A., J. Swarbrick, and A. Cammarata, 1983, **Physical Pharmacy**, 3rd ed., Lea & Febiger, Philadelphia,845-850.

Ohwoavworhua, F.O., T.A.Adelakun, and A.O. Okhamafe, 2009, Processing Pharmaceutical Grade Microcrystalline Cellulose from Groundnut Husk: Extraction Methods and Characterization, **International Journal of Green Pharmacy**, 3(2), 97-104.

Parrott, E.L., 1971, **Pharmaceutical Technology Fundamental Pharmaceutics**, 3rd ed., Burgess Publishing Company, Minneapolis, 17-19, 82, 160-162.

Peck, G.E., G.J. Baley., V.E.McCurdy, and G.S.Banker, 1989, Tablet Formulation and Design, in: **Pharmaceutical Dosage Forms: Tablets**, L. Lachman, H.A. Lieberman, and J.B. Schwartz (Eds.), vol. 1, 2nd ed., Marcell Dekker Inc., New York,88-127.

Reynolds, J.E.F., 1982, **Martindale: The Extra Pharmacopoeia**, 28th ed., The Pharmaceutical Press, London, 349.

Rowe, R.C., P.J. Sheskey, and M. E. Quinn, 2009, **Handbook of Pharmaceuticals Excipient**, 6th ed., The Pharmaceutical Press, London, 283, 581

Shargel, L. and A.B.C. Yu, 1988, **Biofarmasetika dan Farmakokinetika Terapan**, terjemahan Fasich dan S. Syamsial, Universitas Airlangga, Surabaya, 6 – 18, 101, 167 – 199.

Shargel, L. and A. B. C. Yu, 1999, **Applied Biopharmaceutics and Pharmacokinetics**, 4th ed. McGraw – Hill. New York, 8, 132, 169-200.

Shervington, L.A. and A. Shervington, 1998, Guaifenesin, In: **Analytical Profiles of Drug Substances and Excipients**, H.G. Brittain (Ed.), vol. 25, Academic Press, London, 152.

Siregar, C. J. P., 1992, **Proses Validasi Manufaktur Sediaan Tablet**, Institut Teknologi Bandung, Bandung, 29-31.

Spireas, S., 2002, Liquisolid System and Methods of Preparation Same, **Pharmaceutical Research**, vol 9, 1-6.

Sutriyo, Rosmaladewi, dan F. H. Filosane, 2005, Pengaruh Polivinil Pirolidon Terhadap Laju Disolusi Furosemid dalam Sistem Dispersi Padat, **Majalah Ilmu Kefarmasian**, 2(1), 30 – 42.

Sweetman, S. C., 2009, **Martindale The Complete Drug Reference**, 36th ed., The Pharmaceutical Press, London, 64

Voigt, R., 1995, **Buku Pelajaran Teknologi Farmasi**, Terjemahan S. Noerono dan M. S. Reksohardiprojo, Gadjah Mada University Press, Yogyakarta, 163-210.

Wade, A. and P.S. Weller, 1994, **Handbook of Pharmaceutical Excipients**, 2nd ed., The Pharmaceutical Press, London, 84 – 86.

Wagner, J.G., 1971, **Biopharmaceutics and Relevant Pharmacokinetics**, 1st ed., Drug Intelligence Publications, Illinois, 64-110.

Wells, J.T., 1988, **Pharmaceutical Formulation: The Physicochemical Properties of Drug Substance**, Ellis Howard, Ltd., Chester, 209-211.

Wilmania, P. F., 1995, Analgesik – Antipiretik Analgesik Anti-Inflamasi Nonsteroid dan Obat Pirai, dalam: **Farmakologi dan Terapi**, Sulistia G. Ganiswarna (Ed.), edisi 4, Gaya Baru, Jakarta, 207-218.

Yadav, V.B. and A.V. Yadav, 2009, Liquisolid Granulation Technique for Tablet Manufacturing: an Overview, **Journal of Pharmacy Research** 2009, 2(4), 670-674.

Zainuddin, M., 2000, **Validasi Metode Analisis Kuantitatif Secara Spektrofotometris Ultra Ungu – Sinar Tampak (UV-VIS)**, Universitas Airlangga, Surabaya.